arXiv:1208.1077v1 [math.GT] 6 Aug 2012

THE EQUIVALENCE OF HEEGAARD FLOER HOMOLOGY AND EMBEDDED CONTACT HOMOLOGY VIA OPEN BOOK DECOMPOSITIONS II

VINCENT COLIN, PAOLO GHIGGINI, AND KO HONDA

ABSTRACT. This paper is the sequel to [CGH2] and is devoted to proving some of the technical parts of the HF=ECH isomorphism.

CONTENTS

1. Introduction 2
2. Gromov-Witten type computations 3
2.1. A brief review of relative Gromov-Witten invariants 3
2.2. First relative Gromov-Witten calculation 4
2.3. Second relative Gromov-Witten calculation 7
2.4. Third relative Gromov-Witten calculation 13
3. Homotopy of cobordisms I 20
3.1. Construction of the homotopy of cobordisms for $\Psi \circ \Phi$ 20
3.2. Holomorphic curves and moduli spaces 23
3.3. Proof of half of Theorem 1.0.1 32
3.4. Degeneration at $+\infty$ 37
3.5. Degeneration at $-\infty$ 56
3.6. Breaking in the middle 67
3.7. Degeneration at $+\infty$, part II 70
3.8. Degeneration at $-\infty$, part II 86
3.9. Breaking in the middle, part II 92
4. Homotopy of cobordisms II 93
4.1. Construction of the homotopy of cobordisms for $\Phi \circ \Psi$ 93
4.2. Holomorphic curves and moduli spaces 95
4.3. Proof of the other half of Theorem 1.0.1 99
4.4. Degeneration at $+\infty$ 100
4.5. Degeneration at $-\infty$ 102
4.6. Breaking in the middle 109

Date: This version: July 24, 2012.

2000 *Mathematics Subject Classification.* Primary 57M50; Secondary 53D10,53D40.

Key words and phrases. contact structure, Reeb dynamics, embedded contact homology, Heegaard Floer homology, open book decompositions.

VC supported by the Institut Universitaire de France, ANR Symplexe, ANR Floer Power, and ERC Geodycon. PG supported by ANR Floer Power and ANR TCGD. KH supported by NSF Grants DMS-0805352 and DMS-1105432.

5. Stabilization 112
5.1. The setup 113
5.2. Morse-Bott theory 114
5.3. A commutative diagram 116
5.4. An isomorphism 118
5.5. Proof of Theorem 1.0.2 121
References 123

1. INTRODUCTION

This paper is the sequel to [CGH2] and is devoted to proving some of the technical parts of the isomorphism between $\widehat{HF}(-M)$ and $\widehat{ECH}(M)$. The notation from [CGH2] carries over to this paper. References from [CGH2] will be written as "Section I.x" to mean "Section x" of [CGH2], for example.

In [CGH2] we defined the chain maps

$$\Phi : \widehat{CF}(S, \mathbf{a}, h(\mathbf{a})) \to PFC_{2g}(N),$$

$$\Psi : PFC_{2g}(N) \to \widehat{CF}(S, \mathbf{a}, h(\mathbf{a})).$$

In this paper we prove the following two results:

Theorem 1.0.1 (Quasi-isomorphism)**.** *The chain maps Φ and Ψ are quasi-isomorphisms, provided the monodromy map h does not have any elliptic periodic points of period $\leq 2g$ in $int(S)$.*

Theorem 1.0.1 is quite involved and takes up most of this paper. The condition on h is a technical assumption which simplifies gluing and one should be able to remove this condition with more work.

Theorem 1.0.2 (Stabilization)**.** $\widehat{ECH}(M) \simeq PFH_{2g}(N)$.

In view of Theorem I.2.5.6, Theorem 1.0.2 would immediately follow from showing that the chain maps

$$\mathfrak{I}_j : ECC_j(N, f_j\alpha) \to ECC_{j+1}(N, f_{j+1}\alpha),$$

defined in Section I.2.5.3, are quasi-isomorphisms for $j \geq 2g$. What we actually prove is slightly weaker, but sufficient: the similarly defined chain maps

$$ECC_{2j}(N, f_{2j}\alpha) \to ECC_{2j+2}(N, f_{2j+2}\alpha)$$

are quasi-isomorphisms. Theorem 1.0.2 is proved in Section 5.

Organization of this paper. In Sections 3 and 4 we prove the chain homotopy between the chain maps $\Psi \circ \Phi$ and id, as well as the chain homotopy between the chain maps $\Phi \circ \Psi$ and id. The necessary Gromov-Witten type calculations are carried out in Section 2. Finally, Section 5 is devoted to proving Theorem 1.0.2.

2. GROMOV-WITTEN TYPE COMPUTATIONS

This section is devoted to Gromov-Witten type calculations which are used in the proof of Theorem 1.0.1.

After a brief review of relative Gromov-Witten invariants in Section 2.1, we treat a slightly simpler model situation in Section 2.2. We then tackle the specific situations of interest in this paper in Sections 2.3 and 2.4.

2.1. **A brief review of relative Gromov-Witten invariants.**

We now briefly review relative Gromov-Witten invariants, following Ionel-Parker [IP1]. The reader is referred to Ionel-Parker [IP1], as well as to Li-Ruan [LR] and McDuff [M2], for more complete discussions of relative Gromov-Witten invariants.

Let (X,ω) be a compact symplectic manifold and J a compatible almost complex structure on (X,ω).

Let $\overline{\mathcal{U}}_{g,n} \to \overline{M}_{g,n}$ be the universal curve over the Deligne-Mumford moduli space of genus g curves with n marked points, with a fixed embedding $\overline{\mathcal{U}}_{g,n} \subset \mathbb{CP}^N$. Then let ν be a section of the bundle

$$\mathrm{Hom}^{0,1}(\pi_2^* T\mathbb{CP}^N, \pi_1^* TX) \to X \times \mathbb{CP}^N$$

of $\mathbb{C}$-anti-linear maps, where π_1 and π_2 are projections onto the first and second factors of $X \times \mathbb{CP}^N$.

Let $(F,j,\mathbf{x})$, where $\mathbf{x}=(x_1,\dots,x_n)$, be a marked nodal Riemann surface, let $st(F,j,\mathbf{x})$ be the stable Riemann surface obtained by collapsing the unstable components to points and let $\phi_0 : st(F,j,\mathbf{x}) \to \overline{\mathcal{U}}_{g,n}$ be a biholomorphism onto a fiber of $\overline{\mathcal{U}}_{g,n}$.

Remark 2.1.1. The notation $\mathbf{x}=(x_1,\dots,x_n)$ as well as its close cousins will only be used in Section 2 and will not conflict with the usage as a generator of a Heegaard Floer chain group in the other sections of this paper.

Definition 2.1.2. A (J,ν)-*holomorphic map* is a map

$$(u, \phi_0 \circ st) : (F,j,\mathbf{x}) \to X \times \overline{\mathcal{U}}_{g,n}$$

which satisfies the perturbed J-holomorphic equation

$$\overline{\partial}_J u = (u, \phi_0 \circ st)^* \nu. \tag{2.1.1}$$

We will usually abbreviate Equation (2.1.1) as $\overline{\partial}_J u = \nu$ and not mention $\phi_0 \circ st$.

Now let V be a codimension 2 symplectic submanifold of X. Then (J,ν) is V-*compatible* if TV is J-invariant, the component of ν in the direction normal to V is zero, and certain tensors given in Definition 3.2 (b) and (c) of [IP1] vanish.

From now on we assume that all (J,ν) are V-compatible without further mention.

Definition 2.1.3. A (J,ν)-holomorphic stable map

$$u : (F,j,\mathbf{x},\mathbf{x}') \to X,$$

with $\mathbf{x}=(x_1,\dots,x_n)$ and $\mathbf{x}'=(x_1',\dots,x_l')$, is said to be V-*regular* if no irreducible component of the domain, no marked point in $\mathbf{x}$, and no double point of u is mapped to V. The points in $\mathbf{x}'$ are mapped to V with multiplicities $\mathbf{s} = (s_1, \dots, s_l)$. We write $l(\mathbf{s}) = l$ and $\deg(\mathbf{s}) = \sum_{i=1}^{l} s_i$.

Let $\mathcal{M}_{g,n+l(\mathbf{s})}(X, A, J, \nu)$ be the moduli space of (J, ν)-holomorphic maps $u : (F, j, \mathbf{x} \cup \mathbf{x}') \to X$ in the class $A \in H_2(X)$ such that (F, j) is a genus g Riemann surface. Then we let

$$\mathcal{M} = \mathcal{M}^V_{g,n,\mathbf{s}}(X, A, J, \nu) \subset \mathcal{M}_{g,n+l(\mathbf{s})}(X, A, J, \nu)$$

be the moduli space of V-regular (J, ν)-holomorphic stable maps $u : (F, j, \mathbf{x}, \mathbf{x}') \to X$ in A such that $st(F, j) \in \overline{\mathcal{M}}_{g,n}$ and $\mathbf{x}'$ is mapped to V with multiplicities $\mathbf{s}$, modulo automorphisms of the domain. We also use the modifier "irr" to denote the subset of maps $u : (F, j, \mathbf{x}, \mathbf{x}') \to X$ such that (F, j) is irreducible. Then, for a generic (J, ν), $\mathcal{M}_{g,n+l(\mathbf{s})}(X, A, J, \nu)$ is transversely cut out and $\mathcal{M}^{irr} \subset \mathcal{M}_{g,n+l(\mathbf{s})}(X, A, J, \nu)$ is transversely cut out by [IP1, Lemma 4.2].

By [IP1, Theorem 7.4], there is a compactification $\overline{\mathcal{M}} = \overline{\mathcal{M}}^V_{g,n,\mathbf{s}}(X, A, J, \nu)$ of $\mathcal{M}$ such that $\overline{\mathcal{M}} - \mathcal{M}$ is the union of (real) codimension ≥ 2 strata, and $\overline{\mathcal{M}}$ carries a fundamental class. Here $\overline{\mathcal{M}}$ is the space of *V-stable maps* $u : (F, j, \mathbf{x}, \mathbf{x}') \to X$ as defined in [IP1, Definition 7.2]. A map $u \in \overline{\mathcal{M}} - \mathcal{M}$ can have irreducible components F_i which are mapped to V. Such components come equipped with solutions ξ_i of Cauchy-Riemann type equations on the pullback of the normal bundle $N_{X/V}$. The sections ξ_i keep track of how a sequence of V-regular maps approaches $u|_{F_i}$.

We then consider the evaluation map:

$$\overline{ev} : \overline{\mathcal{M}} \to X^n \times V^{l(\mathbf{s})}, \tag{2.1.2}$$

$$(u : (F, j, \mathbf{x}, \mathbf{x}') \to X) \mapsto (u(\mathbf{x}), u(\mathbf{x}')).$$

We write $\overline{ev}_{(J,\nu)}$ when we want to emphasize the pair (J, ν). We also write ev for the restriction of $\overline{ev}$ to $\mathcal{M}$.

Definition 2.1.4. The *relative Gromov-Witten invariant* $GW^V_{X,A,g,n,\mathbf{s}}$ is the cycle

$$\overline{ev}_*[\overline{\mathcal{M}}] \in H_*(X^n \times V^{l(\mathbf{s})}; \mathbb{Q}).$$

2.2. **First relative Gromov-Witten calculation.** Let Σ be a surface of genus g. We consider $(X, \omega) = \Sigma \times \mathbb{CP}^1$ with a product symplectic form. Let π_1 and π_2 be the projections of X onto Σ and $\mathbb{CP}^1$, respectively.

The goal of this subsection is to compute the top-dimensional summand of

$$GW^V_{X,A,g,n,\mathbf{s}} \in H_*(X^n \times V^{l(\mathbf{s})}),$$

where $V = \Sigma \times \{\infty\}$, $A = [\Sigma] + 2g[\mathbb{CP}^1]$, $g = g(\Sigma)$, $n = g + 1$, $\mathbf{s} = (1, \dots, 1)$, and $l(\mathbf{s}) = 2g$. In other words, we are computing the degree of $\overline{ev}$, which is an integer that we denote by G_1.

Theorem 2.2.1. $G_1 = 1$.

Since $X, A, g, n, \mathbf{s}$ are fixed in this subsection, we abbreviate

$$\mathcal{M}^V_{(J,\nu)} := \mathcal{M}^V_{g,n,\mathbf{s}}(X, A, J, \nu), \quad \mathcal{M}_{(J,\nu)} := \mathcal{M}_{g,n+l(\mathbf{s})}(X, A, J, \nu).$$

In the proof of Theorem 2.2.1, instead of a generic (J,ν), we will use $(J,0)$ with J a product complex structure to carry out our calculations. This is legitimate because of certain automatic transversality results which will be discussed in Section 2.2.1. From now on assume that J is a product complex structure.

2.2.1. *Regularity.* Let $\mathcal{M}_g(X,A,J)$ be the moduli space of holomorphic maps

$$u:(F,j)\to(X,J)$$

in the class A, modulo automorphisms of the domain. Here F is a closed surface of genus g and j ranges over all complex structures on F.

Recall Hofer-Lizan-Sikorav's automatic transversality theorem [HLS, Theorem 1] (first suggested by Gromov [Gr]), which states that the moduli space $\mathcal{M}_g(X,A,J)$ is regular in a neighborhood of u, provided u is embedded and $c_1(A)=\langle c_1(TM),A\rangle>0$. If N is the normal bundle of an embedded curve u, then $c_1(A)>0$ is equivalent to $c_1(N)>2g-2$ by the adjunction formula $c_1(A)=c_1(T\Sigma)+c_1(N)$.

The following lemma is a consequence of automatic transversality:

Lemma 2.2.2. *The moduli space $\mathcal{M}_g(X,A,J)$ is regular and*

$$\dim_{\mathbb{R}}\mathcal{M}_g(X,A,J)=6g+2.$$

Proof. Any curve $u\in\mathcal{M}_g(X,A,J)$ is clearly embedded, since the projection $\pi_1\circ u$ is a biholomorphism. For an embedded curve u in the class $A=[\Sigma]+2g[\mathbb{CP}^1]$,

$$c_1(N)=A\cdot A=4g>2g-2.$$

This implies the regularity of $\mathcal{M}_g(X,A,J)$.

Now $c_1(A)=c_1(T\Sigma)+c_1(N)=2g+2$, and the Fredholm index is given by:

$$\mathrm{ind}(u)=-\chi(F)+2c_1(A)=(2g-2)+2(2g+2)=6g+2.$$

By the regularity of $\mathcal{M}_g(X,A,J)$, $\dim\mathcal{M}_g(X,A,J)=\mathrm{ind}(u)=6g+2$. □

Let $u\in\mathcal{M}_g(X,A,J)$. Since $\pi_1\circ u:F\to\Sigma$ must have degree one, it follows that (F,j) must be biholomorphic to our chosen Σ. Now fix a particular identification of (F,j) with Σ. This has the effect of eliminating automorphisms of the domain, provided we take Σ to be sufficiently generic. Hence there is a one-to-one correspondence between $u\in\mathcal{M}_g(X,A,J)$ and meromorphic functions (i.e., branched covers)

$$v=\pi_2\circ u:\Sigma\to\mathbb{CP}^1$$

of degree $2g$ and u can be recovered by taking the graph of the function v.

Corollary 2.2.3. *The moduli space $\mathcal{M}^{irr}_{(J,0)}$ is regular and the moduli space $\mathcal{M}^{V,irr}_{(J,0)}$ is transversely cut out in $\mathcal{M}^{irr}_{(J,0)}$.*

Proof. The moduli space $\mathcal{M}^{irr}_{(J,0)}$ is regular by Lemma 2.2.2, since we are just adjoining the variables $\mathbf{x}=(x_1,\dots,x_{g+1}),\mathbf{x}'=(x'_1,\dots,x'_{2g})$ to each $u:(F,j)\to X$. The second statement of the corollary is more precisely stated as:

$$ev(\mathcal{M}^{V,irr}_{(J,0)})\pitchfork(X^n\times V^{l(\mathbf{s})})\ \text{ in }\ X^{n+l(\mathbf{s})}, \tag{2.2.1}$$

where $ev:\mathcal{M}_{(J,0)}\to X^{n+l(\mathbf{s})}$ is the evaluation map $ev(u,\mathbf{x},\mathbf{x}')=(u(\mathbf{x}),u(\mathbf{x}'))$. If $(u,\mathbf{x},\mathbf{x}')\in\mathcal{M}^{V,irr}_{(J,0)}$, then u is transverse to V and intersects it at $2g$ points by definition. Hence Condition (2.2.1) is easily attained at $(u,\mathbf{x},\mathbf{x}')$ by considering the variations of $\mathbf{x}'$ while keeping $(u,\mathbf{x})$ fixed. □

2.2.2. *The Riemann-Roch theorem.* Let Σ be a closed Riemann surface of genus g. The classical Riemann-Roch theorem states that:

$$l(D)-l(K-D)=\deg(D)+1-g,$$

where D is a divisor, K is the canonical divisor, and $l(D)$ is the dimension of the space of meromorphic functions with poles at most at D.

Lemma 2.2.4. *If* $\deg(D)=2g-1+m$ *with* $m\geq 0$, *then* $l(D)=g+m$.

Proof. Since $\deg(K)=2g-2$, we have $\deg(K-D)<0$ and $l(K-D)=0$. Hence $l(D)=\deg(D)+1-g=(2g-1+m)+1-g=g+m$. □

2.2.3. *Proof of Theorem 2.2.1.* Let $\widetilde{\mathbf{x}}=(\widetilde{x}_1,\dots,\widetilde{x}_{g+1})$ and $\widetilde{\mathbf{x}}'=(\widetilde{x}'_1,\dots,\widetilde{x}'_{2g})$ be ordered $(g+1)$- and $2g$-tuples in Σ and let $\mathbf{y}=(y_1,\dots,y_{g+1})$ be an ordered $(g+1)$-tuple in $\mathbb{CP}^1$. We assume that $(\widetilde{\mathbf{x}},\widetilde{\mathbf{x}}')$ and $\mathbf{y}$ are in generic position; in particular, the points of $(\widetilde{\mathbf{x}},\widetilde{\mathbf{x}}')$ are distinct and the points of $\mathbf{y}$ are distinct and different from ∞. If $(u,\mathbf{x},\mathbf{x}')\in\mathcal{M}^{V,irr}_{(J,0)}\cap\overline{ev}^{-1}((\widetilde{\mathbf{x}},\mathbf{y}),\widetilde{\mathbf{x}}')$,[1] where $u:(F,j)\to(X,J)$ is a holomorphic map, then $\mathbf{x}=\widetilde{\mathbf{x}}$ and $\mathbf{x}'=\widetilde{\mathbf{x}}'$, since (F,j) is identified with the Riemann surface Σ. Henceforth, we write $\mathbf{x}$ and $\mathbf{x}'$ instead of $\widetilde{\mathbf{x}}$ and $\widetilde{\mathbf{x}}'$.

The moduli spaces $\mathcal{M}^{V,irr}_{(J,0)}$ and $\mathcal{M}^{irr}_{(J,0)}$ are regular by Corollary 2.2.3. The restriction of the evaluation map

$$ev_{(J,0)}:\mathcal{M}^{V}_{(J,0)}\to X^n\times V^{l(\mathbf{s})}$$

to $\mathcal{M}^{V,irr}_{(J,0)}$ will be denoted by $ev^{irr}_{(J,0)}$. Let us write $\mathfrak{E}=(ev_{(J,0)})^{-1}((\mathbf{x},\mathbf{y}),\mathbf{x}')$ and $\mathfrak{E}^{irr}=(ev^{irr}_{(J,0)})^{-1}((\mathbf{x},\mathbf{y}),\mathbf{x}')$.

Lemma 2.2.5. *If* $(\mathbf{x},\mathbf{x}')$ *and* $\mathbf{y}$ *are generic, then* $\mathfrak{E}^{irr}$ *is compact.*

Proof. Arguing by contradiction, let $u_i\in\mathfrak{E}^{irr}$ be a sequence that converges to a limit curve $u:\Sigma'\to X$ with a bubble component. Since u is $(J,0)$-holomorphic, $\pi_1\circ u$ is holomorphic and one component of Σ' must be biholomorphic to Σ. Hence Σ' is obtained from Σ by attaching spheres which are mapped by u to $\{x''_i\}\times\mathbb{CP}^1$, $1\leq i\leq k$. Moreover, $k\leq 2g$ by the positivity of intersections, since $\langle u(\Sigma'),\Sigma\times\{pt\}\rangle=2g$, where $\langle\cdot,\cdot\rangle$ is the intersection pairing.

(1) Suppose that $u(\Sigma)\neq V=\Sigma\times\{\infty\}$. Since $\langle u(\Sigma'),V\rangle=2g$ and the intersection points are $(x'_1,\infty),\dots,(x'_{2g},\infty)$, we must have $\mathbf{x}''=\{x''_1,\dots,x''_k\}\subset\mathbf{x}'$.[2] We claim the map $u|_\Sigma$ cannot exist. Let us write $v=\pi_2\circ u|_\Sigma:\Sigma\to\mathbb{CP}^1$. Then v is a meromorphic function with poles at $\mathbf{x}'-\mathbf{x}''$, subject to $v(x_i)=y_i$

---

[1] By slight abuse of notation, $(\widetilde{\mathbf{x}},\mathbf{y})$ means $((\widetilde{x}_1,y_1),\dots,(\widetilde{x}_{g+1},y_{g+1}))$.

[2] Here we are slightly abusing notation and viewing $\mathbf{x}'',\mathbf{x}'$ as sets.

for all $i = 1, \dots, g+1$. Hence $v \in l(D' - D'')$, where $D' = x'_1 + \cdots + x'_{2g}$ and $D'' = x''_1 + \cdots + x''_k$. By Lemma 2.2.4, $l(D' - D'') \leq g$. On the other hand, there are more constraints $v(x_i) = y_i$ than there are linearly independent functions. Hence v cannot exist provided $\mathbf{x}$ and $\mathbf{y}$ are generic, a contradiction.

(2) Now suppose that $u(\Sigma) = V$. Then u consists of $u|_\Sigma$, together with $2g$ bubbles $\{x''_i\} \times \mathbb{CP}^1$, $i = 1, \dots, 2g$. The set $\mathbf{x}'' = \{x''_1, \dots, x''_{2g}\}$ contains $\mathbf{x}$ and may also contain $k \leq g-1$ elements of $\mathbf{x}'$. We then apply the renormalization procedure of [IP1, Proposition 6.6] to the sequence u_i restricted to a neighborhood of V. After choosing suitable restrictions and rescalings, the sequence u_i converges to a meromorphic function $\xi : \Sigma \to \mathbb{CP}^1$ which encodes how the curves $u_i(\Sigma)$ approach V. The function ξ has poles at most at $D'' = x''_1 + \cdots + x''_{2g}$ and zeros at $2g - k \geq g+1$ points of $\mathbf{x}'$. The details of this argument are left to the reader. Since $l(D'') = g+1$ by Lemma 2.2.4, we have at least as many constraints as linearly independent functions, which is again a contradiction. $\Box$

Lemma 2.2.6. *If* $(\mathbf{x}, \mathbf{x}')$ *and* $\mathbf{y}$ *are generic, then* $\mathfrak{E} = \mathfrak{E}^{irr}$.

Proof. Suppose $u : \Sigma' \to X$ is a stable curve in $\mathfrak{E} - \mathfrak{E}^{irr}$. Then Σ' is obtained from Σ by attaching spheres which are mapped by u to $\{x''_i\} \times \mathbb{CP}^1$, $1 \leq i \leq k$. By the V-regularity of u, $u(\Sigma')$ does not have a component $V = \Sigma \times \{\infty\}$. Hence the argument from Lemma 2.2.6(1) gives a contradiction. $\Box$

By Corollary 2.2.3 and Lemmas 2.2.5 and 2.2.6, G_1 is the number of meromorphic functions $v : \Sigma \to \mathbb{CP}^1$ with poles at $\mathbf{x}'$ such that $v(x_i) = y_i$ for all $i = 1, \dots, g+1$. By Lemma 2.2.4, $l(x'_1 + \cdots + x'_{2g}) = g+1$. On the other hand, since there are $g+1$ constraints $v(x_i) = y_i$, there is a unique solution for generic $\mathbf{x}$, $\mathbf{y}$. This shows that $G_1 = 1$.

2.3. **Second relative Gromov-Witten calculation.**

2.3.1. *Definitions.* Let $\Sigma = \overline{S}$, where $\overline{S} = S \cup D^2$ is obtained by capping off a page S of an open book as in Section I.5.1.2. Also recall the point at infinity $z_\infty = \{\rho = 0\} \in \overline{S}$.

Consider (X, ω), where $X = \Sigma \times \mathbb{CP}^1$ and ω is a product symplectic form. Let $V = V_1 \cup V_\infty$, where $V_* = \Sigma \times \{*\}$ and $* = 1, \infty$. We take $A = [\Sigma] + 2g[\mathbb{CP}^1]$, $g = g(\Sigma)$, $n = 1$, and $\mathbf{s} = \mathbf{s}_1 \cup \mathbf{s}_\infty$, where:

- $\mathbf{s}_1 = (s_{1,1}, \dots, s_{1,l})$, $l(\mathbf{s}_1) = l$, $\deg(\mathbf{s}_1) = 2g$;
- $\mathbf{s}_\infty = (1, \dots, 1)$, $l(\mathbf{s}_\infty) = 2g$, $\deg(\mathbf{s}_\infty) = 2g$; and
- $l(\mathbf{s}) = l(\mathbf{s}_1) + l(\mathbf{s}_\infty)$.

Let $\mathcal{M}_{(J,\nu)} := \mathcal{M}_{g,n+l(\mathbf{s})}(X, A, J, \nu)$ and let

$$\mathcal{M}^{V_1,V_\infty}_{(J,\nu)} := \mathcal{M}^{V_1,V_\infty}_{g,n,\mathbf{s}_1,\mathbf{s}_\infty}(X, A, J, \nu) \subset \mathcal{M}^V_{g,n,\mathbf{s}}(X, A, J, \nu)$$

be the subset consisting of V-regular (J,ν)-holomorphic stable maps

$$u : (F, j, x, \mathbf{x}'_1, \mathbf{x}'_\infty) \to X$$

in the class A, where

$$\mathbf{x}'_1 = (x'_{1,1}, \dots, x'_{1,l(\mathbf{s}_1)}), \quad \mathbf{x}'_\infty = (x'_{\infty,1}, \dots, x'_{\infty,l(\mathbf{s}_\infty)}),$$

and $\mathbf{x}'_*$, $* = 1, \infty$, is mapped to V_* with multiplicities $\mathbf{s}_*$. For $* = 1, \infty$, let

$$\mathcal{M}^{V_*}_{(J,\nu)} := \mathcal{M}^{V_*}_{g,n,\mathbf{s}_*}(X, A, J, \nu)$$

be the moduli space of V_*-regular (J, ν)-holomorphic stable maps $(u, \mathbf{x}'_*)$ in A such that $\mathbf{x}'_*$ is mapped to V_* with multiplicities $\mathbf{s}_*$.

We define the evaluation map

$$\overline{ev}_{(J,\nu)} : \overline{\mathcal{M}}^{V_1,V_\infty}_{(J,\nu)} \to X \times V_1^{l(\mathbf{s}_1)} \times V_\infty^{l(\mathbf{s}_\infty)},$$

$$(u : (F, j, x, \mathbf{x}'_1, \mathbf{x}'_\infty) \to X) \mapsto (u(x), u(\mathbf{x}'_1), u(\mathbf{x}'_\infty)),$$

and, for $* = 1, \infty$, we define the evaluation maps

$$\overline{ev}^{V_*}_{(J,\nu)} : \overline{\mathcal{M}}^{V_*}_{(J,\nu)} \to V_*^{l(\mathbf{s}_*)}, \quad (u, \mathbf{x}'_*) \mapsto u(\mathbf{x}'_*).$$

Let $ev_{(J,\nu)}$ and $ev^{irr}_{(J,\nu)}$ be the restrictions of $\overline{ev}_{(J,\nu)}$ to $\mathcal{M}^{V_1,V_\infty}_{(J,\nu)}$ and $\mathcal{M}^{V_1,V_\infty,irr}_{(J,\nu)}$. The restrictions $ev^{V_*}_{(J,\nu)}$ and $ev^{V_*,irr}_{(J,\nu)}$ of $\overline{ev}^{V_*}_{(J,\nu)}$ are defined similarly.

Definition 2.3.1. A pair $((z_0, y_0), \mathbf{z}'_1)$ consisting of $(z_0, y_0) \in X$ and $\mathbf{z}'_1 \in \Sigma^{l(\mathbf{s}_1)}$ is *generic* if (z_0, y_0) is a generic point in X and $(z_0, \mathbf{z}'_1)$ is a generic $(l(\mathbf{s}_1) + 1)$-tuple in $\Sigma^{l(\mathbf{s}_1)+1}$. In particular, $y_0 \neq 1, \infty$ and the points of $(z_0, \mathbf{z}'_1)$ are disjoint and do not lie on any $\overline{a}_i$ given in Section I.5.2.2.

Definition 2.3.2. The relative Gromov-Witten invariant $G_2(\Sigma, \mathbf{s}_1)$ is defined as follows:

$$G_2(\Sigma, \mathbf{s}_1) = \langle (\overline{ev}_{(J,\nu)})_*([\overline{\mathcal{M}}^{V_1,V_\infty}_{(J,\nu)}]), \{(z_0, y_0)\} \times \mathbf{z}'_1 \times \overline{a}_1 \times \cdots \times \overline{a}_{2g} \rangle,$$

where $((z_0, y_0), \mathbf{z}'_1)$ is generic. When we want to specify the constraints, we write $G_2(\Sigma, \mathbf{s}_1; \{(z_0, y_0)\} \times \mathbf{z}'_1 \times \overline{a}_1 \times \cdots \times \overline{a}_{2g})$.

The main theorem of this subsection is the following:

Theorem 2.3.3. *If* $\deg(\mathbf{s}_1) = 2g$, *then* $G_2(\Sigma, \mathbf{s}_1)$ *does not depend on* $\mathbf{s}_1$ *and* $G_2(\Sigma, \mathbf{s}_1) = 1$.

2.3.2. *First steps in the calculation of* G_2. We will use $(J, \nu) = (J, 0)$, where $J = j_\Sigma \times j_{\mathbb{CP}^1}$ is a split complex structure on $\Sigma \times \mathbb{CP}^1$.

There is a one-to-one correspondence between $(u, x, \mathbf{x}'_1, \mathbf{x}'_\infty) \in \mathcal{M}^{V_1,V_\infty,irr}_{(J,0)}$ and meromorphic functions

$$v : (F, j, x, \mathbf{x}'_1, \mathbf{x}'_\infty) \to \mathbb{CP}^1$$

satisfying $\Sigma = (F, j)$, $v(\mathbf{x}'_1) = 1$ and $v(\mathbf{x}'_\infty) = \infty$, where the correspondence is given by $v = \pi_2 \circ u$. Moreover, if

$$ev^{irr}_{(J,0)}((u, x, \mathbf{x}'_1, \mathbf{x}'_\infty)) \in \{(z_0, 0)\} \times \mathbf{z}'_1 \times \overline{a}_1 \times \cdots \times \overline{a}_{2g},$$

then $(x,\mathbf{x}'_1)=(z_0,\mathbf{z}'_1)$ and $x'_{2g+i}\in\overline{a}_i$, $i=1,\dots,2g$. Henceforth we identify $(x,\mathbf{x}'_1)=(z_0,\mathbf{z}'_1)$. Since we have the freedom to postcompose v by a fractional linear transformation, we may (and we will) assume that $y_0=0$.

The analog of Corollary 2.2.3 is the following:

Lemma 2.3.4. *The moduli space* $\mathcal{M}^{V_1,irr}_{(J,0)}$ *is regular.*

Proof. Let $\mathcal{M}$ be the moduli space of V-regular $(J,0)$-holomorphic maps to X in the class A, which is regular by Lemma 2.2.2. We consider the map

$$\pi:\mathcal{M}\to Sym^{2g}(\Sigma),$$

which sends u to the intersection of u with V_1, counted with multiplicity. Each fiber $\pi^{-1}(D)$ is a dense open subset of the space $H^0(\Sigma,\mathcal{O}_D)$ of meromorphic sections of Σ with poles at most at D. By Lemma 2.2.4, $H^0(\Sigma,\mathcal{O}_D)$ is a complex vector space of dimension $l(D)=g+1$ for each $D\in Sym^{2g}(\Sigma)$. Hence π is the restriction of a holomorphic vector bundle

$$\widetilde{\pi}:\mathcal{E}\to Sym^{2g}(\Sigma)$$

to the dense open subset $\mathcal{M}$, where $\widetilde{\pi}^{-1}(D)=H^0(\Sigma,\mathcal{O}_D)$. In particular, π is a submersion.

Let $Z_{\mathbf{s}_1}\subset \mathrm{Sym}^{2g}(\Sigma)$ be the submanifold consisting of divisors of type $D=\sum_{i=1}^{l(\mathbf{s}_1)}s_{1,i}z'_{1,i}$, $\deg(D)=\deg(\mathbf{s}_1)=2g$, with pairwise distinct $z'_{1,i}$. Then π is transverse to $Z_{\mathbf{s}_1}\subset\mathrm{Sym}^{2g}(\Sigma)$. This implies the regularity of $\mathcal{M}^{V_1,irr}_{(J,0)}$. □

Consider the following genericity conditions:

($\mathfrak{G}_1$) $((z_0,0),\mathbf{z}'_1)$ is generic.

($\mathfrak{G}_2$) $(z_0,0)=(z_\infty,0)$,[3] $\mathbf{z}'_1$ is generic, and $\overline{a}_1,\dots,\overline{a}_{2g}$ are generic.

Let us write

$$\mathfrak{E}=(ev_{(J,0)})^{-1}(\{(z_0,0)\}\times\mathbf{z}'_1\times\overline{a}_1\times\cdots\times\overline{a}_{2g}),$$
$$\mathfrak{E}^{irr}=(ev^{irr}_{(J,0)})^{-1}(\{(z_0,0)\}\times\mathbf{z}'_1\times\overline{a}_1\times\cdots\times\overline{a}_{2g}).$$

The following lemmas are analogous to Lemmas 2.2.5 and 2.2.6:

Lemma 2.3.5. *If* ($\mathfrak{G}_1$) *or* ($\mathfrak{G}_2$) *holds, then* $\mathfrak{E}$ *is compact.*

Proof. Arguing by contradiction, suppose the limit curve $u:\Sigma'\to X$ has a bubble component. A nonconstant sphere bubble must be mapped to some $\{w\}\times\mathbb{CP}^1$. We have the homological constraint:

$$\langle u(\Sigma'),V_1\rangle=\langle u(\Sigma'),V_\infty\rangle=2g. \tag{2.3.1}$$

(1) Suppose ($\mathfrak{G}_1$) holds. If $w\notin\mathbf{z}'_1$, then V_1 must be a component of $u(\Sigma')$; otherwise, $\langle u(\Sigma'),V_1\rangle\geq g+1$, which contradicts Equation (2.3.1). Hence $u(\Sigma')$ consists of V_1 and $2g$ bubbles $\{z'_{\infty,i}\}\times\mathbb{CP}^1$, where $z'_{\infty,i}\in\overline{a}_i$, $i=1,\dots,2g$, and w is equal to some $z'_{\infty,i}$. This contradicts the fact that $u(\Sigma')$ passes through $(z_0,0)$.

---

[3] Here z_∞ is the point at infinity.

On the other hand, if $w \in \mathbf{z}'_1$, then $w \notin \overline{\mathbf{a}}$. Again, by Equation (2.3.1), $u(\Sigma')$ consists of V_∞ and bubbles $\{z'_{1,i}\} \times \mathbb{CP}^1$, $i = 1, \dots, l(\mathbf{s}_1)$, of total multiplicity $2g$, and this contradicts the fact that $u(\Sigma')$ passes through $(z_0, 0)$.

(2) Suppose ($\mathfrak{G}_2$) holds. If $w \in \mathbf{z}'_1$, then the proof is the same as (1).

On the other hand, if $w \notin \mathbf{z}'_1$, then $u(\Sigma')$ consists of V_1 and $2g$ bubbles $\{z'_{\infty,i}\} \times \mathbb{CP}^1$, $z'_{\infty,i} \in \overline{a}_i$. Since $u(\Sigma')$ passes through $(z_\infty, 0)$, we must have $z'_{\infty,i} = z_\infty$ for one or more $i = 1, \dots, 2g$. Let $D(\overline{\mathbf{a}})$ be the set of divisors $z'_{\infty,1} + \dots + z'_{\infty,2g}$, $z'_{\infty,i} \in \overline{a}_i$, such that $z'_{\infty,i} = z_\infty$ for at least one i. Applying the renormalization procedure of [IP1, Proposition 6.6] as in Lemma 2.2.5, we obtain a meromorphic function $\xi : \Sigma \to \mathbb{CP}^1$ with divisor $D(\xi) = D_2 - D_1$, where $D_1 \in D(\overline{\mathbf{a}})$ and $D_2 \in Z_{\mathbf{s}_1}$.

By the Abel-Jacobi theorem, the Picard group $\mathrm{Pic}^{2g}(\Sigma)$ of divisors of degree $2g$ modulo linear equivalence is isomorphic to a g-dimensional complex torus. Since $\dim_{\mathbb{R}} D(\overline{\mathbf{a}}) < 2g$, its image in $\mathrm{Pic}^{2g}(\Sigma)$ has $\dim_{\mathbb{R}} < 2g$. Hence, a generic $D_2 \in Z_{\mathbf{s}_1}$ is not linearly equivalent to any $D_1 \in D(\overline{\mathbf{a}})$, provided $\overline{a}_1, \dots, \overline{a}_{2g}$ are generic. This contradicts the existence of ξ. □

Lemma 2.3.6. *If ($\mathfrak{G}_1$) or ($\mathfrak{G}_2$) holds, then $\mathfrak{C} = \mathfrak{C}^{irr}$.*

Proof. Suppose $u : \Sigma' \to X$ is a stable curve in $\mathfrak{C} - \mathfrak{C}^{irr}$. Then $u(\Sigma')$ has a sphere bubble $\{w\} \times \mathbb{CP}^1$. By the V-regularity of u, V_1 and V_∞ cannot be components of $u(\Sigma')$. Suppose ($\mathfrak{G}_1$) or ($\mathfrak{G}_2$) holds; the key point is that the points of $\mathbf{z}'_1$ are disjoint and do not lie on any $\overline{a}_i$. If $w \notin \mathbf{z}'_1$, then V_1 must be a component of $u(\Sigma')$ by Equation (2.3.1), and if $w \in \mathbf{z}'_1$, then V_∞ must be a component of $u(\Sigma')$ by Equation (2.3.1). Both are contradictions. □

Lemma 2.3.7. *If ($\mathfrak{G}_1$) or ($\mathfrak{G}_2$) holds, then the invariant $G_2(\Sigma, \mathbf{s}_1)$ is the (mod 2) count of holomorphic maps*

$$v : (\Sigma, z_0, \mathbf{z}'_1, \mathbf{z}'_\infty) \to \mathbb{CP}^1,$$

such that $v(z_0) = 0$, $v(z'_{1,i}) = 1$ with multiplicity $s_{1,i}$ for $i = 1, \dots, l(\mathbf{s}_1)$, $v(\mathbf{z}'_\infty) = \infty$, and $z'_{\infty,i} \in \overline{a}_i$ for $i = 1, \dots, 2g$.

Proof. This follows from Lemmas 2.3.4, 2.3.5, and 2.3.6. □

Lemma 2.3.8. *$G_2(\Sigma, \mathbf{s}_1)$ does not depend on $\mathbf{s}_1$.*

Proof. Let us write $ev^{V_1, irr, \mathbf{s}_1}_{(J,0)}$ and $\mathcal{M}^{V_1, irr, \mathbf{s}_1}_{(J,0)}$ for $ev^{V_1, irr}_{(J,0)}$ and $\mathcal{M}^{V_1, irr}_{(J,0)}$ corresponding to $\mathbf{s}_1$. By the proof of Lemma 2.3.4, the moduli spaces $\mathcal{M}^{V_1, irr, \mathbf{s}_1}_{(J,0)}$ for all the $\mathbf{s}_1$ can be smoothly glued into $\pi : \mathcal{M} \to \mathrm{Sym}^{2g}(\Sigma)$. The lemma then follows from Lemma 2.3.7 by generically intersecting with $(z_0, 0)$ and $(\overline{a}_i, \infty)$, $i = 1, \dots, 2g$. □

In view of Lemma 2.3.8, from now on we may assume that $\mathbf{s}_1 = (1, \dots, 1)$ with $l(\mathbf{s}_1) = 2g$. For simplicity we write $G_2(\Sigma)$ for $G_2(\Sigma, \mathbf{s}_1)$.

The proof strategy for Theorem 2.3.3 is to first compute $G_2(T^2)$ and then reduce the higher genus case to $g = 1$.

2.3.3. *Case of* T^2. We consider the situation where $\Sigma = T^2$. Suppose $(\mathfrak{G}_1)$ holds with $g = 1$.

Lemma 2.3.9. *If* $z_1' \neq z_2'$*, then every meromorphic function on* T^2 *with poles at* z_1' *and* z_2' *of order* 1 *can be written as* $a_1 + a_2 f$*, where* $a_1, a_2 \in \mathbb{C}$*,* $a_2 \neq 0$*, and* f *is a function on* T^2 *with poles at* z_1' *and* z_2' *of order* 1.

In other words, f is unique up to a postcomposition by a fractional linear transformation $\mathbb{CP}^1 \to \mathbb{CP}^1$ which fixes ∞.

Proof. By Lemma 2.2.4, $l(z_i') = 1$ and is given by the constants, and $l(z_1'+z_2') = 2$ and one dimension is taken by the constants. The lemma follows. □

Lemma 2.3.10. $G_2(T^2) = 1$.

Proof. Suppose $(\mathfrak{G}_1)$ holds with $g = 1$. In view of Lemmas 2.3.7 and 2.3.8, $G_2(T^2)$ is the count of irreducible curves $u \in \mathcal{M}_{g=1}(X, A, J)$ that pass through $(z_0, 0)$, $(z_1', 1)$, $(z_2', 1)$, $\overline{a}_1 \times \{\infty\}$, and $\overline{a}_2 \times \{\infty\}$, where u intersects $\overline{a}_1 \times \{\infty\}$ and $\overline{a}_2 \times \{\infty\}$ at distinct points.



FIGURE 1. The branched double cover v of the torus T^2. The branch points $w_1, \dots, w_4$ of v are indicated by $\circ$.

According to Lemma 2.3.9, a branched double cover $v : T^2 \to \mathbb{CP}^1$ satisfying $v(z_0) = 0$ and $v(z_1') = v(z_2') = 1$ is determined up to postcomposition by $\eta \in PSL(2,\mathbb{C})$ which fixes 0 and 1; see Figure 1. Let $w_1, \dots, w_4 \in \mathbb{CP}^1$ be the branch points of v. Since z_1' and z_2' are generic, we may assume that $w_i \neq 0, 1$ for $i = 1, \dots, 4$. Let C_{ij} be a simple closed curve which bounds a small neighborhood of an arc from w_i to w_j. We may assume that C_{12} and C_{23} intersect only at 2 points y_1, y_2. Let $\overline{a}_1, \overline{a}_2$ be lifts of C_{12}, C_{23} such that $\#(\overline{a}_1 \cap \overline{a}_2) = 1$ and $v(\overline{a}_1 \cap \overline{a}_2) = y_1$. We now impose the condition $\eta \circ v(z_3') = \eta \circ v(z_4') = \infty$, where $z_3' \in \overline{a}_1$, $z_4' \in \overline{a}_2$, and η is a fractional linear transformation which fixes $0, 1$. This is possible only if $z_3', z_4' \in v^{-1}(y_i)$ for $i = 1$ or 2. On the other hand, since y_1, y_2 are not branch points, z_3' and z_4' must be distinct. Hence the only possibility is $\{z_3', z_4'\} = v^{-1}(y_2)$ and $\eta \in PSL(2, \mathbb{C})$ is the uniquely determined by $\eta(0) = 0$, $\eta(1) = 1$, $\eta(y_2) = \infty$. This completes the proof of the lemma. $\square$

2.3.4. *Proof of Theorem 2.3.3.* We now reduce to the torus case and prove Theorem 2.3.3. Suppose that $g(\Sigma) = 2$; the general case is similar.

The strategy is to degenerate the fiber Σ into a nodal surface Σ^∞ consisting of two irreducible components of genus one. More precisely, let $\Sigma^\tau = (\overline{S}, \overline{i}_\tau)$, $\tau \in [0, \infty)$, be a 1-parameter family of $g = 2$ Riemann surfaces which degenerates into a nodal Riemann surface

$$(\Sigma^\infty, \mathfrak{n}) = ((\Sigma_-, \mathfrak{n}_-) \sqcup (\Sigma_+, \mathfrak{n}_+)) / \mathfrak{n}_- \sim \mathfrak{n}_+,$$

where Σ_-, Σ_+ have genus one and $\mathfrak{n} = \{\mathfrak{n}_-, \mathfrak{n}_+\}$. The nodal surface Σ^∞ is obtained topologically as follows: Let γ be a separating curve in $\overline{S}$ and let $\sim_0$ be an equivalence relation on $\overline{S}$ such that $x \sim_0 y$ if $x, y \in \gamma$. Then $\Sigma^\infty = \overline{S}/\sim_0$. We also set $\widetilde{\mathbf{a}} = \overline{\mathbf{a}}/\sim_0$. Here $\overline{\mathbf{a}}$ and γ are fixed, while the complex structure $\overline{i}_\tau$ varies as $\tau \to \infty$.

Since $G_2(\Sigma)$ only depends on the homology classes of the $\overline{a}_j$, $j = 1, \dots, 4$, we may assume that the following hold:

(i) γ is disjoint from the $\overline{a}_j$, $j = 1, \dots, 4$, $\widetilde{a}_1, \widetilde{a}_2 \subset \Sigma_+$, and $\widetilde{a}_3, \widetilde{a}_4 \subset \Sigma_-$.
(ii) The points $z_0, z_1', \dots, z_4' \in \overline{S} - \gamma$ are fixed for all τ and in the limit $z_0, z_1', z_2' \in \Sigma_+$ and $z_3', z_4' \in \Sigma_-$.
(iii) $\overline{a}_1, \dots, \overline{a}_4$ are generic and $z_0, z_1', \dots, z_4'$ are generic.

Let $u^\tau : \Sigma^\tau \to \Sigma^\tau \times \mathbb{CP}^1$, $\tau \to \infty$, be a sequence of curves in the class $A = [\Sigma^\tau \times \{pt\}] + 4[\{pt\} \times \mathbb{CP}^1]$ that pass through

$$(z_0, 0), (z_1', 1), \dots, (z_4', 1), \overline{a}_1 \times \{\infty\}, \dots, \overline{a}_4 \times \{\infty\}.$$

By Lemma 2.3.5, no bubbling can occur at finite τ. Let $u = u_- \cup u_0 \cup u_+$ be the limit of u^τ, where $\mathrm{Im}(u_\pm) \subset \Sigma_\pm \times \mathbb{CP}^1$, $u_\pm$ has no components of $\{\mathfrak{n}\} \times \mathbb{CP}^1$, and u_0 maps to $\{\mathfrak{n}\} \times \mathbb{CP}^1$. If $u_\pm$ is irreducible, then we write $v_\pm = \pi_2 \circ u_\pm$, where π_2 is the projection to $\mathbb{CP}^1$.

Lemma 2.3.11. *The levels u_+ and u_- are irreducible and $u_0 = \varnothing$.*

Proof. The proof is similar to that of Lemma 2.3.5. Arguing by contradiction, suppose u_+ has a bubble component. The bubble is necessarily of the form $\{z\} \times \mathbb{CP}^1$. When τ is finite, $\mathrm{Im}(u^\tau) \cap (\Sigma^\tau \times \{1\}) = \{(z_1', 1), \dots, (z_4', 1)\}$.

We claim that $\Sigma_+ \times \{1\}$ or $\Sigma_+ \times \{\infty\}$ must be a component of u_+. Otherwise, suppose that $\Sigma_+ \times \{w\}$ is not a component of u_+ for $w = 1$ or ∞. Since z

cannot be in $\{z_1', z_2'\}$ and $\widetilde{a}_1 \cup \widetilde{a}_2$ at the same time, $\langle u_+, \Sigma_+ \times \{w\}\rangle \geq 3$ for $w \in \mathbb{CP}^1$. We also have $\langle u_-, \Sigma_- \times \{w\}\rangle \geq 2$ and $\langle u_0, \Sigma_\pm \times \{w\}\rangle \geq 0$. Hence $\langle u^\tau, \Sigma^\tau \times \{w\}\rangle > 4$ for $w \in \mathbb{CP}^1$, which is a contradiction. This implies the claim.

Now, the claim implies that u_+ consists of $\Sigma_+ \times \{w\}$, $w = 1$ or ∞, and 3 bubbles: $\{z_0\} \times \mathbb{CP}^1$ and either $\{z_1', z_2'\} \times \mathbb{CP}^1$ or $\{z_3', z_4'\} \times \mathbb{CP}^1$, where $z_{j+2}' \in \widetilde{a}_j$, $j = 1, 2$. Since $\langle u_-, \Sigma_- \times \{w\}\rangle \geq 2$ and $\langle u_0, \Sigma_\pm \times \{w\}\rangle \geq 0$, we again obtain $\langle u^\tau, \Sigma^\tau \times \{w\}\rangle > 4$ for $w \in \mathbb{CP}^1$. Hence u_+ does not have bubble components and is irreducible.

We can similarly prove that u_- is irreducible and $u_0 = \varnothing$. □

By Lemma 2.3.11, u_+ and u_- are irreducible. Hence we are counting pairs (v_-, v_+) such that $v_+(\mathfrak{n}_+) = v_-(\mathfrak{n}_-)$. By Lemma 2.3.10,

$$G_2(\Sigma_+) = G_2(\Sigma_+; \{(z_0, 0)\}, z_1', z_2', \widetilde{a}_1, \widetilde{a}_2) = 1$$

and is represented by a unique map v_+. Now the point $(\mathfrak{n}_-, v_+(\mathfrak{n}_+)) \in \Sigma_- \times \mathbb{CP}^1$ plays the role of $(z_0, 0)$ for v_-. Hence, by Lemma 2.3.10,

$$G_2(\Sigma_-) = G_2(\Sigma_-; \{(\mathfrak{n}_-, v_+(\mathfrak{n}_+))\}, z_3', z_4', \widetilde{a}_3, \widetilde{a}_4) = 1.$$

By (iii), the gluing conditions for u_+ and u_- are transverse (verification left to the reader) and

$$G_2(\Sigma) = G_2(\Sigma_+) \times G_2(\Sigma_-) = 1. \tag{2.3.2}$$

This completes the proof of Theorem 2.3.3.

2.4. **Third relative Gromov-Witten calculation.**

2.4.1. *Definitions.* Let D^2 be the closed unit disk $\{|z| \leq 1\} \subset \mathbb{C}$ and let $\Sigma = \overline{S}$. We consider (X, ω), where $X = \Sigma \times D^2$ and ω is a product symplectic form. Then $L_{\widehat{\mathbf{a}}} = \widehat{\mathbf{a}} \times \partial D^2 \subset \partial X$ is a Lagrangian submanifold of (X, ω). We write π_i, $i = 1, 2$, for the projection of X onto the ith factor. Let A be the relative homology class

$$A = [\Sigma \times \{pt\}] + 2g \cdot [\{pt\} \times D^2] \in H_2(X, \partial X),$$

and let $(F, j, x, \mathbf{x}')$ be a compact Riemann surface of genus g with $2g$ boundary components, an interior marked point x, and boundary marked points $\mathbf{x}' = (x_1', \dots, x_{4g}')$, where x_i', x_{2g+i}' are on the ith boundary component $\partial_i F$.

Let $\mathcal{J}$ be the space of compatible almost complex structures on (X, ω) and let $\mathcal{J}^{prod} \subset \mathcal{J}$ be the subset of product complex structures on X. Given $J \in \mathcal{J}$, let $\mathcal{M}(X, A, J)$ be the moduli space of holomorphic maps

$$u : (F, j, x, \mathbf{x}') \to (X, J)$$

in the class A such that $u(\partial_i F) \subset L_{\widehat{a}_i}$, modulo automorphisms of the domain.

Consider the evaluation map:

$$ev : \mathcal{M}(X, A, J) \to \mathfrak{X} = X \times L_{\widehat{a}_1} \times \cdots \times L_{\widehat{a}_{2g}} \times L_{\widehat{a}_1} \times \cdots \times L_{\widehat{a}_{2g}}, \tag{2.4.1}$$

$$(u, x, \mathbf{x}') \mapsto (u(x), u(x_1'), \dots, u(x_{4g}')).$$

Let $\mathbf{z} = (z_1, \dots, z_{4g})$ be a $4g$-tuple of points such that $z_i, z_{2g+i} \in \widehat{a}_i$, $i = 1, \dots, 2g$. Then we write

$$\mathcal{M}_J(z_\infty, \mathbf{z}) := ev^{-1}((z_\infty, 0), (z_1, 1), \dots, (z_{2g}, 1), (z_{2g+1}, -1), \dots, (z_{4g}, -1)),$$

where z_∞ is the point at infinity.

Definition 2.4.1. The relative Gromov-Witten type invariant $G_3(\Sigma; J; z_\infty, \mathbf{z})$ is the mod 2 count of equivalence classes $(u, x, \mathbf{x}') \in \mathcal{M}_J(z_\infty, \mathbf{z})$, where $J \in \mathcal{J}^{prod}$ and $\mathbf{z}$ are generic.

Theorem 2.4.2. *If $J \in \mathcal{J}^{prod}$ and $\mathbf{z}$ is generic, then $\mathcal{M}_J(z_\infty, \mathbf{z})$ is regular and compact. The count $G_3(\Sigma; J; z_\infty, \mathbf{z})$ is independent of the choice of $J \in \mathcal{J}^{prod}$ and $\mathbf{z}$ generic and is equal to* 1.

Proof. The first statement is proved in Lemmas 2.4.5 and 2.4.6. The independence of $G_3(\Sigma; J; z_\infty, \mathbf{z})$ is proved in Lemma 2.4.7 and $G_3(\Sigma; J; z_\infty, \mathbf{z})$ is computed in Sections 2.4.4–2.4.6. ☐

2.4.2. *Transversality.* We recall the following automatic transversality result from [HLS]; see Theorem 2 and Remark (1) following the statement of Theorem 2 in [HLS]:

Theorem 2.4.3 (Hofer-Lizan-Sikorav). *Let (F, j) be a compact Riemann surface with nonempty boundary, (M, J) be an almost complex 4-manifold, and $L \subset (M, J)$ be a totally real surface. Let*

$$u_0 : (F, \partial F, j) \to (M, L, J)$$

be a holomorphic map which sends ∂F to L. If u_0 is immersed and the sum of the Maslov indices of $u_0|_{\partial F}$ with respect to any unitary trivialization of $u_0^* TM$ is positive, then the set of holomorphic maps $u : (F, \partial F, j) \to (M, L, J)$ near u_0 is regular.

Using Theorem 2.4.3, we prove the following key result:

Lemma 2.4.4. *If $J \in \mathcal{J}^{prod}$, then the moduli space $\mathcal{M}(X, A, J)$ is regular and*

$$\dim_{\mathbb{R}} \mathcal{M}(X, A, J) = 8g + 4.$$

Proof. Let $(u, x, \mathbf{x}) \in \mathcal{M}(X, A, J)$, where $u : (F, j) \to (X, J)$. Since there are no branch points of $\pi_2 \circ u$ along ∂F, the curve u is immersed near ∂F. On the other hand, the restriction of $\pi_1 \circ u$ to $int(F)$ is a biholomorphism onto its image. Hence u is an immersion on all of F.

The total Maslov index $\mu(u)$ relative to the Lagrangian $L_{\widehat{\mathbf{a}}}$, with respect to any unitary trivialization of $u^* TX$ is:

$$\mu(u) = 2g \cdot \mu(\{pt\} \times D^2) + 2c_1(T\Sigma),$$

where $\{pt\} \times D^2$ has boundary on $L_{\widehat{\mathbf{a}}}$ and $\mu(\{pt\} \times D^2)$ is computed with respect to a trivialization of $TX|_{\{pt\} \times D^2}$. An easy calculation gives $\mu(\{pt\} \times D^2) = 2$ and $c_1(T\Sigma) = \chi(\Sigma) = 2 - 2g$. Hence

$$\mu(u) = 2g(2) + 2(2 - 2g) = 4 > 0. \tag{2.4.2}$$

The regularity of $\mathcal{M}(X, J, A)$ then follows from Theorem 2.4.3.

Using the usual index formula for holomorphic curves with Lagrangian boundary, we obtain:

$$\begin{aligned}\dim_{\mathbb{R}} \mathcal{M}(X, A, J) &= -\chi(F) + \mu(u) + (4g + 2)\\ &= -(2 - 4g) + 4 + (4g + 2) = 8g + 4,\end{aligned}$$

where the summand $4g+2$ comes from the marked points x and $\mathbf{x}'$. This completes the proof of Lemma 2.4.4. □

Lemma 2.4.5. *If $J \in \mathcal{J}^{prod}$, then $\mathcal{M}_J(z_\infty, \mathbf{z})$ is regular for generic $\mathbf{z}$.*

Proof. This is basically Sard's theorem, except that z_∞ is nongeneric. The modification is as follows: Let $(u, x, \mathbf{x}') \in \mathcal{M}_J(z_\infty, \mathbf{z})$. Then $ev_*(u, x, \mathbf{x}')$ composed with the projection onto $T_{(z_\infty,0)}X$ is surjective due to two inputs: (i) postcomposition of u by a fractional linear transformation and (ii) variation of the point x. Hence $\mathcal{M}' = \mathcal{M}(X, A, J) \cap \{u(x) = (z_\infty, 0)\}$ is transversely cut out. The rest of the proof follows from applying Sard's theorem to

$$ev' : \mathcal{M}' \to L_{\widehat{a}_1} \times \cdots \times L_{\widehat{a}_{2g}} \times L_{\widehat{a}_1} \times \cdots \times L_{\widehat{a}_{2g}},$$

$$(u, x, \mathbf{x}') \mapsto (u(x'_1), \ldots, u(x'_{4g})).$$

□

2.4.3. *Compactness.*

Lemma 2.4.6. *If $J \in \mathcal{J}^{prod}$, then $\mathcal{M}_J(z_\infty, \mathbf{z})$ is compact for generic $\mathbf{z}$.*

Proof. Let $u_i : F_i \to X$ be a sequence of curves in $\mathcal{M}_J(z_\infty, \mathbf{z})$ that limits to $u : F \to X$. We will eliminate all possible degenerations in the limit to show that $u \in \mathcal{M}_J(z_\infty, \mathbf{z})$. Note in particular that we are assuming that all the points of $\mathbf{z}$ are distinct.

(i) We first claim that u can have no disk bubbles. Arguing by contradiction, suppose that u has a disk bubble u'. Since $\pi_1 \circ u_i|_{int(F_i)}$ is a biholomorphism onto its image for all i, $\mathrm{Im}(u') = \{z\} \times D^2$ for some z. The point z must be an element of $\mathbf{z}$, say z_1, since $\langle u, \Sigma \times \{w\}\rangle = 2g$ for generic $w \in D^2$. On the other hand, if $z = z_1$, then $\langle u, \Sigma \times \{-1\}\rangle > 2g + 1$ unless the image of one component of u is $\Sigma \times \{-1\}$, since the points of $\mathbf{z}$ are distinct. This in turn implies that u has $2g$ disk bubbles that pass through the $2g + 1$ points $(z_\infty, 0), (z_1, 1), \ldots (z_{2g}, 1)$, a contradiction.

(ii) Next we claim that $u(\partial F)$ does not intersect $\{z_\infty\} \times \partial D^2$. Arguing by contradiction, suppose that $u(\partial F) \cap (\{z_\infty\} \times \partial D^2) \neq \varnothing$. Since u cannot have a disk bubble $\{z_\infty\} \times D^2$ by (i), there exists a point $z \in \Sigma - \overline{\mathbf{a}}$ close to z_∞, such u intersects $\{z\} \times D^2$ at two points: once near $(z_\infty, 0)$ and once near $\{z_\infty\} \times \partial D^2$. Hence $\langle u, \{z\} \times D^2\rangle \geq 2$, which is a contradiction. A similar argument also implies that u has no nodes along the boundary: if there are two components of ∂F that map to the same $\widehat{a}_i \times \partial D^2$, then $\langle u, \{z\} \times D^2\rangle \geq 2$, which is a contradiction.

(iii) Finally we eliminate interior nodes. Since u has no disk bubbles and sphere bubbles (the latter easily follows by projecting to D^2), u must have a component $u' : F' \to X$ of genus $< g$ if u has interior nodes. Since $\pi_1 \circ u'|_{int(F')}$ cannot map onto the genus g surface $\Sigma - (\widehat{\mathbf{a}} - N(z_\infty))$, we have a contradiction. $\square$

Lemma 2.4.7. *The count $G_3(\Sigma; J; z_\infty, \mathbf{z})$ is independent of the choice of $J \in \mathcal{J}^{prod}$ and $\mathbf{z}$ generic.*

Proof. We prove the compactness of $\coprod_{t\in[0,1]} \mathcal{M}_{J_t}(z_\infty, \mathbf{z}_t)$ where $(J_t, \mathbf{z}_t)$, $t \in [0, 1]$, is a generic 1-parameter family and we write $\mathbf{z}_t = ((z_1)_t, \dots, (z_{4g})_t)$. The proof is similar to that of Lemma 2.4.6. The only difference is that in Lemma 2.4.6(i), two of the points of $\mathbf{z}_{t_0}$ may come together at isolated $t_0 \in [0, 1]$. Hence it is possible that a limit holomorphic curve u consists of a disk bubble u' and a component $u'' : F'' \to X$ where F'' is a compact surface of genus g with $2g - 1$ boundary components. By a calculation similar to that of Lemma 2.4.4, u'' is regular and $\operatorname{ind}(u'') = 8g - 1$. On the other hand, passing through z_∞ and $4g - 2$ points of $\mathbf{z}_{t_0}$ is a constraint of dimension $4 + 2(4g - 2) = 8g$, which is overdetermined. Hence u must be irreducible, and the proof of Lemma 2.4.6 carries over. $\square$

2.4.4. *Reduction to a torus.* Without loss of generality, assume that $g(\Sigma) = 2$. We explain how to reduce to the case of a torus. As in Section 2.3.4, we degenerate $\Sigma^\tau = (\Sigma, i_\tau)$, $\tau \in [0, \infty)$, into a nodal Riemann surface

$$(\Sigma^\infty, \mathfrak{n}) = ((\Sigma_-, \mathfrak{n}_-) \sqcup (\Sigma_+, \mathfrak{n}_+)) / \mathfrak{n}_- \sim \mathfrak{n}_+$$

by pinching along a separating curve γ. Let $(u^\tau, x^\tau, (\mathbf{x}')^\tau) \in \mathcal{M}_{J_\tau}(z_\infty, \mathbf{z})$ be a sequence where

$$u^\tau : (F_\tau, j_\tau) \to (\Sigma^\tau \times D^2, J_\tau)$$

and J_τ is the product complex structure (i_τ, i_{D^2}). Here we assume the following:

(i) z_∞, $\mathbf{z}$, and $\widehat{\mathbf{a}}$ are fixed for all $\tau \in [0, \infty)$;
(ii) γ is disjoint from $\mathbf{z}, \widehat{a}_1, \widehat{a}_2$, but intersects $\widehat{a}_3, \widehat{a}_4$ at two points each;
(iii) in the limit $z_\infty, z_1, z_2, z_5, z_6 \in \Sigma_+$ and $z_3, z_4, z_7, z_8 \in \Sigma_-$.

Let $\widetilde{\mathbf{a}}$ be the quotient of $\widehat{\mathbf{a}}$ at $\tau = \infty$. Let $v^\tau = \pi_2 \circ u^\tau$, where π_2 is the projection to D^2. We also write $\mathbf{z}_1 = (z_1, z_2, z_5, z_6)$ and $\mathbf{z}_2 = (z_3, z_4, z_7, z_8)$.

We first note that u^τ cannot degenerate as τ approaches a finite τ_0. This is as argued in Lemma 2.4.6.

Let $u = u_- \cup u_0 \cup u_+$ be the limit of the sequence u^τ as $\tau \to \infty$, where $\operatorname{Im}(u_\pm) \subset \Sigma_\pm \times D^2$, $u_\pm$ has no components of $\{\mathfrak{n}\} \times D^2$, and u_0 maps to $\{\mathfrak{n}\} \times D^2$, and let $v_\pm = \pi_2 \circ u_\pm$. By the argument of Lemma 2.4.6(i), disks $\{z\} \times D^2$ cannot bubble off, since otherwise they would introduce extra intersection points with

$$\Sigma^\infty \times \{w\} = ((\Sigma_+ \sqcup \Sigma_-)/\sim) \times \{w\},$$

where $w \in D^2$ is generic. In particular, this implies that $u_0 = \varnothing$.

Let $F_\pm$ be the domain of $u_\pm$. By Lemma 2.4.6(ii), $u_+(\partial F_+)$ does not intersect $\{z_\infty\} \times \partial D^2$. Let $\#(\partial F_\pm)$ be the number of boundary components of $F_\pm$.

We claim that $\#(\partial F_+) = \#(\partial F_-) = 2$. First observe that $\#(\partial F_+) \geq 2$, since two boundary components are needed to map to $\widetilde{a}_i$, $i = 1, 2$; similarly $\#(\partial F_-) \geq$

2. The restriction of v_+ to each component C of ∂F_+ is either a positive degree map $C \to \partial D^2$ or is a constant map to a point $w \in \partial D^2$. If v_+ maps C to a point $w \in \partial D^2$, then u_+ maps an irreducible component of F_+ to a fiber $\Sigma_+ \times \{w\}$. This in turn implies that u_+ has disk bubbles, a contradiction. Hence the restriction of v_+ to each C is a positive degree map. If $\#(\partial F_+) > 2$, then $\deg(v_+|_{\partial F_+}) > 2$. Similarly we obtain that $\deg(v_-|_{\partial F_-}) \geq 2$. This implies that $\deg((v_+ \cup v_-)|_{\partial F}) > 4$, which is a contradiction. Hence $\#(\partial F_+) = \#(\partial F_-) = 2$.

The above claim implies that there exist $x_i, \mathbf{x}'_i$, $i = 1, 2$, such that

$$(u_+, x_1, \mathbf{x}'_1) \in \mathcal{M}_{J_1}(z_\infty, \mathbf{z}_1), \quad (u_-, x_2, \mathbf{x}'_2) \in \mathcal{M}_{J_2}(\mathfrak{n}_-, \mathbf{z}_2).$$

By the usual gluing argument we obtain the following:

Lemma 2.4.8.

$$G_3(\Sigma^\tau; J^\tau; z_\infty, \mathbf{z}) = G_3(\Sigma_+; z_\infty, \mathbf{z}_1) \times G_3(\Sigma_-; \mathfrak{n}_-, \mathbf{z}_2).$$

2.4.5. *Two calculations on* $\mathbb{CP}^1 \times D^2$. We now calculate two model situations which are key ingredients in the proof of Lemma 2.4.12 below. Note that all the holomorphic curves on $\mathbb{CP}^1 \times D^2$ that are considered below satisfy automatic transversality; see Hofer-Lizan-Sikorav [HLS, Theorem 2′].

Fix real numbers $a > b > 0$. Let $\mathcal{S}_1$ be the set of pairs (v_1, w), where v_1 is a degree 1 holomorphic map $D^2 \to \mathbb{CP}^1$ such that $v_1(\partial D^2) \subset \mathbb{R}^+$, $v_1(1) = a$, $v_1(-1) = b$, $v_1(0) = \infty$, and w is a point in D^2 such that $v_1(w) = 0$, and let C_1 be the set of points w for which there is some v_1 with $(v_1, w) \in \mathcal{S}_1$. Similarly, let $\mathcal{S}_2$ be the set of pairs (v_2, w), where v_2 is a degree 1 holomorphic map $D^2 \to \mathbb{CP}^1$ such that $v_2(\partial D^2) \subset \mathbb{R}$, $v_2(1) = a$, $v_2(-1) = b$, $v_2(0) = \infty$, and w is a point on D^2 such that $v_2(w) = -i$, and let C_2 be the set of points w for which there is some v_2 with $(v_2, w) \in \mathcal{S}_2$.

Let $\mathcal{R}_\theta$ be the restriction to D^2 of the radial ray which passes through $0 \in D^2$ and makes an angle of θ with the positive real axis.

Lemma 2.4.9. *There exists* $0 < \theta_0 < \frac{\pi}{2}$ *such that* $C_1 \subset D^2$ *is a curve which is parametrized by* $\theta \in (\pi - \theta_0, \pi + \theta_0)$, $C_1(\theta) = C_1 \cap int(\mathcal{R}_\theta)$, *and*

$$\lim_{\theta \to (\pi+\theta_0)^-} C_1(\theta) = e^{i(\pi+\theta_0)}, \qquad \lim_{\theta \to (\pi-\theta_0)^+} C_1(\theta) = e^{i(\pi-\theta_0)}.$$

See Figure 2 for the curve C_1.

Proof. By the Schwarz reflection principle, a map v_1 with $(v_1, w) \in \mathcal{S}_1$ extends to a degree 2 branched cover $\mathbb{CP}^1 \to \mathbb{CP}^1$ with two branch points which lie on $\mathbb{R}^+ \subset \mathbb{CP}^1$. Hence v_1 admits a factorization

$$D^2 \xrightarrow{f} \mathbb{H} \xrightarrow{\tilde{v}_1} \mathbb{CP}^1 \xrightarrow{g} \mathbb{CP}^1,$$

where $\mathbb{H} = \{\mathrm{Im}(z) \geq 0\}$ is the upper half plane, $f : D^2 \overset{\sim}{\to} \mathbb{H}$ is a fractional linear transformation, $\tilde{v}_1(z) = z^2$, and $g \in PSL(2, \mathbb{R})$. In order for $v_1(0) = \infty$ and $v_1(w) = 0$ to hold, f must map the ray $\mathcal{R}_\theta$ through 0 and w to the line $\{\mathrm{Re}(z) = 0\} \subset \mathbb{H}$. We may set $f(e^{i\theta}) = 0$ and $f(0) = i$; these conditions uniquely determine $f = f_\theta$. We leave it to the reader to verify that for each θ, there



FIGURE 2

is a one-to-one correspondence between $(v_1, w) \in \mathcal{S}_1$ with $w \in \mathcal{R}_\theta$ and $w \in \mathcal{R}_\theta$ satisfying the following equality of cross ratios:

$$(\tilde{v}_1 \circ f_\theta(w), \tilde{v}_1 \circ f_\theta(1); \tilde{v}_1 \circ f_\theta(-1), -1) = (0, b; a, \infty). \tag{2.4.3}$$

Note that $\tilde{v}_1 \circ f_\theta(\mathcal{R}_\theta) = [-1, 0]$ and that there is at most one $w \in \mathcal{R}_\theta$ such that Equation (2.4.3) holds.

When $\theta = \pi$, $\tilde{v}_1 \circ f_\pi(1) = \infty$ and $\tilde{v}_1 \circ f_\pi(-1) = 0$. Hence there is a unique $w \in \mathcal{R}_\pi$ satisfying Equation (2.4.3). As θ moves from π to $\frac{3\pi}{2}$, the points $\tilde{v}_1 \circ f_\theta(1)$ and $\tilde{v}_1 \circ f_\theta(-1)$ approach each other and become equal when $\theta = \frac{3\pi}{2}$. Hence there exists $0 < \theta_0 < \frac{\pi}{2}$ such that there is no $w \in \mathcal{R}_\theta$ for $\pi + \theta_0 < \theta < \frac{3\pi}{2}$ and there is a unique $w \in \mathcal{R}_\theta$ for $\pi \leq \theta < \pi + \theta_0$. The situation of $\theta \in (\frac{\pi}{2}, \pi]$ is symmetric. The lemma then follows. □

Lemma 2.4.10. *There exists a parametrization of $C_2 \subset D^2$ by $\theta \in (\frac{\pi}{2}, \frac{3\pi}{2})$ such that*

$$\lim_{\theta \to \frac{\pi}{2}+} C_2(\theta) = 1, \qquad \lim_{\theta \to \frac{3\pi}{2}-} C_2(\theta) = -1.$$

See Figure 2 for the curve C_2.

Proof. As in the proof of Lemma 2.4.9, v_2 can be factored as

$$D^2 \xrightarrow{f_\theta} \mathbb{H} \xrightarrow{\tilde{v}_2} \mathbb{CP}^1 \xrightarrow{g} \mathbb{CP}^1,$$

where $\tilde{v}_2(z) = z^2$ and f_θ maps 0 to i and $\mathcal{R}_\theta$ to $\{\mathrm{Re}(z) = 0, 0 \leq \mathrm{Im}(z) \leq 1\} \subset \mathbb{H}$. Note that there is a unique fractional linear transformation g such that $g(-1) = \infty$, $g(0) = b$, and $g(\infty) = a$. For each $\theta \in (\frac{\pi}{2}, \frac{3\pi}{2})$, there exists $(v_2, C_2(\theta)) \in \mathcal{S}_2$ such that $C_2(\theta)$ is in one of the half-disks of D^2 divided by $\mathcal{R}_\theta \cup \mathcal{R}_{\theta+\pi}$. As a reference point, $C_2(\pi)$ is in the upper half-disk. As θ approaches $\frac{\pi}{2}$ from above, the corresponding v_2 sends -1 and 1 to arbitrarily close points. Hence $\lim_{\theta \to \frac{\pi}{2}+} C_2(\theta) = 1$. Similarly, $\lim_{\theta \to \frac{3\pi}{2}-} C_2(\theta) = -1$. □

The following lemma is immediate from Lemmas 2.4.9 and 2.4.10; the key ingredient is that the endpoints of C_1 and C_2 alternate on ∂D^2:

Lemma 2.4.11. *The total signed count of intersections between C_1 and C_2 is ± 1.*

2.4.6. *Reduction to* $\mathbb{CP}^1 \times D^2$. We now explain how to further reduce from T^2 to S^2. We pinch T^2 along three parallel, disjoint, essential closed curves $\gamma_1, \gamma_2, \gamma_3$ to obtain a "sausage"

$$\big((\Sigma_1, z_1, z_2') \sqcup (\Sigma_2, z_2, z_3') \sqcup (\Sigma_3, z_3, z_1')\big) \,/\sim, \tag{2.4.4}$$

where $\Sigma_i \simeq S^2$, $i = 1, 2, 3$, and $z_i \sim z_i'$, $i = 1, 2, 3$. More precisely, pick an oriented identification $T^2 \simeq \mathbb{R}^2/\mathbb{Z}^2$ with coordinates (x, y) so that $\overline{a}_1 = \{y = 0\}$ and $\overline{a}_2 = \{x = 0\}$. Then $\gamma_1 = \{x = \frac{1}{4}\}$, $\gamma_2 = \{x = \frac{1}{2}\}$, and γ_3 is obtained from $\{x = \frac{3}{4}\}$ by applying a finger move along the arc $[\frac{3}{4}, 1 + \varepsilon] \times \{\frac{1}{2}\}$ so that γ_3 has two intersections $y = \frac{1}{2} - \varepsilon, \frac{1}{2} + \varepsilon$ with $\widehat{a}_2$. We also assume that $z_1, z_3 \in \widehat{a}_1$ lie on $\{\frac{1}{4} < x < \frac{1}{2}\}$ and $z_2, z_4 \in \widehat{a}_2$ lie on $\{\frac{1}{2} - \varepsilon < y < \frac{1}{2} + \varepsilon\}$. Then Σ_1 is obtained from the closure of the connected component of $T^2 - \cup_{i=1}^3 \gamma_i$ which is bounded by (copies of) γ_1 and γ_2, by identifying all of γ_1 to z_1 and all of γ_2 to z_2'. The other components Σ_2 and Σ_3 are defined similarly. In particular, $z_\infty \in \Sigma_3$. See Figure 3.



FIGURE 3. The top diagram is the torus T^2, where the sides are identified and the top and the bottom are identified. The arrow indicates the projection of T^2 onto $\Sigma_1 \sqcup \Sigma_2 \sqcup \Sigma_3 / \sim$, given in (2.4.4). The components of $\widetilde{a}_i$ which do not contain points in the set $\{z_1, \dots, z_4\}$ are not drawn in Σ_i. Also the two disks on the left and the right of the bottom diagram are glued into Σ_3.

Lemma 2.4.12. *$G_3(T^2; J; z_\infty, \mathbf{z}) = 1$ mod 2, where $\mathbf{z} = \{z_1, \dots, z_4\}$.*

Proof. We degenerate the Riemann surfaces $\Sigma^\tau = (T^2, i_\tau)$, $\tau \to \infty$, by pinching along $\gamma_1 \cup \gamma_2 \cup \gamma_3$. Then a sequence of holomorphic maps

$$u^\tau : (F, j_\tau) \to (\Sigma^\tau \times D^2, J_\tau)$$

in $\mathcal{M}_{J_\tau}(z_\infty, \mathbf{z})$ converges to a cusp curve (u_1, u_2, u_3), where

$$u_i : F_i \to \Sigma_i \times D^2, \quad i = 1, 2, 3,$$

and $F_1 = F_1 = D^2$, $F_3 = \mathbb{CP}^1$. This is because $z_1, z_3 \in \Sigma_1$ and $z_2, z_4 \in \Sigma_2$, and the total number of boundary components $\sum_{i=1}^{3} \#(\partial F_i)$ is equal to two by the argument in Section 2.4.4. Now, u_3 must have image $\Sigma_3 \times \{0\}$. The sets $\mathcal{S}_1$ and $\mathcal{S}_2$ for $v_i = \pi_2 \circ u_i$, $i = 1, 2$, were determined in Section 2.4.5. The gluing of intersecting curves u_1 and u_2 is given by the signed intersection number of C_1 and C_2, which is ± 1 by Lemma 2.4.11. □

3. HOMOTOPY OF COBORDISMS I

In this section and the next we prove Theorem 1.0.1. The chain homotopies that appear in the proof of Theorem 1.0.1 are induced by homotopies of cobordisms $\overline{W}^{\pm}_\tau$ and $\overline{W}^{\mp}_\tau$ which are parametrized by $\tau \in \mathbb{R}$. In this section we treat $\overline{W}^{\pm}_\tau$, leaving $\overline{W}^{\mp}_\tau$ for Section 4. If $\pm$ is understood (as it will be in the rest of this section), then it will be omitted.

We now give a brief description of $\overline{W}_\tau$, leaving precise definitions for later. The base B_τ of $\overline{W}_\tau$ is biholomorphic to an annulus with one puncture on each boundary component; the neighborhoods of the punctures are viewed as strip-like ends. As $\tau \to +\infty$, $\overline{W}_\tau$ degenerates to the stacking of $\overline{W}_+$ "on top of" $\overline{W}_-$. (This is the reason why there is a $\pm$ in $\overline{W}^{\pm}_\tau$.) On the other hand, as $\tau \to -\infty$, $\overline{W}_\tau$ degenerates to $\overline{W}_{-\infty}$, whose base $B_{-\infty}$ is (more or less) given by:

$$B_{-\infty} = ((\mathbb{R} \times [0,1]) \sqcup D) \,/ \sim, \tag{3.0.5}$$

where $D = \{|z| \leq 1\} \subset \mathbb{C}$ and $\sim$ identifies $(0,1) \in \mathbb{R} \times [0,1]$ with $1 \in D$ and $(0,0) \in \mathbb{R} \times [0,1]$ with $-1 \in D$.

3.1. Construction of the homotopy of cobordisms for $\Psi \circ \Phi$.

3.1.1. *Definition of the family* $\overline{W}_\tau$. Let $l, r \in [2, \infty)$. Consider the fibration

$$\pi_r : \mathbb{R} \times \overline{N}_r \to \mathbb{R} \times (\mathbb{R}/r\mathbb{Z}),$$

where

$$\overline{N}_r = (\overline{S} \times [0, r])/(x, r) \sim (\overline{h}(x), 0)$$

and (s, t) are coordinates on $\mathbb{R} \times (\mathbb{R}/r\mathbb{Z})$. Recall that the monodromy map $\overline{h} = \overline{h}_m$ is dependent on the integer $m \gg 0$. We define $\overline{W}_{l,r} = \pi_r^{-1}(B_{l,r})$, where the base $B_{l,r}$ is obtained by smoothing the corners of

$$\{-l \leq s \leq l\} \cup \{0 \leq t \leq 1\} \subset \mathbb{R} \times (\mathbb{R}/r\mathbb{Z}).$$

Next choose a function

$$\eta = (l, r) : \mathbb{R} \to [2, \infty) \times [2, \infty), \tag{3.1.1}$$

which is obtained by smoothing

$$\eta_0(\tau) = \begin{cases} (\tau + 2, 2), & \text{for } \tau \geq 0; \\ (2, 2 - \tau), & \text{for } \tau \leq 0; \end{cases}$$

near $\tau = 0$. We then let $\overline{W}_\tau = \overline{W}_{\eta(\tau)}$ and $B_\tau = B_{\eta(\tau)}$. Let $\pi_{B_\tau} : \overline{W}_\tau \to B_\tau$ be the projection along $\{(s, t)\} \times \overline{S}$.

As $\tau \to +\infty$, the cobordism $\overline{W}_\tau$ approaches the concatenation of $\overline{W}_+$ and $\overline{W}_-$; see Figure 4. On the other hand, as $\tau \to -\infty$, the cobordism $\overline{W}_\tau$ degenerates to a 2-component building $\overline{W}_{-\infty} = \overline{W}_{-\infty,1} \cup \overline{W}_{-\infty,2}$, which we describe now; see Figure 5.

The base of $\overline{W}_{-\infty} = \overline{W}_{-\infty,1} \cup \overline{W}_{-\infty,2}$ is $B_{-\infty} = B_{-\infty,1} \cup B_{-\infty,2}$, where $B_{-\infty,1}$ is obtained from $\{-2 \leq s \leq 2\} \cup \{0 \leq t \leq 1\} \subset \mathbb{R}^2$ by smoothing the corners and $B_{-\infty,2} = [-2,2] \times \mathbb{R}$. Here both $\mathbb{R}^2$ and $[-2,2] \times \mathbb{R}$ have coordinates (s,t). The component $B_{-\infty,1}$ has four strip-like ends: the ends $s \to +\infty$, $s \to -\infty$, $t \to +\infty$, $t \to -\infty$ will be referred to as the top, bottom, left, and right ends. The component $B_{-\infty,2}$ has two strip-like ends: the ends $t \to +\infty$ and $t \to -\infty$ will be referred to as the left and right ends. As usual, $B_{-\infty}$ is endowed with identifications of the compactifications of the strip-like ends. More precisely, if the compactification $\check{B}_{-\infty,1}$ is obtained from $B_{-\infty,1}$ by attaching $\{\pm\infty\} \times [0,1]$ and $[-2,2] \times \{\pm\infty\}$ and the compactification $\check{B}_{-\infty,2}$ is obtained from $B_{-\infty,2}$ by attaching $[-2,2] \times \{\pm\infty\}$, then we identify $(s, \pm\infty) \in \check{B}_{-\infty,1}$ with $(s, \mp\infty) \in \check{B}_{-\infty,2}$.

Let $\overline{W}_{-\infty,i} = B_{-\infty,i} \times \overline{S}$ for $i = 1,2$. The 2-component building $\overline{W}_{-\infty} = \overline{W}_{-\infty,1} \cup \overline{W}_{-\infty,2}$ is endowed with identifications of compactifications of the ends. The compactification $\check{\overline{W}}_{-\infty,1}$ is obtained from $\overline{W}_{-\infty,1}$ by attaching $\{\pm\infty\} \times [0,1] \times \overline{S}$ and $[-2,2] \times \{\pm\infty\} \times \overline{S}$, the compactification $\check{\overline{W}}_{-\infty,2}$ is obtained from $\overline{W}_{-\infty,2}$ by attaching $[-2,2] \times \{\pm\infty\} \times \overline{S}$, and we identify $(s, +\infty, x) \in \check{\overline{W}}_{-\infty,1}$ with $(s, -\infty, x) \in \check{\overline{W}}_{-\infty,2}$ and $(s, +\infty, x) \in \check{\overline{W}}_{-\infty,2}$ with $(s, -\infty, \overline{h}(x)) \in \check{\overline{W}}_{-\infty,1}$. We write $\pi_{B_{-\infty,i}} : \overline{W}_{-\infty,i} \to B_{-\infty,i}$ for the projection along $\overline{S}$.



FIGURE 4. The bases of the family $\overline{W}_\tau$. The parameter τ increases as we go to the right. The sides are identified in this picture, as indicated. The location of $\overline{\mathfrak{m}}^b(\tau)$ is indicated by ×.

We write $cl(B_\tau)$, $cl(B_+)$, $cl(B_-)$ to denote the compactifications of B_τ, B_+, B_-, obtained by adjoining the points at infinity $\pm\infty$. Similarly, we write $cl(B_{-\infty,1})$ for the compactification of $B_{-\infty,1}$, obtained by adding 4 points $s = \pm\infty$ and $t = \pm\infty$, and $cl(B_{-\infty,2})$ for the compactification of $B_{-\infty,2}$, obtained by adding 2 points $t = \pm\infty$.



FIGURE 5. The bases of $\overline{W}_{-\infty,2}$ to the left and $\overline{W}_{-\infty,1}$ to the right.

3.1.2. *Marked points.* We choose a 1-parameter family of marked points

$$\overline{\mathfrak{m}}(\tau)=(\overline{\mathfrak{m}}^b(\tau),\overline{\mathfrak{m}}^f(\tau))=((-l(\tau)+2,(r(\tau)+1)/2),z_\infty)\in\overline{W}_\tau$$

for $\tau\in\mathbb{R}$, such that the following hold:

(i) as $\tau\to+\infty$, $\overline{\mathfrak{m}}(\tau)$ limits to $\overline{\mathfrak{m}}(+\infty)=(\overline{\mathfrak{m}}^b(+\infty),\overline{\mathfrak{m}}^f(+\infty))$, where $\overline{\mathfrak{m}}^b(+\infty)=(0,\frac{3}{2})\in B_-$ and $\overline{\mathfrak{m}}^f(-\infty)=z_\infty$;

(ii) as $\tau\to-\infty$, $\overline{\mathfrak{m}}(\tau)$ limits to $\overline{\mathfrak{m}}(-\infty)=(\overline{\mathfrak{m}}^b(-\infty),\overline{\mathfrak{m}}^f(-\infty))$, where $\overline{\mathfrak{m}}^b(-\infty)=(0,0)\in B_{-\infty,2}$ and $\overline{\mathfrak{m}}^f(-\infty)=z_\infty$.

Convention 3.1.1. In this section and the next, $\overline{\mathfrak{m}}$ will denote a 1-parameter family (as opposed to a single marked point).

Let us write $\mathcal{L}_{t_0}$ for the locus $\{t=t_0\}$, viewed as a subset of B_+, B_-, B_τ, as appropriate. Of particular interest is $\mathcal{L}_{(r(\tau)+1)/2}$, which passes through $\overline{\mathfrak{m}}^b(\tau)\in B_\tau$ or $\overline{\mathfrak{m}}^b(+\infty)\in B_-$.

3.1.3. *Stable Hamiltonian structures and symplectic forms.* We first consider $\overline{W}_\tau$. Let $\overline{\omega}$ be the area form on $\overline{S}$ from Section I.5.1.2. Then the stable Hamiltonian structure on $\overline{N}_{r(\tau)}=(\overline{S}\times[0,r(\tau)])/\sim$ is obtained from $(dt,\overline{\omega})$ on $\overline{S}\times[0,r(\tau)]$ by passing to the quotient. The 2-plane field is $\xi_\tau=\ker dt=T\overline{S}$ and the Hamiltonian vector field is $\overline{R}_\tau=\partial_t$. The symplectic form $\overline{\Omega}_\tau$ is obtained from the symplectic form $ds\wedge dt+\overline{\omega}$ on $\mathbb{R}\times\overline{S}\times[0,r(\tau)]$ by passing to the quotient $\mathbb{R}\times\overline{N}_{r(\tau)}$ and then restricting to $\overline{W}_\tau$.

Next we consider $\overline{W}_{-\infty}=\overline{W}_{-\infty,1}\cup\overline{W}_{-\infty,2}$. Let $\omega_{-\infty,1}$ be the restriction of the area form $ds\wedge dt$ on $\mathbb{R}^2$ to $B_{-\infty,1}$ and let $\omega_{-\infty,2}=ds\wedge dt$ on $B_{-\infty,2}=[-2,2]\times\mathbb{R}$. Then we set $\overline{\Omega}_{-\infty,i}=\omega_{-\infty,i}+\overline{\omega}$. The stable Hamiltonian structure at the $s\to\pm\infty$ ends of $\overline{W}_{-\infty,1}$ are given by $(dt,\overline{\omega})$ and the stable Hamiltonian structure at the $t\to\pm\infty$ ends of $\overline{W}_{-\infty,1}$ are given by $(ds,\overline{\omega})$.

3.1.4. *The choice of hyperbolic orbit h.* Recall that ∂N is a negative Morse-Bott torus. Without loss of generality, we may assume that there is only one holomorphic cylinder Z_γ in $(\mathbb{R}\times\overline{N},\overline{J}')$ from δ_0 to any orbit γ of ∂N, modulo $\mathbb{R}$-translation. Each Z_γ corresponds to a radial ray $\mathcal{R}_{\phi_\gamma}$, which is the asymptotic direction of

$\pi_{D^2}(Z_\gamma)$ at the positive end. Here $\pi_{D^2}: \overline{N} - int(N) \to D^2$ is the projection with respect to the balanced coordinates; see Section I.5.1.2.

The choice of the hyperbolic orbit $h = \gamma_{\phi_h}$ in ∂N is the same as that of Convention I.6.6.4: h is generic and ϕ_h is close to $-\frac{2\pi}{m}$, where m is as defined in Section I.5.2.2. In particular, the radial ray $\mathcal{R}_{\phi_h}$ does not lie on the thin wedges from $\overline{a}_i$ to $\overline{h}(\overline{a}_i)$ for all i. There are no restrictions on the orbit e except that $e \neq h$.

3.2. **Holomorphic curves and moduli spaces.**

3.2.1. *Lagrangian boundary conditions.* Recall that $\overline{\mathbf{a}} = \{\overline{a}_1, \dots, \overline{a}_{2g}\}$ is the extension of the basis $\mathbf{a}$ to $\overline{S}$ and that the monodromy map $\overline{h} = \overline{h}_m : \overline{S} \to \overline{S}$ depends on the integer $m \gg 0$.

We first describe the pushoff $\overline{\mathbf{b}}$ of $\overline{\mathbf{a}}$: Let $\varepsilon_0 = \varepsilon_0(m) > 0$ be sufficiently small and let $\overline{b}_i$ be a ε_0-close transverse pushoff of $\overline{a}_i$ which satisfies the following:

- in a neighborhood of z_∞, $\overline{b}_i$ is obtained from $\overline{a}_i$ by a $-\frac{2\pi}{mK(m)}$-rotation, where $K(m)$ is a positive integer such that $\lim_{m\to\infty} K(m) = \infty$; and
- $\overline{a}_i$ and $\overline{b}_i$ intersect at three points $x^\#_{i1}$, $x^\#_{i2}$ and $x^\#_{i3}$ (besides at z_∞); $x^\#_{i2} \in int(S)$ and $x^\#_{i1}, x^\#_{i3} \in \overline{S} - S$.

See Figure 6. When we want to signal that z_∞ is an intersection point of $\overline{a}_i$ and $\overline{b}_i$, then we write it as $z^\#_\infty$.

Let us write $\phi(\overline{a}_{i,j})$ for the ϕ-coordinate of $\overline{a}_{i,j}$, subject to the condition $0 \leq \phi(\overline{a}_{i,j}) < 2\pi$; similarly define $\phi(\overline{b}_{i,j})$, etc. We additionally assume that

$$0 < \phi(\overline{a}_{i,j}), \phi(\overline{b}_{i,j}), \phi(\overline{h}(\overline{a}_{i,j})), \phi(\overline{h}(\overline{b}_{i,j})) \leq c(m)$$

where $c(m) \to 0$ as $m \to \infty$; cf. Section I.5.2.2.



FIGURE 6. The arcs $\overline{a}_i$, $\overline{b}_i$, $\overline{h}(\overline{a}_i)$, and $\overline{h}(\overline{b}_i)$ near z_∞.

Remark 3.2.1. In view of the choices of $\overline{a}_{i,j}$ and $\overline{h}(\overline{a}_{i,j})$ from Section I.5.2.2 and the choices of $\overline{b}_{i,j}$ above, as $m \to \infty$:

- $\phi(\overline{a}_{i,j}) - \phi(\overline{b}_{i,j}) \to 0$ the fastest;
- $\phi(\overline{h}(\overline{a}_{i,j})) - \phi(\overline{a}_{i,j}) \to 0$ the next fastest;
- $\phi(\overline{a}_{i,j}) - \phi(\overline{a}_{i',j'}) \to 0$ the slowest if $(i,j) \neq (i',j')$.

The symplectic fibration

$$\pi_{B_\tau}:(\overline{W}_\tau,\overline{\Omega}_\tau)\to(B_\tau,ds\wedge dt)$$

induces a symplectic connection, defined as the $\overline{\Omega}_\tau$-orthogonal of the tangent plane to the fibers. We place a copy of $\overline{\mathbf{a}}$ on the fiber $\pi_{B_\tau}^{-1}(s,1)$ with $s>l(\tau)$ and use the symplectic connection to parallel transport $\overline{\mathbf{a}}$ along $\partial\overline{W}_\tau$. (Note that $\partial\overline{W}_\tau$ is equal to the *vertical boundary* $\partial_v\overline{W}_\tau:=\pi_{B_\tau}^{-1}(\partial B_\tau)$.) This gives us a singular Lagrangian submanifold $L^{\tau,+}_{\overline{\mathbf{a}}}$. Similarly, we place a copy of $\overline{\mathbf{b}}$ on the fiber $\pi_{B_\tau}^{-1}(s,1)$ with $s<-l(\tau)$ and use the symplectic connection to parallel transport $\overline{\mathbf{b}}$ along $\partial\overline{W}_\tau$ to construct the singular Lagrangian submanifold $L^{\tau,-}_{\overline{\mathbf{b}}}$. The Lagrangian submanifolds $L^{\tau,+}_{\widehat{\mathbf{a}}}$, $L^{\tau,+}_{\widehat{a}_i}$, $L^{\tau,-}_{\widehat{\mathbf{b}}}$, $L^{\tau,-}_{\widehat{b}_i}$, etc. are defined similarly.

On $(\overline{W}_{-\infty},\overline{\Omega}_{-\infty})$, we define the singular Lagrangian submanifolds as follows: Let us write

$$\partial B_{-\infty,1}=\sqcup_{i=1}^4\partial_i B_{-\infty,1},$$

where the boundary components, in order from $i=1$ to $i=4$, satisfy

$$s>0,t>1/2;\quad s<0,t>1/2;\quad s<0,t<1/2;\quad \text{and } s>0,t<1/2.$$

Also let $\partial_i\overline{W}_{-\infty,1}$ be the component of $\partial\overline{W}_{-\infty,1}$ corresponding to $\partial_i B_{-\infty,1}$. Then we define:

- $L^{-\infty,1}_{\overline{\mathbf{a}},1}=\partial_1 B_{-\infty,1}\times\overline{\mathbf{a}}$ on $\partial_1\overline{W}_{-\infty,1}$;
- $L^{-\infty,1}_{\overline{\mathbf{b}},2}=\partial_2 B_{-\infty,1}\times\overline{\mathbf{b}}$ on $\partial_2\overline{W}_{-\infty,2}$;
- $L^{-\infty,1}_{\overline{h}(\overline{\mathbf{b}}),3}=\partial_3 B_{-\infty,1}\times\overline{h}(\overline{\mathbf{b}})$ on $\partial_3\overline{W}_{-\infty,1}$;
- $L^{-\infty,1}_{\overline{h}(\overline{\mathbf{a}}),4}=\partial_4 B_{-\infty,1}\times\overline{h}(\overline{\mathbf{a}})$ on $\partial_4\overline{W}_{-\infty,1}$;
- $L^{-\infty,2}_{\overline{\mathbf{a}},+}=\{2\}\times\mathbb{R}\times\overline{\mathbf{a}}$ on $\partial\overline{W}_{-\infty,2}$;
- $L^{-\infty,2}_{\overline{\mathbf{b}},-}=\{-2\}\times\mathbb{R}\times\overline{\mathbf{b}}$ on $\partial\overline{W}_{-\infty,2}$.

3.2.2. *Almost complex structures.* Recall the space $\overline{\mathcal{J}}$ of admissible almost complex structures $\overline{J}$ on $\overline{W}$ from Definition I.5.3.2, the space $\overline{\mathcal{J}'}$ of adapted almost complex structures $\overline{J'}$ on $\overline{W'}$ from Definition I.5.3.9, and the spaces $\overline{\mathcal{J}}_+$ and $\overline{\mathcal{J}}_-$ of admissible almost complex structures $\overline{J}_+$ and $\overline{J}_-$ on $\overline{W}_+$ and $\overline{W}_-$ from Definition I.5.4.1.

Definition 3.2.2. An almost complex structure $\overline{J}_{-\infty,2}$ on $\overline{W}_{-\infty,2}$ is *admissible* if the following hold:

1. $\overline{J}_{-\infty,2}$ is t-invariant, $\overline{J}_{-\infty,2}(\partial_t)=-\partial_s$, and $\overline{J}_{-\infty,2}(T\overline{S})=T\overline{S}$; and
2. there exists $\varepsilon>0$ such that $\overline{J}_{-\infty,2}$ restricts to the standard complex structure on the subsurface $D_\varepsilon=\{\rho\le\varepsilon\}\subset\overline{S}$ of each fiber.

The space of all admissible $\overline{J}_{-\infty,2}$ will be denoted by $\overline{\mathcal{J}}_{-\infty,2}$.

Definition 3.2.3. An almost complex structure $\overline{J}_{-\infty,1}$ on $\overline{W}_{-\infty,1}$ is *admissible* if the following hold:

(1) there is a complex structure $j_{-\infty,1}$ on $B_{-\infty,1}$ such that the projection $\pi_{B_{-\infty,1}}$ is $(\overline{J}_{-\infty,1}, j_{-\infty,1})$-holomorphic;
(2) there exists $\varepsilon > 0$ such that $\overline{J}_{-\infty,1}$ restricts to the standard complex structure on the subsurface $D_\varepsilon = \{\rho \leq \varepsilon\} \subset \overline{S}$ of each fiber;
(3) there exist $\overline{J} \in \overline{\mathcal{J}}$ and $\overline{J}_{-\infty,2} \in \overline{\mathcal{J}}_{-\infty,2}$ such that $\overline{J}_{-\infty,1}$ agrees with $\overline{J}$ on $\overline{W} = \mathbb{R} \times [0,1] \times \overline{S}$, with $\overline{J}_{-\infty,2}$ on $[-2,2] \times [3,+\infty) \times \overline{S}$, and with $(id \times \overline{h})_*(\overline{J}_{-\infty,2})$ on $[-2,2] \times [-2,-\infty) \times \overline{S}$.

If (3) holds, we say that $\overline{J}_{-\infty,1}$ is *compatible* with $\overline{J} \in \overline{\mathcal{J}}$ and $\overline{J}_{-\infty,2} \in \overline{\mathcal{J}}_{-\infty,2}$. The space of admissible $\overline{J}_{-\infty,1}$ will be denoted by $\overline{\mathcal{J}}_{-\infty,1}$.

Definition 3.2.4. An almost complex structure $\overline{J}_{-\infty} = \overline{J}_{-\infty,1} \cup \overline{J}_{-\infty,2}$ on $\overline{W}_{-\infty} = \overline{W}_{-\infty,1} \cup \overline{W}_{-\infty,2}$ is *admissible* if $\overline{J}_{-\infty,i} \in \overline{\mathcal{J}}_{-\infty,i}$ for $i = 1,2$ and $\overline{J}_{-\infty,1}$ is compatible with $\overline{J}_{-\infty,2}$. The space of admissible $\overline{J}_{-\infty}$ will be denoted by $\overline{\mathcal{J}}_{-\infty}$.

Definition 3.2.5. An almost complex structure $\overline{J}_\tau$ on $\overline{W}_\tau$ is *admissible* if the following hold:

(1) there is a complex structure j_τ on B_τ such that the projection π_{B_τ} is $(\overline{J}_\tau, j_\tau)$-holomorphic;
(2) there exists $\varepsilon > 0$ such that $\overline{J}_\tau$ restricts to the standard complex structure on the subsurface $D_\varepsilon = \{\rho \leq \varepsilon\} \subset \overline{S}$ of each fiber;
(3) if $\tau \geq 0$, then $\overline{J}_\tau$ is the restriction of some $\overline{J'} \in \overline{\mathcal{J}'}$;
(4) if $\tau \leq 0$, then $\overline{J}_\tau$ agrees with some $\overline{J} \in \overline{\mathcal{J}}$ on $\overline{W} = \mathbb{R} \times [0,1] \times \overline{S}$ and with some $\overline{J}_{-\infty,2} \in \overline{\mathcal{J}}_{-\infty,2}$ on $[-2,2] \times [3, r(\tau) - 2] \times \overline{S}$.

If (3) holds, we say that $\overline{J}_\tau$ is *compatible* with $\overline{J'} \in \overline{\mathcal{J}'}$ and if (4) holds, we say that $\overline{J}_\tau$ is *compatible* with $\overline{J} \in \overline{\mathcal{J}}$ and $\overline{J}_{-\infty,2} \in \overline{\mathcal{J}}_{-\infty,2}$. The space of all admissible $\overline{J}_\tau$ on $\overline{W}_\tau$ will be denoted by $\overline{\mathcal{J}}_\tau$.

Definition 3.2.6. A family $\{\overline{J}_\tau \in \overline{\mathcal{J}}_\tau\}_{\tau\in\mathbb{R}}$ of almost complex structures is *admissible* if there exist $\overline{J'} \in \overline{\mathcal{J}'}$, $\overline{J} \in \overline{\mathcal{J}}$, $\overline{J}_+ \in \overline{\mathcal{J}}_+$, $\overline{J}_- \in \overline{\mathcal{J}}_-$ and $\overline{J}_{-\infty} = \overline{J}_{-\infty,1} \cup \overline{J}_{-\infty,2} \in \overline{\mathcal{J}}_{-\infty}$ such that the following hold:

(1) $\overline{J}_\tau$ converges to $\overline{J}_{-\infty}$ as $\tau \to -\infty$;
(2) $\overline{J}_\tau$ converges to $\overline{J}_+$ and $\overline{J}_-$ as $\tau \to +\infty$;
(3) $\overline{J}_+$ and $\overline{J}_-$ are compatible with $\overline{J'}$ and $\overline{J}$; and
(4) $\overline{J}_\tau$ is compatible with $\overline{J}$ and $\overline{J}_{-\infty,2}$ for $\tau \leq 0$ and with $\overline{J'}$ for $\tau \geq 0$.

The space of all admissible $\{\overline{J}_\tau \in \overline{\mathcal{J}}_\tau\}_{\tau\in\mathbb{R}}$ will be denoted by $\overline{\mathcal{I}}$.

3.2.3. *Some notation and conventions.* We now collect some notation and conventions.

Convention. When we write $\mathbf{y}$ or γ (with possible superscripts, subscripts and other decorations), it is assumed that $\mathbf{y} \subset S$ and $\gamma \subset N$. In particular, $\mathbf{y}$ and γ do not contain any multiples of z_∞ or δ_0.

Sections at ∞. The sections $\{\rho = 0\}$ of $\overline{W}$, $\overline{W'} = \mathbb{R} \times \overline{N}$, $\overline{W}_\tau$, $\overline{W}_+$, $\overline{W}_-$ and $\overline{W}_{-\infty,i}$ are holomorphic with respect to almost complex structures in $\overline{\mathcal{J}}$, $\overline{\mathcal{J}'}$, $\overline{\mathcal{J}}_\tau$, etc. They are called *sections at* ∞ and are denoted by σ_∞, σ'_∞, σ^τ_∞, σ^+_∞, σ^-_∞ and $\sigma^{-\infty,i}_\infty$.

The intersection numbers $n^*(\overline{u})$ *and* $n^{*,alt}(\overline{u})$. Let δ_{ρ_0,ϕ_0} be a closed orbit of the Hamiltonian vector field ∂_t which lies on the torus $\{\rho=\rho_0\}\subset \overline{S}\times[0,r]/\sim$ for appropriate r and $\rho_0>0$ sufficiently small and which passes through the point $(t,\phi)=(0,\phi_0)$. Here we are using balanced coordinates on $\overline{N}_r$. We assume additionally that δ_{ρ_0,ϕ_0} does not intersect the projections of the Lagrangians of $\overline{W}_\tau$, $\overline{W}_+$ and $\overline{W}_-$ to $\overline{N}_r$.

Let us write $\phi(\overline{a}_{i,j})$ for the ϕ-coordinate of $\overline{a}_{i,j}$ such that $0\leq \phi(\overline{a}_{i,j})<2\pi$. Also let $\varepsilon_0=\frac{2\pi}{Km}$ be the constant appearing in the definition of $\overline{\mathbf{b}}$ and let ε_1 be a constant satisfying $0<\varepsilon_1<\varepsilon_0$. We consider two possibilities for ϕ_0:

$$\phi_0^\pm=\phi(\overline{a}_{i,j})\pm\varepsilon_1.$$

Hence $\phi(\overline{b}_{i,j})<\phi_0^-<\phi(\overline{a}_{i,j})<\phi_0^+$.

We write $(\sigma^*_\infty)^{\dagger,\pm}$ for the restriction of $\mathbb{R}\times\delta_{\rho_0,\phi_0^\pm}$ to $\overline{W}_*$, where $*=\varnothing,{}'$, τ, $+$, or $-$. For $\overline{W}_{-\infty,i}$, we write

$$(\sigma^{-\infty,i}_\infty)^{\dagger,\pm}=B_{-\infty,i}\times\{\rho=\rho_0,\phi=\phi_0^\pm+2\pi k/m,k\in\mathbb{Z}\},$$

for some ϕ_0. Finally we define:

$$n^*(\overline{u})=\langle\overline{u},(\sigma^*_\infty)^{\dagger,+}\rangle,\quad n^{*,alt}(\overline{u})=\langle\overline{u},(\sigma^*_\infty)^{\dagger,-}\rangle, \tag{3.2.1}$$

where $*=\varnothing,{}'$, τ, $+$, $-$, or $(-\infty,i)$. The two quantities n^* and $n^{*,alt}$ can be used interchangeably, except when $\tau=-\infty$; for the most part we will use n^*.

Notation. Given a holomorphic curve $\overline{u}$ in $\overline{W}$, $\overline{W'}$, etc., we write

$$\overline{u}=\overline{u}'\cup\overline{u}''=\overline{u}'\cup\overline{u}^\sharp\cup\overline{u}^\flat\cup\overline{u}^f,$$

where

- $\overline{u}'$ is a possibly disconnected branched cover of σ^*_∞;
- $\overline{u}''$ is the union of irreducible components which do not branch cover σ^*_∞;
- $\overline{u}^\sharp$ is the union of components of $\overline{u}''$ which are asymptotic to a multiple of δ_0 or z_∞ at one or more ends;
- $\overline{u}^\flat$ is the union of the remaining non-fiber components of $\overline{u}''$; and
- $\overline{u}^f$ is the union of fiber components of $\overline{u}$, including ghosts.

If $\overline{u}$ is a multisection, then $\deg\overline{u}$ is the degree of $\overline{u}$ as a multisection.

3.2.4. *Holomorphic maps to* $\overline{W}_\tau$. Let (F,j) be a compact Riemann surface, possibly disconnected, with two k-tuples of boundary punctures $\mathbf{q}^+=\{q_1^+,\dots,q_k^+\}$ and $\mathbf{q}^-=\{q_1^-,\dots,q_k^-\}$ on $\partial F=\partial_+F\sqcup\partial_-F$, such that:

(i) each component of F nontrivially intersects ∂_+F and ∂_-F;
(ii) each of ∂_+F and ∂_-F is a union of connected components of ∂F; and
(iii) on each component of ∂_+F (resp. ∂_-F) there is at least one puncture from $\mathbf{q}^+$ (resp. $\mathbf{q}^-$) and none from $\mathbf{q}^-$ (resp. $\mathbf{q}^+$).

We write $\dot{F} = F - \mathbf{q}^+ - \mathbf{q}^-$, $\partial_+\dot{F} = \partial_+ F - \mathbf{q}^+$ and $\partial_-\dot{F} = \partial_- F - \mathbf{q}^-$.

Let $\mathbf{z} = \{z_\infty^p(\overrightarrow{\mathcal{D}})\} \cup \mathbf{y}$, $p \geq 0$, be a k-tuple of points of $\overline{\mathbf{a}} \cap \overline{h}(\overline{\mathbf{a}})$, where $k \leq 2g$, z_∞ has multiplicity p, and $\overrightarrow{\mathcal{D}}$ is the data at z_∞^p with respect to $\overline{\mathbf{a}} \cap \overline{h}(\overline{\mathbf{a}})$. (Recall that each arc of $\{\overline{a}_i, \overline{h}(\overline{a}_i)\}_{i=1}^{2g}$ is used at most once.) Also let $\mathbf{z}' = \{z_\infty^q(\overrightarrow{\mathcal{D}}')\} \cup \mathbf{y}'$, $q \geq 0$, be a k-tuple of points of $\overline{\mathbf{b}} \cap \overline{h}(\overline{\mathbf{b}})$, where z_∞ has multiplicity q and $\overrightarrow{\mathcal{D}}'$ is the data at z_∞^q with respect to $\overline{\mathbf{b}} \cap \overline{h}(\overline{\mathbf{b}})$.

Let $\overline{J}_\tau \in \overline{\mathcal{J}}_\tau$. If $\overline{u}' : \dot{F} \to (\overline{W}_\tau, \overline{J}_\tau)$ is a branched cover of σ_∞^τ, then $\overline{u}'$ comes equipped with data $\mathcal{C}$, which is a map from $\pi_0(\partial_+\dot{F})$ (resp. $\pi_0(\partial_-\dot{F})$) to the set of arcs $\overline{a}_{i,j}$ (resp. $\overline{b}_{i,j}$). In words, $\overline{u}'$ is viewed as mapping each component of $\partial_+\dot{F}$ (resp. $\partial_-\dot{F}$) to some $L^{\tau,+}_{\overline{a}_{i,j}}$ (resp. $L^{\tau,-}_{\overline{b}_{i,j}}$). Then $\mathcal{C}$ determines the data $\overrightarrow{\mathcal{D}}$ and $\overrightarrow{\mathcal{D}}'$ at the positive and negative ends.

We then make the following definition:

Definition 3.2.7. Let $\overline{J}_\tau \in \overline{\mathcal{J}}_\tau$, $\mathbf{z} = \{z_\infty^p(\overrightarrow{\mathcal{D}})\} \cup \mathbf{y}$ be a k-tuple of $\overline{\mathbf{a}} \cap \overline{h}(\overline{\mathbf{a}})$ and $\mathbf{z}' = \{z_\infty^q(\overrightarrow{\mathcal{D}}')\} \cup \mathbf{y}'$ be a k-tuple of $\overline{\mathbf{b}} \cap \overline{h}(\overline{\mathbf{b}})$.

A *degree k multisection of $(\overline{W}_\tau, \overline{J}_\tau)$ from $\mathbf{z}$ to $\mathbf{z}'$* is a pair $(\overline{u}, \mathcal{C})$ consisting of a holomorphic map

$$\overline{u} = \overline{u}' \cup \overline{u}'' : (\dot{F} = \dot{F}' \sqcup \dot{F}'', j) \to (\overline{W}_\tau, \overline{J}_\tau)$$

which is a degree k multisection of $\pi_{B_\tau} : \overline{W}_\tau \to B_\tau$ and data $\mathcal{C}$ for $\overline{u}'$, and which additionally satisfies the following:

1. $\overline{u}''(\partial_+\dot{F}'') \subset L^{\tau,+}_{\widehat{\mathbf{a}}}$ and $\overline{u}''(\partial_-\dot{F}'') \subset L^{\tau,-}_{\widehat{\mathbf{b}}}$;
2. $\overline{u}$ maps each connected component of $\partial_+\dot{F}$ to a different $L^{\tau,+}_{\overline{a}_i}$ and each connected component of $\partial_-\dot{F}$ to a different $L^{\tau,-}_{\overline{b}_i}$ (here we are using $\mathcal{C}$ to assign some $L^{\tau,+}_{\overline{a}_i}$ or $L^{\tau,-}_{\overline{b}_i}$ to each component of $\partial_\pm\dot{F}'$);
3. $\lim_{w \to q_i^+} \pi_\mathbb{R} \circ \overline{u}(w) = +\infty$ and $\lim_{w \to q_i^-} \pi_\mathbb{R} \circ \overline{u}(w) = -\infty$;
4. $\overline{u}$ converges to a strip over $[0,1] \times \mathbf{z}$ near $\mathbf{q}^+$ and to a strip over $[0,1] \times \mathbf{z}'$ near $\mathbf{q}^-$;
5. the positive and negative ends of $\overline{u}$ which limit to z_∞ are described by $\overrightarrow{\mathcal{D}}$ and $\overrightarrow{\mathcal{D}}'$.

Here $\pi_\mathbb{R} : \overline{W}_\tau \to \mathbb{R}$ is the projection to the s-coordinate.

A *$(\overline{W}_\tau, \overline{J}_\tau)$-curve from $\mathbf{y}$ to $\mathbf{y}'$* is a degree $2g$ multisection of $(\overline{W}_\tau, \overline{J}_\tau)$ satisfying $n^*(\overline{u}) = m$.

Let $\mathcal{M}_{\overline{J}_\tau}(\mathbf{z}, \mathbf{z}')$ be the moduli space of degree k multisections of $(\overline{W}_\tau, \overline{J}_\tau)$ from $\mathbf{z}$ to $\mathbf{z}'$ and let $\mathcal{M}_{\overline{J}_\tau}(\mathbf{z}, \mathbf{z}'; \overline{\mathfrak{m}}(\tau)) \subset \mathcal{M}_{\overline{J}_\tau}(\mathbf{z}, \mathbf{z}')$ be the subset consisting of $\overline{W}_\tau$-curves that pass through $\overline{\mathfrak{m}}(\tau)$. We write

$$\mathcal{M}_{\{\overline{J}_\tau\}}(\mathbf{z}, \mathbf{z}') := \coprod_{\tau \in \mathbb{R}} \mathcal{M}_{\overline{J}_\tau}(\mathbf{z}, \mathbf{z}'),$$

$$\mathcal{M}_{\{\overline{J}_\tau\}}(\mathbf{z},\mathbf{z}';\overline{\mathfrak{m}}) := \coprod_{\tau\in\mathbb{R}} \mathcal{M}_{\overline{J}_\tau}(\mathbf{z},\mathbf{z}';\overline{\mathfrak{m}}(\tau)).$$

Modifiers. For any moduli space $\mathcal{M}_{\star_1}(\star_2)$, we may place modifiers $*$ as in $\mathcal{M}^*_{\star_1}(\star_2)$ to denote the subset of $\mathcal{M}_{\star_1}(\star_2)$ satisfying $*$. Typical self-explanatory modifiers are $I=i$, $n^*=m$, and $\deg = k$. Note that the degree can be inferred from $\star_2$.

The following is a list of non-self-explanatory modifiers:

$\dagger$ = no component of $\overline{u}$ branch covers σ^*_∞ with possibly empty branch locus.
s = all the components of $\overline{u}$ are simply covered.
irr = the curve $\overline{u}$ is irreducible.

3.2.5. *Holomorphic maps to $\overline{W}_{-\infty}$.* We first discuss holomorphic curves without ends at z_∞.

Definition 3.2.8. Let $\overline{J}_{-\infty,1}\in\overline{\mathcal{J}}_{-\infty,1}$, $\mathbf{y}_1\in\mathcal{S}_{\mathbf{a},h(\mathbf{a})}$, $\mathbf{y}_2\in\mathcal{S}_{\mathbf{b},\mathbf{a}}$, $\mathbf{y}_3\in\mathcal{S}_{\mathbf{b},h(\mathbf{b})}$ and $\mathbf{y}_4\in\mathcal{S}_{h(\mathbf{a}),h(\mathbf{b})}$.

A *degree $k\le 2g$ multisection $\overline{u}$ of $(\overline{W}_{-\infty,1},\overline{J}_{-\infty,1})$ with ends* $\mathbf{y}_1,\mathbf{y}_2,\mathbf{y}_3,\mathbf{y}_4$ is a holomorphic map

$$\overline{u}:(\dot{F},j)\to(\overline{W}_{-\infty,1},\overline{J}_{-\infty,1})$$

which is degree k multisection of $\pi_{B_{-\infty,1}}:\overline{W}_{-\infty,1}\to B_{-\infty,1}$ and which additionally satisfies the following:

(1) $\overline{u}(\partial\dot{F})\subset L^{-\infty,1}_{\overline{\mathbf{a}},1}\cup L^{-\infty,1}_{\overline{\mathbf{b}},2}\cup L^{-\infty,1}_{\overline{h(\mathbf{b})},3}\cup L^{-\infty,1}_{\overline{h(\mathbf{a})},4}$ and $\overline{u}$ maps each component of $\partial\dot{F}$ to a different $L^{-\infty,1}_{\overline{a}_i,1}$, $L^{-\infty,1}_{\overline{b}_i,2}$, $L^{-\infty,1}_{\overline{h(\overline{b}_i)},3}$, or $L^{-\infty,1}_{\overline{h(\overline{a}_i)},4}$;
(2) $\overline{u}$ converges to a strip over $[0,1]\times\mathbf{y}_1$ as $s\to+\infty$; $[-2,2]\times\mathbf{y}_2$ as $t\to+\infty$; $[0,1]\times\mathbf{y}_3$ as $s\to-\infty$; and $[-2,2]\times\mathbf{y}_4$ as $t\to+\infty$.

A $(\overline{W}_{-\infty,1},\overline{J}_{-\infty,1})$*-curve with ends* $\mathbf{y}_1,\dots,\mathbf{y}_4$ is a degree $2g$ multisection of $(\overline{W}_{-\infty,1},\overline{J}_{-\infty,1})$ satisfying $n^*(\overline{u})=0$.

We will use the convention to list the ends of a multisection $\overline{u}$ of $\overline{W}_{-\infty,1}$ in counterclockwise order, starting with the top end.

Definition 3.2.9. Let $\overline{J}_{-\infty,2}\in\overline{\mathcal{J}}_{-\infty,2}$ and $\mathbf{y},\mathbf{y}'\in\mathcal{S}_{\mathbf{b},\mathbf{a}}$. A *degree $k\le 2g$ multisection $\overline{u}$ of $(\overline{W}_{-\infty,2},\overline{J}_{-\infty,2})$ with ends $\mathbf{y}$ and $\mathbf{y}'$* is a holomorphic map

$$\overline{u}:(\dot{F},j)\to(\overline{W}_{-\infty,2},\overline{J}_{-\infty,2})$$

which is degree k multisection of $\pi_{B_{-\infty,2}}:\overline{W}_{-\infty,2}\to B_{-\infty,2}$ and which additionally satisfies the following:

(1) $\overline{u}(\partial\dot{F})\subset L^{-\infty,2}_{\overline{\mathbf{a}},+}\cup L^{-\infty,2}_{\overline{\mathbf{b}},-}$ and $\overline{u}$ maps each component of $\partial\dot{F}$ to a different $L^{-\infty,2}_{\overline{a}_i,+}$ or a different $L^{-\infty,2}_{\overline{b}_i,-}$;
(2) $\overline{u}$ converges to a cylinder over $[-2,2]\times\mathbf{y}$ as $t\to+\infty$ and to a cylinder over $[-2,2]\times\mathbf{y}'$ as $t\to-\infty$.

A $(\overline{W}_{-\infty,2},\overline{J}_{-\infty,2})$*-curve with ends $\mathbf{y}$ and $\mathbf{y}'$* is a degree $2g$ multisection of $(\overline{W}_{-\infty,2},\overline{J}_{-\infty,2})$ satisfying $n^*(\overline{u})=m$.

Next we discuss holomorphic curves with ends at z_∞. For $s = 1, \dots, 4$, the data $\overrightarrow{\mathcal{D}}_s$ at $z_\infty^{p_s}$ corresponding to the s^{th} end is given by

$$\overrightarrow{\mathcal{D}}_s = \{(i'_{s,k}, j'_{s,k}) \to (i_{s,k}, j_{s,k})\}_{k=1}^{p_s}.$$

When $s = 1$, $\mathcal{D}_{s=1}^{from} = \{(i'_{1,k}, j'_{1,k})\}_{k=1}^{p_1}$ and $\mathcal{D}_{s=1}^{to} = \{(i_{1,k}, j_{1,k})\}_{k=1}^{p_1}$ specify the initial points on $\overline{h}(\overline{a}_{i'_{1,k}, j'_{1,k}})$ and terminal points on $\overline{a}_{i_{1,k}, j_{1,k}}$, respectively; the cases $s = 2, 3, 4$ are analogous.

We then extend the definition of a degree k multisection $\overline{u}$ of $\overline{W}_{-\infty,1}$ to include those with ends $\mathbf{z}_s = \{z_\infty^{p_s}(\overrightarrow{\mathcal{D}}_s)\} \cup \mathbf{y}_s$, $s = 1, \dots, 4$, by attaching data $\mathcal{C}$ to $\overline{u}'$ and modifying Definition 3.2.8 in the same way Definition I.5.7.2 modifies Definition I.4.3.1. Degree k multisections of $\overline{W}_{-\infty,2}$ with ends $\mathbf{z} = \{z_\infty^p(\overrightarrow{\mathcal{D}})\} \cup \mathbf{y}$ and $\mathbf{z}' = \{z_\infty^q(\overrightarrow{\mathcal{D}}')\} \cup \mathbf{y}'$ are defined similarly.

Let $\mathcal{M}_{\overline{J}_{-\infty,1}}(\mathbf{z}_1, \mathbf{z}_2, \mathbf{z}_3, \mathbf{z}_4)$ be the moduli space of degree k multisections of $(\overline{W}_{-\infty,1}, \overline{J}_{-\infty,1})$ with ends $\mathbf{z}_s$, $s = 1, \dots, 4$, and let $\mathcal{M}_{\overline{J}_{-\infty,2}}(\mathbf{z}, \mathbf{z}')$ be the moduli space of degree k multisections of $(\overline{W}_{-\infty,2}, \overline{J}_{-\infty,2})$ with ends $\mathbf{z}$ and $\mathbf{z}'$. Also let

$$\mathcal{M}_{\overline{J}_{-\infty,2}}(\mathbf{z}, \mathbf{z}'; \overline{\mathfrak{m}}(-\infty)) \subset \mathcal{M}_{\overline{J}_{-\infty,2}}(\mathbf{z}, \mathbf{z}')$$

be the subset of curves which pass through $\overline{\mathfrak{m}}(-\infty)$. We then define the extended moduli spaces $\mathcal{M}^{\dagger,ext}_{\overline{J}_{-\infty,1}}(\mathbf{z}_1, \mathbf{z}_2, \mathbf{z}_3, \mathbf{z}_4)$ and $\mathcal{M}^{\dagger,ext}_{\overline{J}_{-\infty,2}}(\mathbf{z}, \mathbf{z}')$ in a manner similar to that of Section I.5.7.11. The precise definitions will be omitted.

3.2.6. *Indices.* We now discuss the Fredholm index $\operatorname{ind}(\overline{u})$ and the ECH index $I(\overline{u})$ of a $\overline{W}_\tau$-curve $\overline{u} : \dot{F} \to \overline{W}_\tau$ from $\mathbf{y}$ to $\mathbf{y}'$. The discussion will be brief since all the key ingredients have already been discussed in Sections I.5.5 and I.5.6.

We remark that, once again, $\operatorname{ind}(\overline{u})$ and $I(\overline{u})$ do not take into account the point constraint $\overline{\mathfrak{m}}(\tau)$ and that the condition "passing through $\overline{\mathfrak{m}}(\tau)$" is a codimension 2 condition.

Let $\check{\overline{W}}_\tau = \overline{W}_\tau - \{s > l(\tau) + 1\} - \{s < -l(\tau) - 1\}$, where $l(\tau)$ is given in Equation (3.1.1).[4] Let $\check{\overline{u}} : \check{F} \to \check{\overline{W}}_\tau$ be the compactification of $\overline{u}$, where $\check{F}$ is obtained by performing a real blow-up of F at its boundary punctures. We also define

$$\begin{aligned} Z_{\mathbf{y},\mathbf{y}'} = {} & (\{l(\tau) + 1\} \times [0,1] \times \mathbf{y}) \cup (\{-l(\tau) - 1\} \times [0,1] \times \mathbf{y}') \\ & \cup ((L^{\tau,+}_{\widehat{\mathbf{a}}} \cup L^{\tau,-}_{\widehat{\mathbf{b}}}) \cap \check{\overline{W}}_\tau). \end{aligned}$$

The trivialization $\tau^\star$ of $T\overline{S}$ along $Z_{\mathbf{y},\mathbf{y}'}$ is defined in a manner similar to that of Section I.4.4.2:[5] First we define the trivialization $\tau^\star$ of $T\overline{S}|_{\pi_{B_\tau}^{-1}(l(\tau)+1,1)}$ (resp. $T\overline{S}|_{\pi_{B_\tau}^{-1}(-l(\tau)-1,1)}$) along the $\widehat{a}_i$ (resp. $\widehat{b}_i$) by choosing a nonsingular tangent vector field along $\widehat{a}_i$ (resp. $\widehat{b}_i$). We then parallel transport $\tau^\star$ along $\partial\overline{W}_\tau$ and extend $\tau^\star$ arbitrarily to $\{l(\tau) + 1\} \times [0,1] \times \mathbf{y}$ and $\{-l(\tau) - 1\} \times [0,1] \times \mathbf{y}'$.

---

[4] Note that the definition of $\check{\overline{W}}_\tau$ is slightly different from that of Section 3.1.1.

[5] Here we are write $\tau^\star$ instead of τ due to the notational conflict with $\tau \in \mathbb{R}$.

Let $Q_{\tau^\star}(\breve{\overline{u}})$ be the relative intersection form given by intersecting $\breve{\overline{u}}$ and a pushoff of $\breve{\overline{u}}$ in the direction of $J\tau^\star$ along $\partial\breve{\overline{W}}_\tau$. Then

$$I(\overline{u}) = c_1(\breve{\overline{u}}^* T\breve{\overline{W}}_\tau, (\tau^\star, \partial_t)) + Q_{\tau^\star}(\breve{\overline{u}}) + \mu_{\tau^\star}(\mathbf{y}) - \mu_{\tau^\star}(\mathbf{y}') - 2g, \tag{3.2.2}$$

$$\operatorname{ind}(\overline{u}) = -\chi(\dot{F}) + 2c_1(\breve{\overline{u}}^* T\breve{\overline{W}}_\tau, (\tau^\star, \partial_t)) + \mu_{\tau^\star}(\mathbf{y}) - \mu_{\tau^\star}(\mathbf{y}') - 2g, \tag{3.2.3}$$

and the index inequality holds as usual:

$$\operatorname{ind}(\overline{u}) + 2\delta(\overline{u}) \leq I(\overline{u}), \tag{3.2.4}$$

where $\delta(\overline{u}) \geq 0$ and equals zero if and only if $\overline{u}$ is an embedding.

One can easily verify the following:

$$I(\sigma^+_\infty) = \operatorname{ind}(\sigma^+_\infty) = -1; \tag{3.2.5}$$

$$I(\sigma^-_\infty) = \operatorname{ind}(\sigma^-_\infty) = 0; \tag{3.2.6}$$

$$I(\sigma^\tau_\infty) = \operatorname{ind}(\sigma^\tau_\infty) = -1. \tag{3.2.7}$$

The Fredholm and ECH indices for $\overline{W}_{-\infty,i}$-curves can be defined and computed similarly. We now prove the following:

Lemma 3.2.10. *If there exists a $\overline{W}_{-\infty,2}$-curve $\overline{u}$ with $I = 2$ and ends $\mathbf{y}$ and $\mathbf{y}'$, then $\mathbf{y} = \{x^{\#}_{ij(i)}\}_{i=1}^{2g}$ and $\mathbf{y}' = \{x^{\#}_{ik(i)}\}_{i=1}^{2g}$, where $j(i)$ is odd and $k(i)$ is even for all i.*

In other words, $\mathbf{y}$ is a summand of the top generator $\Theta_{\overline{\mathbf{a}},\overline{\mathbf{b}}} \in \widehat{HF}(\overline{\mathbf{a}}, \overline{\mathbf{b}})$ and $\mathbf{y}'$ is a summand of the top generator $\Theta_{\overline{\mathbf{b}},\overline{\mathbf{a}}} \in \widehat{HF}(\overline{\mathbf{b}}, \overline{\mathbf{a}})$.

Proof. The proof is similar to the index calculation of Lemma 2.2.2. Let $\overline{u}$ be a $\overline{W}_{-\infty,2}$-curve. First consider the situation where $\overline{u}$ is in the homology class consisting of a copy $\{pt\} \times \overline{S}$ of the fiber and $2g$ trivial strips. Then

$$\operatorname{ind}(\{pt\} \times \overline{S}) = -\chi(\overline{S}) + 2c_1(T\overline{S}) = -(2-2g) + 2(2-2g) = 2-2g.$$

The strips contribute 0 to ind and there are $2g$ intersection points. Hence

$$I(\overline{u}) = (2-2g) + 0 + 2(2g) = 2g+2$$

when $\mathbf{y}' = \mathbf{y}$. The only way to lower $I(\overline{u})$ to 2 is to take $\mathbf{y} = \{x^{\#}_{ij(i)}\}_{i=1}^{2g}$ and $\mathbf{y}' = \{x^{\#}_{ik(i)}\}_{i=1}^{2g}$ so that all the $j(i)$ are odd and all the $k(i)$ are even. □

3.2.7. *Regularity.* We now discuss the regularity of the family $\{\overline{J}_\tau\}_{\tau\in\mathbb{R}}$.

Definition 3.2.11.

1. $\overline{J}_{-\infty,2} \in \overline{\mathcal{J}}_{-\infty,2}$ is *regular* if all the moduli spaces $\mathcal{M}^{\dagger,ext}_{\overline{J}_{-\infty,2}}(\mathbf{z}, \mathbf{z}')$ are transversely cut out.
2. $\overline{J}_{-\infty,1} \in \overline{\mathcal{J}}_{-\infty,1}$ is *regular* if all the moduli spaces $\mathcal{M}^{\dagger,ext}_{\overline{J}_{-\infty,1}}(\mathbf{z}_1, \dots, \mathbf{z}_4)$ are transversely cut out and the restrictions $\overline{J}_{-\infty,2}$ and $\overline{J}$ of $\overline{J}_{-\infty,1}$ to the ends are regular.

(3) $\overline{J}_{-\infty} = \overline{J}_{-\infty,1} \cup \overline{J}_{-\infty,2} \in \overline{\mathcal{J}}_{-\infty}$ is *regular* if $\overline{J}_{-\infty,i}$ are regular for $i = 1, 2$.

Definition 3.2.12. The family $\{\overline{J}_\tau\}_{\tau\in\mathbb{R}} \in \overline{\mathcal{I}}$ is *regular* if:

1. all the moduli spaces $\mathcal{M}^{\dagger,ext}_{\{\overline{J}_\tau\}}(\mathbf{z}, \mathbf{z}')$ are transversely cut out;
2. the restriction $\overline{J}$ of $\overline{J}_\tau$ to the positive and negative ends is regular;
3. $\overline{J}_+$ and $\overline{J}_-$ in the limit $\tau \to +\infty$ are regular; and
4. $\overline{J}_{-\infty}$ in the limit $\tau \to -\infty$ is regular.

Let $\overline{\mathcal{I}}^{reg}$ be the space of regular $\{\overline{J}_\tau\} \in \overline{\mathcal{I}}$. As usual, we have:

Lemma 3.2.13. *The generic $\{\overline{J}_\tau\} \in \overline{\mathcal{I}}$ is regular.*

Next we discuss the regularity of moduli spaces passing through $\overline{\mathfrak{m}}$. Since the point constraints $\overline{\mathfrak{m}}(\tau)$ are nongeneric, we need to introduce a perturbation of the family $\{\overline{J}_\tau\}$:

Definition 3.2.14. Let $\varepsilon > 0$ and let $\{U_\tau\}$, $U_\tau \subset \overline{W}_\tau$, be a family of open sets such that $U_\tau \not\ni \overline{\mathfrak{m}}(\tau)$. Then a family $\{\overline{J}^\Diamond_\tau\}$ of almost complex structures on $\{\overline{W}_\tau\}$ is $(\varepsilon, \{U_\tau\})$-*close* to a regular $\{\overline{J}_\tau\}$ if:

- $\overline{J}^\Diamond_\tau = \overline{J}_\tau$ on $\overline{W}_\tau - U_\tau$; and
- $\overline{J}^\Diamond_\tau$ is ε-close to $\overline{J}_\tau$ on U_τ.

Here the ε-closeness is measured with respect to a family $\{g_\tau\}_{\tau\in\mathbb{R}}$ of Riemannian metrics which is defined as follows: Let h be an s-invariant metric on $\mathbb{R} \times \overline{N}_2$ such that h, viewed as a metric on $\mathbb{R}\times[0,2]\times\overline{S}$, is also t-invariant on $\mathbb{R}\times[1,2]\times\overline{S}$. There is an extension of h to h_τ on $\mathbb{R} \times [0, r(\tau)] \times \overline{S}$ which is s- and t-invariant on $\mathbb{R} \times [2, r(\tau)] \times \overline{S}$. We then view h_τ as a metric on $\mathbb{R} \times \overline{N}_{r(\tau)}$ and define g_τ as the restriction of h_τ to $\overline{W}_\tau \subset \mathbb{R} \times \overline{N}_{r(\tau)}$.

We will use the following open sets U_τ: Let $\mathfrak{p}(\tau) \subset int(B_\tau)$ be a family, where:

- for each $\tau \in [-\infty, \infty]$, $\mathfrak{p}(\tau)$ is a finite number of points and $\#\mathfrak{p}(\tau)$ is independent of τ;
- $\mathfrak{p}(+\infty) = \lim_{\tau\to+\infty} \mathfrak{p}(\tau)$ nontrivially intersects $int(B_+)$ and $int(B_-)$;
- $\mathfrak{p}(-\infty) = \lim_{\tau\to-\infty} \mathfrak{p}(\tau)$ nontrivially intersects $int(B_{-\infty,1})$ and $int(B_{-\infty,2})$;
- for each $\tau \in [-\infty, \infty]$, $\overline{\mathfrak{m}}^b(\tau) \not\in \mathfrak{p}(\tau)$.

Then, for $\tau \in \mathbb{R}$, U_τ is an open δ-neighborhood of $K_\tau = \pi^{-1}_{B_\tau}(\mathfrak{p}(\tau)) - \{\rho < 2\delta\}$, where $\delta > 0$ is arbitrarily small. The sets $U_{\pm\infty}$ and $K_{\pm\infty}$ are defined as the limits of U_τ and K_τ as $\tau \to \pm\infty$. When we want to emphasize (ε, U_τ) or $(\varepsilon, \delta, \mathfrak{p})$, we write $\overline{J}^\Diamond_\tau(\varepsilon, U_\tau)$ or $\overline{J}^\Diamond_\tau(\varepsilon, \delta, \mathfrak{p}(\tau))$ for $\overline{J}^\Diamond_\tau$, $U_{\varepsilon,\delta,\mathfrak{p}(\tau)}$ for U_τ, and $K_{\mathfrak{p}(\tau),\delta}$ for K_τ.

We define a degree k *almost multisection* $\overline{u}$ *of* $(\overline{W}_\tau, \overline{J}^\Diamond_\tau)$ in the same way as a degree k multisection of $(\overline{W}_\tau, \overline{J}_\tau)$, except that $\overline{u}$ is a multisection of

$$\pi_{B\tau} : \overline{W}_\tau - \pi^{-1}_{B_-}(V_\tau) \to B_\tau - V_\tau,$$

where $V_\tau = \pi_{B_-}(U_\tau)$. The moduli spaces of almost multisections are defined in the same way as the moduli spaces of multisections, with $\overline{J}^\Diamond_\tau$ replacing $\overline{J}_\tau$. If

$\{K_\tau \not\ni \overline{\mathfrak{m}}(\tau)\}$ is a family of compact sets of $\overline{W}_\tau$, then the modifier $\{K_\tau\}$ means that $\overline{u}$ in $\overline{W}_\tau$ passes through K_τ.

Almost complex structures $\overline{J}^\Diamond_{-\infty,i}$, almost multisections on $(\overline{W}_{-\infty,i},\overline{J}^\Diamond_{-\infty,i})$, and moduli spaces of almost multisections are defined similarly.

Definition 3.2.15. The family $\{\overline{J}^\Diamond_\tau\}$ is *$\{K_\tau\}$-regular with respect to* $\overline{\mathfrak{m}}$ if all the moduli spaces $\mathcal{M}^{\dagger,ext,\{K_\tau\}}_{\{\overline{J}^\Diamond_\tau\}}(\mathbf{z},\mathbf{z}';\overline{\mathfrak{m}})$ are transversely cut out.

Lemma 3.2.16. *A generic family $\{\overline{J}^\Diamond_\tau\}$ is $\{K_\tau\}$-regular with respect to $\overline{\mathfrak{m}}$.*

Proof. The proof is similar to that of [MS, Theorem 3.1.7], with modifications as in Proposition I.5.8.8. ☐

The following can also be proved using a standard regularity argument:

Lemma 3.2.17. *If $\{\overline{J}_\tau\}$ is a generic family, then for $\varepsilon,\delta>0$ sufficiently small, there exist a generic family $\{\overline{J}^\Diamond_\tau(\varepsilon,\delta,\mathfrak{p}(\tau))\}$ which is $\{K_{\mathfrak{p}(\tau),\delta}\}$-regular with respect to $\overline{\mathfrak{m}}$ and disjoint finite subsets $\mathcal{T}_1,\mathcal{T}_2\subset\mathbb{R}$ with the following properties:*

1. *$\tau\in\mathcal{T}_1$ if and only if there exists $\overline{v}_\tau\in\mathcal{M}^{\dagger,s,irr,\mathrm{ind}=-1}_{\overline{J}_\tau}(\mathbf{z},\mathbf{z}')$ for some $\mathbf{z}$ and $\mathbf{z}'$.*
2. *$\tau\in\mathcal{T}_2$ if and only if there exists $\overline{v}_\tau\in\mathcal{M}^{\dagger,s,irr,\{K_\tau\},\mathrm{ind}=1}_{\overline{J}^\Diamond_\tau(\varepsilon,\delta,\mathfrak{p}(\tau))}(\mathbf{z},\mathbf{z}';\overline{\mathfrak{m}})$ for some $\mathbf{z}$ and $\mathbf{z}'$.*

Moreover, for each $\tau\in\mathcal{T}_i$ there is a unique such irreducible curve $\overline{v}_\tau$.

By shrinking $\overline{\mathcal{I}}$, we assume that all $\{\overline{J}_\tau\}\in\overline{\mathcal{I}}$ satisfy Lemma 3.2.17.

3.3. **Proof of half of Theorem 1.0.1.**

In this subsection we prove that $\Psi\circ\Phi$ is an isomorphism on the level of homology.

Recall the chain complexes $(\widehat{CF'}(S,\mathbf{a},h(\mathbf{a})),\partial')$ and $(\widetilde{CF}(S,\mathbf{a},h(\mathbf{a})),\widetilde{\partial})$ with isomorphic homology groups from Sections I.4.9.3 and I.6.6, and the maps $\widetilde{\Phi}$ and Ψ,Ψ' from Sections I.6.6 and I.7.1. Let q be the quotient map

$$q:\widehat{CF'}(S,\mathbf{a},h(\mathbf{a}))\to\widehat{CF}(S,\mathbf{a},h(\mathbf{a})).$$

The following is proved in this subsection:

Theorem 3.3.1. *Suppose $m\gg 0$. Then there exist maps*

$$H',\Theta_0':\widetilde{CF}(S,\mathbf{a},h(\mathbf{a}))\to\widehat{CF'}(S,\mathbf{b},h(\mathbf{b})),$$
$$\widetilde{V}:\widetilde{CF}(S,\mathbf{a},h(\mathbf{a}))\to\widetilde{CF}(S,\mathbf{b},h(\mathbf{b})),$$

which satisfy the following:

$$\Psi'\circ\widetilde{\Phi}-\Theta_0'=(\partial'H'+H'\widetilde{\partial})+\widetilde{\partial}_1\circ\widetilde{V}, \tag{3.3.1}$$

where $\widetilde{\partial}_1$ is as defined in Equation (I.7.1.1). Composing with q, we obtain the chain homotopy

$$\Psi\circ\widetilde{\Phi}-\Theta_0=\partial H+H\widetilde{\partial}, \tag{3.3.2}$$

where $\Theta_0=q\circ\Theta_0'$ induces an isomorphism on homology.

Since Θ_0 is an isomorphism on the level of homology, so is $\Psi \circ \widetilde{\Phi}$. In view of Corollary I.6.6.7, $\Psi \circ \Phi$ is also an isomorphism on the level of homology.

Proof. We prove Theorem 3.3.1, assuming the results of Sections 3.4–3.9. In Steps 1–3 we consider the situation where the holomorphic curves $\overline{u}$ in $\overline{W}_\tau$ are asymptotic to some $\mathbf{y} \in \mathcal{S}_{\mathbf{a},h(\mathbf{a})}$ at the positive end. In Step 4 we describe the modifications needed for the situation where $\overline{u}$ is asymptotic to some $\mathbf{z} = \{z_{\infty,i}\}_{i\in\mathcal{I}} \cup \mathbf{y}$ at the positive end and $\mathcal{I}$ is a subset of $\{1,\dots,2g\}$ with $|\mathcal{I}| > 0$. Steps 1–4 prove Equation (3.3.2). In Step 5 we prove that Θ_0 induces an isomorphism on homology by further degenerating $\overline{W}_{-\infty,1}$.

Suppose $m \gg 0$. Fix $\mathfrak{p}(\tau)$ and choose $\{\overline{J}_\tau\} \in \overline{\mathcal{I}}^{reg}$ satisfying Lemma 3.2.17. For sufficiently small $\varepsilon, \delta > 0$ (which depend on the choices of m and $\{\overline{J}_\tau\}$), there exists $\{\overline{J}_\tau^\Diamond(\varepsilon,\delta,\mathfrak{p}(\tau))\}$ so that Lemma 3.2.17 holds.

Fix $\mathbf{y} \in \mathcal{S}_{\mathbf{a},h(\mathbf{a})}$, $\mathbf{y}' \in \mathcal{S}_{\mathbf{b},h(\mathbf{b})}$ and abbreviate

$$\mathcal{M} = \mathcal{M}^{I=2,n^*=m}_{\{\overline{J}_\tau^\Diamond(\varepsilon,\delta,\mathfrak{p}(\tau))\}}(\mathbf{y},\mathbf{y}';\overline{\mathfrak{m}}),\ \mathcal{M}^{\{K_{\mathfrak{p}(\tau),\delta}\}} = \mathcal{M}^{I=2,n^*=m,\{K_{\mathfrak{p}(\tau),\delta}\}}_{\{\overline{J}_\tau^\Diamond(\varepsilon,\delta,\mathfrak{p}(\tau))\}}(\mathbf{y},\mathbf{y}';\overline{\mathfrak{m}}).$$

Let $\overline{\mathcal{M}}$ be the SFT compactification of $\mathcal{M}$ and let $\partial\mathcal{M} = \overline{\mathcal{M}} - \mathcal{M}$ be the boundary of $\mathcal{M}$. If $U \subset [-\infty,+\infty]$, then we write $\partial_U\mathcal{M}$ for the set of $\overline{u}_\infty \in \partial\mathcal{M}$ where $\overline{u}_\infty$ is a building which corresponds to some $\tau \in U$. By Lemma 3.2.16, we may take $\mathcal{M}^{\{K_{\mathfrak{p}(\tau),\delta}\}}$ to be regular.

Step 1 (Breaking at $+\infty$). Recall the definition of a *bad radial ray* $\mathcal{R}_\phi$ from Definition I.7.8.7. We now enlarge the class of bad radial rays as follows: Let

$$\coprod_{\mathbf{y},\delta_0^{l_i}\gamma'} \mathcal{M}^{I=0}_{(\overline{J}_+)_\infty}(\mathbf{y},\delta_0^{l_i}\gamma') = \{C_1,\dots,C_r\},^{6}$$

where $(\overline{J}_+)_\infty$ is the limit of $(\overline{J}_+)_m$ as $m \to \infty$ and the disjoint union is over all $\mathbf{y}$ and $\delta_0^{l_i}\gamma'$ with $l_i > 0$, and let $f_{ij} : \mathbb{R}/2l_{ij}\mathbb{Z} \to \mathbb{C}$ be the asymptotic eigenfunctions corresponding to the ends $\delta_0^{l_{ij}}$ of C_i, where $(l_{i1},\dots,l_{i\lambda_i})$ is the partition of l_i corresponding to C_i. We then add the radial rays which pass through

$$\{f_{ij}(t) \mid i = 1,\dots,r; j = 1,\dots,\lambda_i; 0 < t < 2l_{ij}; t \equiv 3/2 \bmod 2\}$$

to the class of bad radial rays. We can still assume that $\mathcal{R}_\pi$ is a good radial ray.

We then have the following:

---

[6] The notation is the same as that of Section I.7.8.2, except the C_i belong to different moduli spaces.

Lemma 3.3.2. *$\partial_{\{+\infty\}}\mathcal{M}\subset A_1\sqcup A_2$, where*

$$A_1=\coprod_{\gamma\in\widehat{\mathcal{O}}_{2g}}\left(\mathcal{M}^{I=0}_{J^\Diamond_+(\varepsilon,\delta,\mathfrak{p}(+\infty))}(\mathbf{y},\gamma)\times\mathcal{M}^{I=2,n^*=m}_{\overline{J}^\Diamond_-(\varepsilon,\delta,\mathfrak{p}(+\infty))}(\gamma,\mathbf{y}';\overline{\mathfrak{m}}(+\infty))\right);$$

$$A_2=\coprod_{\delta_0\gamma,\{z_\infty\}\cup\mathbf{y}''}\left(\mathcal{M}^{I=1,n^*=m-1,f_{\delta_0}}_{\overline{J}^\Diamond_+(\varepsilon,\delta,\mathfrak{p}(+\infty))}(\mathbf{y},\delta_0\gamma)\times\mathcal{M}^{I=0,n^*=0}_{\overline{J}^\Diamond_-(\varepsilon,\delta,\mathfrak{p}(+\infty))}(\delta_0\gamma,\{z_\infty\}\cup\mathbf{y}'')\right.$$
$$\left.\times\,\mathcal{M}^{I=1,n^*=1}_{\overline{J}}(\{z_\infty\}\cup\mathbf{y}'',\mathbf{y}')\right),$$

if $\mathbf{y}'=\{x_i^j\}\times\mathbf{y}''$ for some x_i^j and $A_2=\varnothing$ otherwise. The disjoint union for A_2 ranges over all $\delta_0\gamma$ such that $\gamma\in\widehat{\mathcal{O}}_{2g-1}$ and all $\{z_\infty\}\cup\mathbf{y}''$ such that $\mathbf{y}'$ can be written as $\{x_i^j\}\cup\mathbf{y}''$ for some x_i^j. Here we have omitted the potential contributions of connector components and we are writing $x_i^0:=x_i$ and $x_i^1:=x_i'$.

Here f_{δ_0} is a nonzero normalized asymptotic eigenfunction of δ_0 at the negative end such that $f_{\delta_0}(\frac{3}{2})$ lies on the good radial ray $\mathcal{R}_\pi$. Used as a modifier, f_{δ_0} stands for "the normalized asymptotic eigenfunction at the negative end δ_0 is f_{δ_0}".

Remark 3.3.3. If $\overline{u}=\overline{u}'\cup\overline{u}''\in\mathcal{M}^{I=0,n^*=0}_{\overline{J}^\Diamond_-(\varepsilon,\delta,\mathfrak{p}(+\infty))}(\delta_0\gamma,\{z_\infty\}\cup\mathbf{y}'')$, then $\overline{u}$ consists of $\overline{u}'=\sigma^-_\infty$ and a curve $\overline{u}''$ from γ to $\mathbf{y}''$ which is arbitrarily close to a curve with image in W_-. If $\overline{u}\in\mathcal{M}^{I=1,n^*=1}_{\overline{J}}(\{z_\infty\}\cup\mathbf{y}'',\{x_i^j\}\cup\mathbf{y}'')$, then $\overline{u}$ consists of one thin strip from z_∞ to x_i^j and $2g-1$ trivial strips.

Lemma 3.3.2 will be proved in Section 3.4. Gluing the pairs in A_1 using the Hutchings-Taubes gluing theorem [HT1, HT2] (see Section I.6.5) accounts for the term $\Psi'\circ\widetilde{\Phi}$ in Equation (3.3.1).

Gluing the triples in A_2 accounts for the term $\partial_1\circ\widetilde{V}$. This is similar to Section I.7.13, and the details will be omitted. For each triple $(\overline{v}_+,\overline{v}_-,\overline{v}_{-1,1})\in A_2$, where the nontrivial component of $\overline{v}_{-1,1}$ is a thin strip from $z_{\infty,i,0}$ to x_i, there is another triple $(\overline{v}_+,\overline{v}_-,\overline{v}^*_{-1,1})\in A_2$, where the nontrivial component of $\overline{v}^*_{-1,1}$ is a thin strip from $z_{\infty,i,1}$ to x_i'.

Since $\mathcal{M}^{\{K_{\mathfrak{p}(\tau)},\delta\}}$ is regular but $\mathcal{M}-\mathcal{M}^{\{K_{\mathfrak{p}(\tau)},\delta\}}$ is not a priori regular, it remains to verify the following:

Claim 3.3.4. *For $\varepsilon,\delta>0$ sufficiently small, the restriction of $\partial\mathcal{M}^{\{K_{\mathfrak{p}(\tau)},\delta\}}$ to a neighborhood of $\tau=+\infty$ is $A_1\cup\widetilde{A}_2$, where $\#A_2\equiv\#\widetilde{A}_2$ mod 2.*

Proof. If $\varepsilon,\delta>0$ are sufficiently small, then $\overline{v}_-$ passes through $K_{\mathfrak{p}(\tau),\delta}$ whenever $(\overline{v}_+,\overline{v}_-)\in A_1$. Hence if $\overline{u}\in\mathcal{M}$ is close to a curve in A_1, then $\overline{u}\in\mathcal{M}^{\{K_{\mathfrak{p}(\tau)},\delta\}}$. This accounts for the term A_1 in the claim.

Next suppose that $\overline{u}\in\mathcal{M}-\mathcal{M}^{\{K_{\mathfrak{p}(\tau)},\delta\}}$ for τ near $+\infty$. Then $\overline{u}$ is arbitrarily close to a building of type A_2. By Lemma I.7.2.3 and Theorem I.7.2.2, $\mathcal{M}^{\{K_{\mathfrak{p}(\tau)},\delta\}}$ is obtained from $\mathcal{M}$ by truncating ends that are close to A_2; hence $\widetilde{A}_2:=\partial\mathcal{M}^{\{K_{\mathfrak{p}(\tau)},\delta\}}-A_1$ satisfies $\#A_2\equiv\#\widetilde{A}_2$ mod 2. □

Step 2 (Breaking at $-\infty$).

Lemma 3.3.5. *$\partial_{\{-\infty\}}\mathcal{M} \subset A_3$, where*

$$A_3 = \coprod_{\mathbf{y}_2,\mathbf{y}_4} \left(\mathcal{M}^{I=0,n^*=0}_{\overline{J}^{\Diamond}_{-\infty,1}(\varepsilon,\delta,\mathfrak{p}(-\infty))}(\mathbf{y},\mathbf{y}_2,\mathbf{y}',\overline{h}(\mathbf{y}_4)) \times \mathcal{M}^{I=2,n^*=m}_{\overline{J}^{\Diamond}_{-\infty,2}(\varepsilon,\delta,\mathfrak{p}(-\infty))}(\mathbf{y}_4,\mathbf{y}_2;\overline{\mathfrak{m}}(-\infty)) \right).$$

Here the union is over all $\mathbf{y}_2$, $\mathbf{y}_4$ such that $\mathbf{y}_4 = \{x^{\#}_{ij(i)}\}_{i=1}^{2g}$ and $\mathbf{y}_2 = \{x^{\#}_{ik(i)}\}_{i=1}^{2g}$, where $j(i)$ is odd and $k(i)$ is even.

Lemma 3.3.5 will be proved in Section 3.5. Gluing the pairs $(\overline{v}_1, \overline{v}_2)$ in A_3 accounts for the term Θ'_0 in Equation (3.3.1). The map Θ'_0 is given by:

$$\langle \Theta'_0(\mathbf{y}), \mathbf{y}' \rangle = \sum_{\mathbf{y}_2,\mathbf{y}_4} \# \mathcal{M}^{I=0,n^*=0}_{\overline{J}^{\Diamond}_{-\infty,1}(\varepsilon,\delta,\mathfrak{p}(-\infty))}(\mathbf{y},\mathbf{y}_2,\mathbf{y}',\overline{h}(\mathbf{y}_4)),$$

where $\mathbf{y}_2$, $\mathbf{y}_4$ are as in A_3, and Theorem 2.4.2 can be rephrased as (verification left to the reader):

Theorem 3.3.6. *Suppose $\mathbf{y}_4 = \{x^{\#}_{ij(i)}\}_{i=1}^{2g}$ and $\mathbf{y}_2 = \{x^{\#}_{ik(i)}\}_{i=1}^{2g}$, where $j(i)$ is odd and $k(i)$ is even. Then*

$$\# \mathcal{M}^{I=2,n^*=m}_{\overline{J}^{\Diamond}_{-\infty,2}(\varepsilon,\delta,\mathfrak{p}(-\infty))}(\mathbf{y}_4,\mathbf{y}_2;\overline{\mathfrak{m}}(-\infty)) \equiv 1 \textit{ mod } 2.$$

The argument of Claim 3.3.4 gives:

Claim 3.3.7. *For $\varepsilon, \delta > 0$ sufficiently small, the restriction of $\partial\mathcal{M}^{\{K_{\mathfrak{p}(\tau),\delta}\}}$ to a neighborhood of $-\infty$ is A_3.*

Step 3 (Breaking in the middle).

Lemma 3.3.8. *$\partial_{(-\infty,+\infty)}\mathcal{M} \subset A_4 \sqcup A_5$, where:*

$$A_4 = \coprod_{\mathbf{y}''\in\mathcal{S}_{\mathbf{a},h(\mathbf{a})}} \left(\mathcal{M}^{I=1}_{J}(\mathbf{y},\mathbf{y}'') \times \mathcal{M}^{I=1,n^*=m}_{\{\overline{J}^{\Diamond}_{\tau}(\varepsilon,\delta,\mathfrak{p}(\tau))\}}(\mathbf{y}'',\mathbf{y}';\overline{\mathfrak{m}}) \right);$$

$$A_5 = \coprod_{\mathbf{y}'''\in\mathcal{S}_{\mathbf{b},h(\mathbf{b})}} \left(\mathcal{M}^{I=1,n^*=m}_{\{\overline{J}^{\Diamond}_{\tau}(\varepsilon,\delta,\mathfrak{p}(\tau))\}}(\mathbf{y},\mathbf{y}''';\overline{\mathfrak{m}}) \times \mathcal{M}^{I=1}_{J}(\mathbf{y}''',\mathbf{y}') \right).$$

Lemma 3.3.8 will be proved in Section 3.6. Using the technique of [Li, Prop. A.1 and A.2], we can glue each of the pairs in A_4 and A_5. This gluing accounts for the term $\partial' H' + H'\widetilde{\partial}$ in Equation (3.3.1).

Claim 3.3.9. *For $\varepsilon, \delta > 0$ sufficiently small,*

$$\partial\mathcal{M}^{\{K_{\mathfrak{p}(\tau),\delta}\}} = A_1 \cup \widetilde{A}_2 \cup A_3 \cup A_4 \cup A_5.$$

Step 4 (Additional degenerations). In this step we give the necessary modifications for

$$\mathcal{M} = \mathcal{M}^{I=2,n^*=m+|\mathcal{I}|}_{\{\overline{J}^{\Diamond}_{\tau}(\varepsilon,\delta,\mathfrak{p}(\tau))\}}(\mathbf{z},\mathbf{y}';\overline{\mathfrak{m}}),$$

where $\mathbf{z} = \{z_{\infty,i}\}_{i\in\mathcal{I}} \cup \mathbf{y}$, $\mathbf{y}$ and $\mathbf{y}'$ are tuples in S, and $\mathcal{I}$ is a subset of $\{1,\ldots,2g\}$.

The following is proved in Section 3.7:

Lemma 3.3.10. *$\partial_{\{+\infty\}}\mathcal{M} \subset A_2'$, where*

$$A_2' = \coprod_{\delta_0\gamma, \{z_\infty\}\cup \mathbf{y}''} \Big(\mathcal{M}^{I=1, n^*=m+|\mathcal{I}|-1, f_{\delta_0}}_{\overline{J}^\Diamond_+(\varepsilon,\delta,\mathfrak{p}(+\infty))}(\mathbf{z}, \delta_0\gamma) \times \mathcal{M}^{I=0,n^*=0}_{\overline{J}^\Diamond_-(\varepsilon,\delta,\mathfrak{p}(+\infty))}(\delta_0\gamma, \{z_\infty\}\cup\mathbf{y}'') \\ \times \mathcal{M}^{I=1,n^*=1}_{\overline{J}}(\{z_\infty\}\cup\mathbf{y}'', \mathbf{y}')\Big),$$

if $\mathbf{y}' = \{x_i^j\} \times \mathbf{y}''$ for some x_i^j and $A_2' = \varnothing$ otherwise. The disjoint union for A_2' ranges over all $\delta_0\gamma$ such that $\gamma \in \widehat{\mathcal{O}}_{2g-1}$ and all $\{z_\infty\}\cup\mathbf{y}''$ such that $\mathbf{y}'$ can be written as $\{x_i^j\}\cup\mathbf{y}''$ for some x_i^j. Here we have omitted the potential contributions of connector components and we are writing $x_i^0 := x_i$ and $x_i^1 := x_i'$.

Since $\widetilde{\Phi}(\mathbf{z}) = 0$ if $|\mathcal{I}| \geq 1$ by Lemma I.6.6.5, the "gluing" accounts for the terms $\Psi \circ \widetilde{\Phi} = 0$ and $q \circ \widetilde{\partial}_1 \circ \widetilde{V}$.

Next, the following is proved in Section 3.8:

Lemma 3.3.11. *$\partial_{\{-\infty\}}\mathcal{M} \subset A_3'$, where:*

$$A_3' = \coprod_{\mathbf{y}_2,\mathbf{y}_4} \left(\mathcal{M}^{I=0,n^*=|\mathcal{I}|}_{\overline{J}^\Diamond_{-\infty,1}(\varepsilon,\delta,\mathfrak{p}(-\infty))}(\mathbf{z}, \mathbf{y}_2, \mathbf{y}', \overline{h}(\mathbf{y}_4)) \times \mathcal{M}^{I=2,n^*=m}_{\overline{J}^\Diamond_{-\infty,2}(\varepsilon,\delta,\mathfrak{p}(-\infty))}(\mathbf{y}_4, \mathbf{y}_2; \overline{\mathfrak{m}}(-\infty))\right)$$

and the summation is over $\mathbf{y}_2$ and $\mathbf{y}_4$ as in Lemma 3.3.5.

The following is proved in Section 3.9. The corresponding gluing accounts for the term $\partial H + H\widetilde{\partial}$ in Equation (3.3.2) when $|\mathcal{I}| \geq 1$.

Lemma 3.3.12. *$\partial_{(-\infty,+\infty)}\mathcal{M} \subset A_4' \sqcup A_5'$, where:*

$$A_4' = \coprod_{\mathbf{z}'} \left(\mathcal{M}^{I=1,n^*\leq|\mathcal{I}|}_{\overline{J}}(\mathbf{z}, \mathbf{z}') \times \mathcal{M}^{I=1,n^*\leq m+|\mathcal{I}|}_{\{\overline{J}^\Diamond_\tau(\varepsilon,\delta,\mathfrak{p}(\tau))\}}(\mathbf{z}', \mathbf{y}'; \overline{\mathfrak{m}})\right);$$

$$A_5' = \coprod_{\mathbf{y}'''\in\mathcal{S}_{\mathbf{b},h(\mathbf{b})}} \left(\mathcal{M}^{I=1,n^*=m+|\mathcal{I}|}_{\{\overline{J}^\Diamond_\tau(\varepsilon,\delta,\mathfrak{p}(\tau))\}}(\mathbf{z}, \mathbf{y}'''; \overline{\mathfrak{m}}) \times \mathcal{M}^{I=1}_{J}(\mathbf{y}''', \mathbf{y}')\right).$$

Moreover, if $\overline{v}_{1,1} \in \mathcal{M}^{I=1,n^*\leq|\mathcal{I}|}_{\overline{J}}(\mathbf{z}, \mathbf{z}')$, then either (i) $\overline{v}^\sharp_{1,1}$ is a thin strip and $\overline{v}^\flat_{1,1}$ is a union of trivial strips, or (ii) $\overline{v}^\sharp_{1,1} = \varnothing$ and $\overline{v}^\flat_{1,1}$ has image in W.

As before, the analog of Lemma 3.3.9 holds.

Step 5 (The map Θ_0). The map Θ_0 is defined as follows:

$$\Theta_0 : \widetilde{CF}(\mathbf{a}, h(\mathbf{a})) \to \widehat{CF}(\mathbf{b}, h(\mathbf{b})),$$

$$\langle \Theta_0(\mathbf{z}), \mathbf{y}'\rangle = \sum_{\mathbf{y}_2,\mathbf{y}_4} \#\mathcal{M}^{I=0,n^*=|\mathcal{I}|}_{\overline{J}_{-\infty,1}}(\mathbf{z}, \mathbf{y}_2, \mathbf{y}', \overline{h}(\mathbf{y}_4)).$$

Lemma 3.3.13. *The map Θ_0 induces an isomorphism on the level of homology.*


$\overline{\mathbf{a}}$
$\overline{h}(\overline{\mathbf{a}})$
$\overline{\mathbf{b}}$
$\overline{h}(\overline{\mathbf{b}})$

FIGURE 7. Degeneration of the base $B_{-\infty,1}$.

Proof of Lemma 3.3.13. We degenerate the base $B_{-\infty,1}$ as given in Figure 7.

The map Θ_0 is chain homotopic to $\Theta_B \circ \Theta_T$, where:

$$\Theta_T : \widetilde{CF}(\mathbf{a}, h(\mathbf{a})) \to \widehat{CF}(\mathbf{b}, h(\mathbf{a})),$$

$$\Theta_B : \widehat{CF}(\mathbf{b}, h(\mathbf{a})) \to \widehat{CF}(\mathbf{b}, h(\mathbf{b}))$$

are defined by tensoring with the top generators $\Theta_{\overline{\mathbf{b}},\overline{\mathbf{a}}} \in \widehat{CF}(\overline{\mathbf{b}}, \overline{\mathbf{a}})$ and $\Theta_{\overline{h}(\overline{\mathbf{a}}),\overline{h}(\overline{\mathbf{b}})} \in \widehat{CF}(\overline{h}(\overline{\mathbf{a}}), \overline{h}(\overline{\mathbf{b}}))$. During the chain homotopy there is one undesirable type of breaking: a two-level building $(\overline{u}_1, \overline{u}_2)$, where $\overline{u}_1$ is an index $I = -1$, $n^* = |\mathcal{I}|$ curve with ends $\mathbf{z}, \mathbf{y}_2', \mathbf{y}', \overline{h}(\mathbf{y_4})$; $\overline{u}_2 \in \mathcal{M}^{I=1,n^*=0}_{\overline{J}_{-\infty,2}}(\mathbf{y}_2, \mathbf{y}_2')$; and $\mathbf{y}_2 - \mathbf{y}_2' = \{x^\#_{i1}\}$ or $\{x^\#_{i3}\}$. One can verify that the only possible component of $\overline{u}_1$ with left end $x^\#_{i1}$ projects to the quadrilateral Q with edges $\overline{a}_i, \overline{b}_i, \overline{h}(\overline{b}_i), \overline{h}(\overline{a}_i)$ in Figure 6. However, the component corresponding to Q has ECH index $I = 0$, which is a contradiction. This implies that the undesirable breaking does not exist.

Both Θ_T and Θ_B — and hence Θ_0 — induce isomorphisms on the level of homology. $\square$

This completes the proof of Theorem 3.3.1, assuming the results from Sections 3.4–3.9. $\square$

3.4. **Degeneration at $+\infty$.** In this subsection we study the limit of holomorphic maps to $\overline{W}_\tau$ as $\tau \to \infty$, i.e., when $\overline{W}_\tau$ degenerates into the concatenation of $\overline{W}_+$ with $\overline{W}_-$ along the ECH-type end, in order to prove Lemma 3.3.2.

We assume that $m \gg 0$; $\varepsilon, \delta > 0$ are sufficiently small; and $\{\overline{J}_\tau\} \in \overline{\mathcal{I}}^{reg}$ and $\{\overline{J}^\Diamond_\tau(\varepsilon, \delta, \mathfrak{p}(\tau))\}$ satisfy Lemma 3.2.17. Fix $\mathbf{y} \in \mathcal{S}_{\mathbf{a},h(\mathbf{a})}$, $\mathbf{y}' \in \mathcal{S}_{\mathbf{b},h(\mathbf{b})}$ and let

$$\mathcal{M} = \mathcal{M}^{I=2,n^*=m}_{\{\overline{J}^\Diamond_\tau(\varepsilon,\delta,\mathfrak{p}(\tau))\}}(\mathbf{y}, \mathbf{y}'; \overline{\mathfrak{m}}), \quad \mathcal{M}_\tau = \mathcal{M}^{I=2,n^*=m}_{\overline{J}^\Diamond_\tau(\varepsilon,\delta,\mathfrak{p}(\tau))}(\mathbf{y}, \mathbf{y}'; \overline{\mathfrak{m}}(\tau)).$$

We will analyze $\partial_{\{+\infty\}}\mathcal{M}$.

Let $\overline{u}_i$, $i \in \mathbb{N}$, be a sequence of curves in $\mathcal{M}$ such that $\overline{u}_i \in \mathcal{M}_{\tau_i}$ and $\lim\limits_{i\to\infty} \tau_i = +\infty$, and let

$$\overline{u}_\infty = (\overline{v}_{-1,1} \cup \cdots \cup \overline{v}_{-1,c}) \cup \overline{v}_- \cup (\overline{v}_{0,1} \cup \cdots \cup \overline{v}_{0,b}) \cup \overline{v}_+ \cup (\overline{v}_{1,1} \cup \cdots \cup \overline{v}_{1,a})$$

be the limit holomorphic building in order from bottom to top, where $\overline{v}_{-1,j}$, $j = 1,\dots,c$, maps to $\overline{W}$; $\overline{v}_-$ maps to $\overline{W}_-$; $\overline{v}_{0,j}$, $j = 1,\dots,b$, maps to $\overline{W'}$; $\overline{v}_+$ maps to $\overline{W}_+$; and $\overline{v}_{1,j}$, $j = 1,\dots,a$, maps to $\overline{W}$. Here we are allowing the possibility that a, b, or $c = 0$. For notational convenience, sometimes we will refer to $\overline{v}_+$ as $\overline{v}_{0,b+1}$ or $\overline{v}_{1,0}$ and $\overline{v}_-$ as $\overline{v}_{-1,c+1}$ or $\overline{v}_{0,0}$.

Notation.

- We write $\dot{F}_*$, $\dot{F}'_*$, $\dot{F}''_*$ for the domains of $\overline{v}_*$, $\overline{v}'_*$, $\overline{v}''_*$.
- We write p_* for the covering degree of $\overline{v}'_*$.
- If $\widetilde{v}_{*_i}$, $i = 1, 2$, is a union of components of $\overline{v}_{*_i}$, then we write $\widetilde{v}_{*_1} \succ \widetilde{v}_{*_2}$ (resp. $\widetilde{v}_{*_1} \succeq \widetilde{v}_{*_2}$) to indicate that the level $\overline{v}_{*_1}$ is above (resp. equal to or above) the level $\overline{v}_{*_2}$.

Since ghost components can be eliminated by the discussion in Lemma I.6.1.7 they will not be explicitly mentioned in the rest of the paper.

We have the following two constraints:

$$n^*(\overline{u}_i) = \sum_{\overline{v}_*} n^*(\overline{v}_*) = m; \tag{3.4.1}$$

$$I(\overline{u}_i) = \sum_{\overline{v}_*} I(\overline{v}_*) = 2, \tag{3.4.2}$$

where the summations are over all the levels $\overline{v}_*$ of $\overline{u}_\infty$.

Outline of proof of Lemma 3.3.2. The proof of Lemma 3.3.2 follows the same general outline of Section I.7.5–I.7.12: First we calculate the contributions to n^* of the ends that limit to multiples of z_∞ or δ_0 in Section 3.4.1 and obtain lower bounds on the ECH indices of the levels $\overline{v}_*$ in Section 3.4.2, under the assumption that there are no boundary punctures that are removable at z_∞. Boundary punctures are treated in Sections 3.4.3 and 3.4.4. The main new difficulty is to show that $I(\overline{v}_*) \geq I(\overline{v}'_*) + I(\overline{v}''_*)$ for $\overline{v}_* \succeq \overline{v}_+$; this uses the more complicated version of the ECH index inequality given in Lemma I.5.7.18. We then use Equations (3.4.1) and (3.4.2) to obtain Lemma 3.4.17, which describes the case when $\overline{v}'_* \cup \overline{v}^\sharp_* = \varnothing$ for all levels $\overline{v}_*$, and Lemma 3.4.21, which gives a preliminary list when $\overline{v}'_* \cup \overline{v}^\sharp_* \neq \varnothing$ for some $\overline{v}_*$. The renormalization argument from Section I.7.9–7.11, given in Lemma 3.4.24, eliminates all the possibilities with the exception of Case (2_1) of Lemma 3.4.21 when $\overline{v}'_* \cup \overline{v}^\sharp_* \neq \varnothing$ for some $\overline{v}_*$.

3.4.1. *Intersection numbers.* In this subsection we give the analogs of Lemmas I.7.5.1–I.7.5.4 for $\overline{u}_\infty \in \partial_{\{+\infty\}}\mathcal{M}$:

Lemma 3.4.1.

1. *If $\overline{v}''_*$ has a negative end $\mathcal{E}_-$ that converges to δ_0^p, then $n^*(\mathcal{E}_-) \geq m - p$.*
2. *If $\overline{v}''_*$ has a positive end $\mathcal{E}_+$ that converges to δ_0^p, then $n^*(\mathcal{E}_+) \geq p$.*

Proof. This is analogous to Lemma I.7.5.1 and is proved in the same way. □

Definition 3.4.2. An end $\mathcal{E}$ is *nontrivial* if it is not an end of a trivial cylinder or trivial strip.

The following Definition 3.4.3 describes the breaking of the component $\overline{u}_i|_{\partial \dot{G}_i}$ which maps to $L^{\tau_i,+}_{\overline{a}_k}$, into a sequence $g^1_a, \dots, g^1_1, g_0, g^0_1, \dots, g^0_a$ as $\tau_i \to \infty$. Here $\dot{G}_i$ is the domain of $\overline{u}_i$. Definition 3.4.3 plays an important role in the proof of Lemma 3.4.4, which is a refined version of Lemma I.7.5.2, where the ends are considered collectively as well as individually. The proof strategy will usually be referred to as the *continuation argument*.

Let $\partial_+ B_\tau$ (resp. $\partial_+ \overline{W}_\tau$) be the $s > 0$ boundary of B_τ (resp. $\overline{W}_\tau$) and let $\mathcal{C}_{1,j}$, $j \geq 0$, be the data for $\overline{v}'_{1,j}$; see Section I.5.7.2 for the definition of the data $\mathcal{C}_{1,j}$.

Definition 3.4.3 (Continuations)**.** Fix some $k > 0$ and consider the sequence

$$g^1_a, \dots, g^1_1, g_0, g^0_1, \dots, g^0_a \tag{3.4.3}$$

which satisfies the following:

1. $g^*_j = \overline{v}_{1,j}(f^*_j)$ where $j \geq 0$, $* = 0, 1, \varnothing$, and f^*_j is a component of $\partial \dot{F}_{1,j}$.
2. If f^*_j is a component of $\partial \dot{F}''_{1,j}$, then $g^1_j \subset \mathbb{R} \times \{1\} \times \overline{a}_k$ and $g^0_j \subset \mathbb{R} \times \{0\} \times \overline{h}(\overline{a}_k)$ if $j > 0$ and $g_0 \subset L^+_{\overline{a}_k}$ if $j = 0$.
3. If f^*_j is a component of $\partial \dot{F}'_{1,j}$, then f^*_j comes with extra data $\mathcal{C}_{1,j}$ which assigns:
$$f^1_j \mapsto \mathbb{R} \times \{1\} \times \overline{a}_k, \quad f^0_j \mapsto \mathbb{R} \times \{0\} \times \overline{h}(\overline{a}_k), \quad f_0 \mapsto L^+_{\overline{a}_k}.$$

If g^*_j is any element of Sequence (3.4.3), then Sequence (3.4.3) is the *continuation of g^*_j along* $\partial_+ B_\tau$ and the terms to the right of g^*_j in Sequence (3.4.3) form the *continuation of g^*_j in the direction of* $\partial_+ B_\tau$. An element g^*_j is *trivial* if the corresponding f^*_j satisfies (3). For convenience we write $g_0 = g^1_0 = g^0_0$.

Let $\rho_0 > 0$ be small and let $\pi_{D_{\rho_0}}$ be the projection of

$$\{\rho \leq \rho_0\} \subset \overline{W} - int(W) \ \text{ or } \ \{\rho \leq \rho_0, |s| \geq l(\tau) + 1\} \subset \overline{W}_\tau - int(W_\tau)$$

to D_{ρ_0} along the stable Hamiltonian vector field $\overline{R}_\tau$.

Lemma 3.4.4. *Suppose $\overline{v}'_{1,j} \cup \overline{v}^\sharp_{1,j} \neq \varnothing$ for some $j > 0$. Let $\mathcal{E}_{-,i}$, $i = 1, \dots, q$, be the negative ends of $\cup_{j=1}^a \overline{v}^\sharp_{1,j}$ that converge to z_∞ and let $\mathcal{E}_{+,i}$, $i = 1, \dots, r$, be the positive ends of $\cup_{j=0}^{a-1} \overline{v}^\sharp_{1,j}$ that converge to z_∞. Then*

$$n^*(\mathcal{E}_{-,i}) \geq k_0 - 1 \gg 2g \tag{3.4.4}$$

for each i, where the constant k_0 is as given in Section I.5.2.2.

If there are no boundary punctures that are removable at z_∞, then

$$n^*((\cup_{i=1}^q \mathcal{E}_{-,i}) \cup (\cup_{i=1}^r \mathcal{E}_{+,i})) \geq m - p_+, \tag{3.4.5}$$

where $p_+ = \deg(\overline{v}'_+)$, and

$$D_{\rho_0} - (\cup_{i=1}^q \pi_{D_{\rho_0}}(\mathcal{E}_{-,i})) \cup (\cup_{i=1}^r \pi_{D_{\rho_0}}(\mathcal{E}_{+,i}))$$

consists of at most p_+ thin sectors.

Proof. We consider the $\pi_{D_{\rho_0}}$-projections of the positive and negative ends of $\overline{v}_{1,j}$, $j > 0$, and the positive ends of $\overline{v}_{1,0}$ that limit to z_∞. Equation (3.4.4) follows immediately from looking at the projections $\pi_{D_{\rho_0}}(\mathcal{E}_{-,i})$ as in Lemma I.7.5.2.

We start at a nontrivial negative end of some $\overline{v}_{1,j_1}$ limiting to z_∞. Without loss of generality we call it $\mathcal{E}_{-,1}$. Then

$$\pi_{D_{\rho_0}}(\mathcal{E}_{-,1}) = \mathfrak{S}(\overline{h}(\overline{a}_{k_1,l_1}), \overline{a}_{k_2,l_2}),$$

where $\mathfrak{S}(A,B)$ denotes a counterclockwise sector from the radial ray A to the radial ray B.

We analyze the continuation

$$g^1_{j_1-1,1}, \dots, g^1_{1,1}, g_{0,1}, g^0_{1,1}, \dots, g^0_{a,1}$$

of $g^1_{j_1,1} \supset \partial_1\mathcal{E}_{-,1}$ in the direction of $\partial_+ B_\tau$. Here $\partial_k\mathcal{E}_{-,i}$, $k = 0,1$, is the $t = k$ boundary of $\mathcal{E}_{-,i}$, and $f^{*_1}_{*_2}$ corresponds to $g^{*_1}_{*_2}$ as in Definition 3.4.3.

(i) Suppose there is some $0 \leq j < j_1$ such that $g^1_{j,1}$ is nontrivial. Let $j_2 \geq 0$ be the first such occurrence in the continuation. Then $\overline{v}^\sharp_{1,j_2}$ has some nontrivial end which we call $\mathcal{E}_{+,1}$, such that

$$\pi_{D_{\rho_0}}(\mathcal{E}_{+,1}) = \mathfrak{S}(\overline{a}_{k_2,l_2}, \overline{h}(\overline{a}_{k_3,l_3})).$$

(ii) On the other hand, if $g^1_{j,1}$ is trivial for all $j_1 \geq j \geq 0$, then we set $j_2 = 0$ and $\overline{h}(\overline{a}_{k_2,l_2}) = \overline{h}(\overline{a}_{k_3,l_3})$ and skip $\mathfrak{S}(\overline{a}_{k_2,l_2}, \overline{h}(\overline{a}_{k_3,l_3}))$.

Next we consider the continuation

$$g^0_{j_2+1,2}, \dots, g^0_{a,2}$$

of $g^0_{j_2,2}$ in the direction of $\partial_+ B_\tau$. Here $g^0_{j_2,2} \supset \partial_0\mathcal{E}_{+,1}$ if (i) holds; and $j_2 = 0$, $g_{0,2} = g_{0,1}$, and $f_{0,2} = f_{0,1}$ if (ii) holds. There must exist some nontrivial $g^0_{j,2}$, $j > j_2$, and we write j_3 for the first such occurrence in the continuation. Then $\overline{v}^\sharp_{1,j_3}$ has some nontrivial end which we call $\mathcal{E}_{-,2}$, such that

$$\pi_{D_{\rho_0}}(\mathcal{E}_{-,2}) = \mathfrak{S}(\overline{h}(\overline{a}_{k_3,l_3}), \overline{a}_{k_4,l_4}).$$

Continuing in the same manner, the sectors sweep out a neighborhood of z_∞ in D_{ρ_0}, with the possible exception of p_+ thin sectors. This gives Equation (3.4.5). □

Lemma 3.4.5. *If $\mathfrak{p} \in \partial F_*$ is a boundary puncture of $\overline{v}_*$ which is removable at z_∞, then $n^*(\overline{v}_*(N(\mathfrak{p}))) \geq k_0 - 1 \gg 2g$, where $N(\mathfrak{p}) \subset F_*$ is a sufficiently small neighborhood of $\mathfrak{p}$.*

Proof. The proof is the same as that of Lemma I.7.5.4. □

3.4.2. *Bounds on ECH indices.* The goal of this subsection is to show the non-negativity of $I(\overline{v}_*)$ except when $\overline{v}_* = \overline{v}_+$, *under the assumption that there are no boundary punctures that are removable at* z_∞; see Lemma 3.4.9. The main new difficulty is to show that $I(\overline{v}_*) \geq 0$ for $\overline{v}_* \succ \overline{v}_+$. If $\overline{v}_* \succ \overline{v}_+$ and $\overline{v}'_* \cup \overline{v}^\sharp_* \neq \varnothing$, then we need to apply the version of the ECH index inequality in the presence of ends that limit to z_∞ (Lemma I.5.7.18). To apply Lemma I.5.7.18, we need to verify a certain "alternating property" for the ends of $\overline{v}_*$ that limit to z_∞; this is the content of Lemma 3.4.8.

Let $A_\varepsilon = \partial D_\varepsilon \times [0,1]$ for $0 < \varepsilon < \rho_0$ small and let $\pi_{[0,1]\times\overline{S}}$ be the projection of $\overline{W}$ or the positive end of $\overline{W}_\tau$ to $[0,1]\times\overline{S}$. By the proof of Lemma 3.4.4, the intersection $\mathfrak{c} := \pi_{[0,1]\times\overline{S}}(\cup_i \mathcal{E}_{-,i}) \cap A_\varepsilon$ is groomed and the sets P_0 and P_1 of initial and terminal points of $\mathfrak{c}$ alternate along $(0, 2\pi)$.

Definition 3.4.6. A *cycle* $\mathcal{Z} = (\mathfrak{z}_1 \to \mathfrak{z}_2 \to \cdots \to \mathfrak{z}_k \to \mathfrak{z}_1)$ *on* $\partial D_\varepsilon = \mathbb{R}/2\pi\mathbb{Z}$ is a sequence of points $\mathfrak{z}_i \in \mathbb{R}/2\pi\mathbb{Z}$, together with chords in $\mathbb{R}/2\pi\mathbb{Z}$ from $\mathfrak{z}_i$ to $\mathfrak{z}_{i+1}$, where the indices are taken modulo k.

The continuation method from Lemma 3.4.4 gives a cycle

$$\mathcal{Z} = (\mathfrak{z}_{10} \to \mathfrak{z}_{11} \to \cdots \to \mathfrak{z}_{k0} \to \mathfrak{z}_{k1} \to \mathfrak{z}_{10}),$$

where $P_i = \{\mathfrak{z}_{1i}, \dots, \mathfrak{z}_{ki}\}$, the chords correspond to the sectors, the cycle winds around $\mathbb{R}/2\pi\mathbb{Z}$ once, and each point of P_i appears only once in $\mathcal{Z}$. Note that some of the $\mathfrak{z}_{i0} \to \mathfrak{z}_{i1}$ may correspond to sectors $\mathfrak{S}(\overline{a}_{k',l'}, \overline{h}(\overline{a}_{k',l'}))$ of the type that are skipped in Step (ii) of Lemma 3.4.4.

Next let $\overrightarrow{\mathcal{D}}_{\pm,j}$ be the data at z_∞ for the ± end of $\overline{v}_{1,j}$ and let $P_{\pm,j,0}$ and $P_{\pm,j,1}$ be the initial and terminal points on A_ε determined by $\overrightarrow{\mathcal{D}}_{\pm,j}$. Then we write

$$P_{\pm,j,i} = P'_{\pm,j,i} \sqcup P''_{\pm,j,i}, \tag{3.4.6}$$

where $P'_{\pm,j,i}$ corresponds to $\overline{v}'_{1,j}$ and $P''_{\pm,j,i}$ corresponds to $\overline{v}''_{1,j}$. Note that

$$P_{+,j-1,i} = P_{-,j,i} \text{ and } P'_{+,j-1,i} = P'_{-,j-1,i}. \tag{3.4.7}$$

Definition 3.4.7. Let P be a finite subset of $S^1 = \mathbb{R}/2\pi\mathbb{Z}$. If $[a] \neq [b] \in P$, then we write $[a] \lessdot_P [b]$ if there exist $a \in [a]$, $b \in [b]$ such that $a < b < a + 2\pi$ and there are no representatives of P in the open interval (a, b).[7]

Lemma 3.4.8. *For each* $* = (\pm, j)$, $P_{*,0} \subset P_0$ *and* $P_{*,1} \subset P_1$ *and the points of* $P_{*,0}$ *and* $P_{*,1}$ *alternate along* $(0, 2\pi)$.

Proof. The proof is by induction on the level; see Figure 8 for an example. We will inductively define $P_i^{(0)} \supset P_i^{(1)} \supset P_i^{(2)} \supset \dots$ and the corresponding cycles $\mathcal{Z}^{(0)}, \mathcal{Z}^{(1)}, \dots$ and show that the following hold for each $j = 0, 1, \dots$:

($j0$) the points of $P_0^{(j)}$ and $P_1^{(j)}$ alternate along $(0, 2\pi)$;

($j1$) $P_{-,j_0-j,i} = P_{+,j_0-j-1,i} \subset P_i^{(j)}$, $i = 0, 1$;

---

[7] The relation $\lessdot_P$ is not symmetric, i.e., $[a] \lessdot_P [b]$ does not necessarily imply that $[b] \lessdot_P [a]$.

($j2$) there is a partition of $P_{-,j_0-j,0}\cup P_{-,j_0-j,1}$ into pairs $\{\mathfrak{p}_0,\mathfrak{p}_1\}$, $\mathfrak{p}_i\in P_{-,j_0-j,i}$, such that $\mathfrak{p}_0\lessdot_{P_0^{(j)}\cup P_1^{(j)}}\mathfrak{p}_1$; in particular, the points of $P_{-,j_0-j,0}$ and $P_{-,j_0-j,1}$ alternate along $(0,2\pi)$;

($j3$) $P'_{+,j_0-j-1,i}=P'_{-,j_0-j-1,i}\subset P_i^{(j+1)}$, $i=0,1$;

($j4$) there is a partition of $P'_{+,j_0-j-1,0}\cup P'_{+,j_0-j-1,1}$ into pairs $\{\mathfrak{p}_0,\mathfrak{p}_1\}$, $\mathfrak{p}_i\in P'_{+,j_0-j-1,i}$, such that $\mathfrak{p}_0\lessdot_{P_0^{(j+1)}\cup P_1^{(j+1)}}\mathfrak{p}_1$; in particular, the points of $P'_{+,j_0-j-1,0}$ and $P'_{+,j_0-j-1,1}$ alternate along $(0,2\pi)$.

Note that $P_{-,j_0-j,i}=P_{+,j_0-j-1,i}$ in ($j1$) and $P'_{+,j_0-j-1,i}=P'_{-,j_0-j-1,i}$ in ($j3$) by Equation (3.4.7).



FIGURE 8. Each rectangular box, with the sides identified, is A_ε; the top is $\partial D_\varepsilon\times\{1\}$ and the bottom is $\partial D_\varepsilon\times\{0\}$. For each figure, $P^\star_{*,0}$ and $P^\star_{*,0}$ are the sets of initial and terminal points of the arcs drawn, where the arcs are always oriented from bottom to top. The extra dots are drawn only for reference. The kth row corresponds to Step k. The solid and dashed arcs in the left column together indicate $\mathcal{Z}^{(0)}$ and $\mathcal{Z}^{(1)}$. The dashed arcs are potential arcs whose endpoints give pairs of points in $P''_{+,j,0}$ and $P''_{+,j,1}$.

Step 1. Let $\overline{v}_{1,j_0}$ be the highest level which has a negative end at z_∞, let $P_i^{(0)}=P_i$, $i=0,1$, and let $\mathcal{Z}^{(0)}=\mathcal{Z}$. The properties of $\mathcal{Z}^{(0)}$ (which in turn follow from the proof of Lemma 3.4.4) imply (00)–(02).

Step 2. We consider the positive ends of $\overline{v}_{1,j_0-1}$ that limit to z_∞. There is a partition of $P''_{+,j_0-1,0}\cup P''_{+,j_0-1,1}$ into pairs $\{\mathfrak{p}_0,\mathfrak{p}_1\}$, $\mathfrak{p}_i\in P''_{+,j_0-1,i}$, such that $\mathfrak{p}_1\lessdot_{P_0^{(0)}\cup P_1^{(0)}}\mathfrak{p}_0$, i.e., there is a chord $\mathfrak{p}_1\to\mathfrak{p}_0$ in $\mathcal{Z}^{(0)}$. Let $P_i^{(1)}=P_i^{(0)}-P''_{+,j_0-1,i}$ and let $\mathcal{Z}^{(1)}$ be obtained from $\mathcal{Z}^{(0)}$ by inductively replacing $\mathfrak{q}\to\mathfrak{p}_1\to\mathfrak{p}_0\to\mathfrak{q}'$ by $\mathfrak{q}\to\mathfrak{q}'$, given by concatenation. This makes sense since each point of $P_i^{(0)}$

appears only once in $\mathcal{Z}^{(0)}$. Then $P_0^{(1)} \cup P_1^{(1)}$ is the set of (alternating) points of $\mathcal{Z}^{(1)}$, each point of $P_i^{(1)}$ appears only once in $\mathcal{Z}^{(1)}$, and $\mathcal{Z}^{(1)}$ winds around $\mathbb{R}/2\pi\mathbb{Z}$ once. Then (10) follows from the description of $\mathcal{Z}^{(1)}$. Similarly, since $P'_{+,j_0-1,i} = P_{+,j_0-1,i} - P''_{+,j_0-1,i}$ and (02) holds, (03) and (04) follow immediately.

Step 3. We consider the negative ends of $\overline{v}_{1,j_0-1}$ that limit to z_∞. There is a partition of $P''_{-,j_0-1,0} \cup P''_{-,j_0-1,1}$ into pairs $\{\mathfrak{p}_0, \mathfrak{p}_1\}$, $\mathfrak{p}_i \in P''_{-,j_0-1,i}$, such that $\mathfrak{p}_0 \lessdot_{P_0^{(1)} \cup P_1^{(1)}} \mathfrak{p}_1$, i.e., there is a chord $\mathfrak{p}_0 \to \mathfrak{p}_1$ in $\mathcal{Z}^{(1)}$. Since $P_{-,j_0-1,i} = P'_{-,j_0-1,i} \cup P''_{-,j_0-1,i}$ and (04) holds, (11) and (12) follow immediately.

Repeated application of the above then implies the lemma. □

Lemma 3.4.9. *If fiber components are removed from $\overline{u}_\infty$ and there are no boundary punctures that are removable at z_∞, then the ECH index of each level $\overline{v}_* \neq \overline{v}_+$ is nonnegative, the only components of $\overline{u}_\infty$ which have negative ECH index are the branched covers of σ_∞^+, and $I(\overline{v}_{1,j}) \geq I(\overline{v}'_{1,j}) + I(\overline{v}''_{1,j})$ for $0 \leq j \leq a$, with equality for $j > 0$.*

Proof. The proof is analogous to that of Lemma I.7.6.5. By [HT1, Proposition 7.15(a)] and the regularity of $\{\overline{J}_\tau\}$, we have $I(\overline{v}_{0,j}) \geq 0$ for $1 \leq j \leq b$, where equality holds if and only if $\overline{v}_{0,j}$ is a connector. Also note that the only section at infinity that has negative ECH index is σ_∞^+ by Equations (3.2.5) and (3.2.6); this implies that $I(\overline{v}'_+) < 0$ if $\overline{v}'_+$ is nonempty.

We claim that $I(\overline{v}_{1,j}) \geq 0$ for $0 < j \leq a$ and $I(\overline{v}_{1,j}) \geq I(\overline{v}'_{1,j}) + I(\overline{v}''_{1,j})$ for $0 \leq j \leq a$, with equality for $j > 0$. Suppose z_∞ does not appear at an end of $\overline{v}_{1,j}$. With the possible exception of fiber components, $\overline{v}_{1,j}$ is simply-covered and regular since there is at least one HF end. Hence the claim follows from the regularity of $\{\overline{J}_\tau\}$ and the usual index inequality (Lemmas I.4.5.13 and I.5.6.9).

On the other hand, if z_∞ appears at an end of $\overline{v}_{1,j}$, then we use the version of the index inequality given by Lemma I.5.7.18. Recall that $\mathcal{E}_{-,i}$, $i = 1, \dots, q$, are the negative ends of $\cup_{j=1}^{a} \overline{v}_{1,j}^\sharp$ that converge to z_∞ and $\mathcal{E}_{+,i}$, $i = 1, \dots, r$, are the positive ends of $\cup_{j=0}^{a-1} \overline{v}_{1,j}^\sharp$ that converge to z_∞. Let $I_{-,j}$ be the subset of $\{1, \dots, q\}$ such that $\mathcal{E}_{-,i}$ is a negative end of $\overline{v}_{1,j}^\sharp$ if and only if $i \in I_{-,j}$, and let $I_{+,j}$ be the subset of $\{1, \dots, r\}$ such that $\mathcal{E}_{+,i}$ is a positive end of $\overline{v}_{1,j}^\sharp$ if and only if $i \in I_{+,j}$. Then let $\mathfrak{c}_{\pm,j}$ (resp. $\mathfrak{c}'_{\pm,j}$, $\mathfrak{c}''_{\pm,j}$) be the groomings corresponding to the ± ends of $\overline{v}_{1,j}$ (resp. $\overline{v}'_{1,j}$, $\overline{v}''_{1,j}$) at z_∞, subject to the following:

(α_1) $\mathfrak{c}_{\pm,j} = \mathfrak{c}'_{\pm,j} \cup \mathfrak{c}''_{\pm,j}$;
(α_2) $\mathfrak{c}_{-,j}$ has winding $w(\mathfrak{c}_{-,j}) \geq 0$ and $\mathfrak{c}_{+,j}$ has winding $w(\mathfrak{c}_{+,j}) \leq 0$;
(α_3) $\mathfrak{c}''_{-,j} = \pi_{[0,1]\times\overline{S}}(\cup_{k\in I_{-,j}} \mathcal{E}_{-,k}) \cap A_\varepsilon$ and $\mathfrak{c}''_{+,j} = \pi_{[0,1]\times\overline{S}}(\cup_{k\in I_{+,j}} \mathcal{E}_{+,k}) \cap A_\varepsilon$.

By Lemma 3.4.8, the sets $P_{\pm,j,0}$ and $P_{\pm,j,1}$ of initial and terminal points of $\mathfrak{c}_{\pm,j}$ alternate along $(0, 2\pi)$. Hence, by Lemma I.5.7.18,

$$I(\overline{v}_{1,j}) \geq I(\overline{v}'_{1,j}) + I(\overline{v}''_{1,j}) \geq \operatorname{ind}(\overline{v}''_{1,j}) \geq 0$$

with equality for $j > 0$. This proves the claim.

The case of $\overline{v}_{-1,j}$, $1 \leq j \leq c+1$, is similar, and the lemma follows. $\square$

3.4.3. *Boundary punctures.* In this subsection we describe the necessary modifications when $\overline{u}_\infty$ has boundary punctures that are removable at z_∞.

Suppose the following for simplicity:

(S) There is only one boundary puncture $\mathfrak{r} \in F''_{*_0}$ which is removable at z_∞ and $\overline{v}''_{*_0}(\mathfrak{r}) \in \{(0,1)\} \times \overline{\mathbf{a}} \subset \overline{W} = \mathbb{R} \times [0,1] \times \overline{S}$.

For the purposes of computing the Fredholm and ECH indices, we make the following modifications which allows us to use the considerations from Section I.5.7. We view the base $B = \mathbb{R} \times [0,1]$ with the puncture $(0,1)$ as a two-level building consisting of a disk B_1° with three boundary punctures and a disk B_2° with one boundary puncture. (What we are doing here is bubbling off a neighborhood of the puncture $(0,1) \in B$ when taking the limit $\overline{u}_i \to \overline{u}_\infty$.) All the punctures are viewed as strip-like ends: the punctures of B_1° are called the positive, left negative, and right negative ends, corresponding to the positive end, $(0,1)$, and the negative end of B, and the puncture of B_2° is called the positive end and is identified with the left negative end of B_1°. See Figure 9. We write $\partial B_1^\circ = \sqcup_{i=1}^3 \partial_i B_1^\circ$, arranged in counterclockwise order, such that $\partial_1 B_1^\circ$, $\partial_2 B_1^\circ$, and ∂B_2° correspond to $\mathbb{R} \times \{1\}$ and $\partial_3 B_1^\circ$ corresponds to $\mathbb{R} \times \{0\}$. The cobordism $\overline{W} = B \times \overline{S}$ decomposes into $\overline{W}_1^\circ = B_1^\circ \times \overline{S}$ and $\overline{W}_2^\circ = B_2^\circ \times \overline{S}$, the Lagrangian submanifold $\mathbb{R} \times \{1\} \times \overline{\mathbf{a}}$ decomposes into $\partial_1 B_1^\circ \times \overline{\mathbf{a}}$, $\partial B_2^\circ \times \overline{\mathbf{a}}$, and $\partial_2 B_1^\circ \times \overline{\mathbf{a}}$, and the Lagrangian submanifold $\mathbb{R} \times \{0\} \times \overline{h}(\overline{\mathbf{a}})$ corresponds to $\partial_3 B_1^\circ \times \overline{h}(\overline{\mathbf{a}})$.



FIGURE 9

Denote the sections at infinity of $\overline{W}_i^\circ$, $i = 1,2$, by $\sigma_\infty^{\circ,i} = B_i^\circ \times \{z_\infty\}$. A curve $\overline{v}_{*_0}$ in $\overline{W}$ decomposes into $\overline{v}_{*_0}^{\circ,1} \cup \overline{v}_{*_0}^{\circ,2}$, where $\overline{v}_{*_0}^{\circ,i} = \overline{v}_{*_0}^{\circ,i,'} \cup \overline{v}_{*_0}^{\circ,i,''}$, $i = 1,2$, is a curve in $\overline{W}_i^\circ$, $\overline{v}_{*_0}^{\circ,i,'}$ is a possibly branched cover of $\sigma_\infty^{\circ,i}$, and $\overline{v}_{*_0}^{\circ,2,''}$ is a union of constant sections $B_1^\circ \times \{pt\}$, $pt \in \widehat{\mathbf{a}}$. We view the chords at the positive ends of $\overline{v}_{*_0}^{\circ,2}$ (including z_∞) as negative Morse-Bott Reeb chords. The reader can verify that:

$$\operatorname{ind}(\sigma_\infty^{\circ,i}) = 0, \quad \operatorname{ind}(\overline{v}_{*_0}^{\circ,2,''}) = 0, \quad \operatorname{ind}(\overline{v}''_{*_0}) = \operatorname{ind}(\overline{v}_{*_0}^{\circ,1,''}). \tag{3.4.8}$$

We apply the continuation method to $\overline{v}_{1,j}$, $0 \leq j \leq a$, $(1,j) \neq *_0$, and $\overline{v}_{*_0}^{\circ,i}$, $i = 1,2$, to obtain a sequence of sectors, where the sectors are of type $\mathfrak{S}(\overline{a}_{k,l}, \overline{a}_{k',l'})$ with $(k,l) \neq (k',l')$ (corresponding to the left negative end of $\overline{W}_1^\circ$), $\mathfrak{S}(\overline{h}(\overline{a}_{k,l}), \overline{a}_{k',l'})$, or $\mathfrak{S}(\overline{a}_{k',l'}, \overline{h}(\overline{a}_{k,l}))$. This gives rise to a cycle

$$\mathcal{Z} = (\mathfrak{z}_1 \to \mathfrak{z}_2 \to \cdots \to \mathfrak{z}_k \to \mathfrak{z}_1)$$

which winds around $\mathbb{R}/2\pi\mathbb{Z}$ once. We leave it to the reader to make the proper generalizations of $\mathcal{Z}$ when we do not assume (S), i.e., there are several punctures that are removable at z_∞; $*_0 = +$ or $(-1, j_0)$, $1 \leq j_0 \leq c+1$; and/or the puncture lies on $L_{\overline{h}(\overline{\mathbf{a}})}$, $L_{\overline{\mathbf{a}}}^+$, $L_{\overline{\mathbf{b}}}$, $L_{\overline{h}(\overline{\mathbf{b}})}$, or $L_{\overline{\mathbf{b}}}^-$.

We identify the positive end of $\overline{W}_1^\circ$ with $[0,\infty) \times [0,1] \times \overline{S}$, the left negative end with $(-\infty, 0] \times [\frac{1}{2}, 1] \times \overline{S}$, and the right negative end with $(-\infty, 0] \times [0, \frac{1}{2}] \times \overline{S}$. Let $A_\varepsilon^{[a,b]}$ be the annulus $\{\rho = \varepsilon\} \subset [a,b] \times D_{\rho_0}$. Writing $*_0 = (1, j_0)$, let $\overrightarrow{\mathcal{D}}_{+,j_0}$, $\overrightarrow{\mathcal{D}}_{L-,j_0}$, $\overrightarrow{\mathcal{D}}_{R-,j_0}$ be the data at z_∞ of the positive, left negative, and right negative ends, and let $P_{L-,j_0,1/2}$ and $P_{L-,j_0,1}$ be the initial and terminal points of $A_\varepsilon^{[1/2,1]}$ determined by $\overrightarrow{\mathcal{D}}_{L-,j_0}$, etc. We also decompose $P_{*,i} = P'_{*,i} \cup P''_{*,i}$, $* = (L-, j_0)$, $(R-, j_0)$, or $(+, j_0)$, as before so that $P'_{*,i}$ corresponds to $\overline{v}_*^{\circ,1,'}$ and $P''_{*,i}$ corresponds to $\overline{v}_*^{\circ,1,''}$.

Definition 3.4.10. We classify boundary punctures $\mathfrak{r} \in \partial F_{*_0}$, $*_0 = (1, j_0)$, $0 \leq j_0 \leq a$, which are removable at z_∞ into three types:

- (P_1) $\overline{v}'_{*_0} = \varnothing$;
- (P_2) $\overline{v}'_{*_0} \neq \varnothing$ and all the vertices of $\mathcal{Z}$ correspond to points of type $\overline{a}_{k,l}$ or all the vertices of $\mathcal{Z}$ correspond to points of type $\overline{h}(\overline{a}_{k,l})$;
- (P_3) $\overline{v}'_{*_0} \neq \varnothing$ and $\mathcal{Z}$ has vertices of both types $\overline{a}_{k,l}$ and $\overline{h}(\overline{a}_{k,l})$.

Here we are allowing the possibility of multiple punctures. Boundary punctures $\mathfrak{r} \in \partial F_{*_0}$, $*_0 = (-1, j_0)$, $1 \leq j_0 \leq c+1$, are classified similarly.

Remark 3.4.11. The existence of boundary punctures of type (P_1), (P_2), and (P_3) are mutually exclusive.

The following is a strengthening of Lemma 3.4.4 when $\overline{v}'_{1,j} \cup \overline{v}^\sharp_{1,j} \neq \varnothing$ for some $j > 0$ and there are boundary punctures of type (P_3) that are removable at z_∞:

Lemma 3.4.12. *Suppose $\overline{v}'_{1,j} \cup \overline{v}^\sharp_{1,j} \neq \varnothing$ for some $j > 0$. Let $\mathcal{E}_{-,i}$, $i = 1, \dots, q$, be the negative ends of $\cup_{j=1}^a \overline{v}^\sharp_{1,j}$ that converge to z_∞, let $\mathcal{E}_{+,i}$, $i = 1, \dots, r$, be the positive ends of $\cup_{j=0}^{a-1} \overline{v}^\sharp_{1,j}$ that converge to z_∞, and let $\mathcal{E}'_i$, $i = 1, \dots, s$, be the neighborhoods of the punctures of type (P_3). Then*

$$n^*((\cup_{i=1}^q \mathcal{E}_{-,i}) \cup (\cup_{i=1}^r \mathcal{E}_{+,i}) \cup (\cup_{i=1}^s \mathcal{E}'_i)) \geq m - p_+. \tag{3.4.9}$$

Proof. Similar to that of Lemma 3.4.4. □

Lemma 3.4.13. *If $\mathfrak{p}_1, \dots, \mathfrak{p}_s$ are the punctures of type (P_1) and (P_2) and $N(\mathfrak{p}_i) \subset F_*$ is a small neighborhood of $\mathfrak{p}_i$, then $\sum_{i=1}^s n^*(\overline{v}_*(N(\mathfrak{p}_i))) \geq m$.*

Proof. (P_2) follows immediately from the description of $\mathcal{Z}$; (P_1) is similar. □

3.4.4. *Bounds on ECH indices, part II.* In this subsection we give bounds on ECH indices in the presence of boundary punctures that are removable at z_∞.

For simplicity we are still assuming (S). The ECH indices of $\overline{v}_{*_0}^{\circ,i}$ are defined in a manner similar to that of Definition I.5.7.2.

Lemma 3.4.14. *If* $\overline{v}'_{*_0}, \overline{v}''_{*_0} \neq \varnothing$*, (S) holds, and the boundary puncture of type (P*$_3$*), then*

$$I(\overline{v}_{*_0}) = I(\overline{v}_{*_0}^{\circ,1,'}) + I(\overline{v}_{*_0}^{\circ,1,''}) + I_+ + I_{L-} + I_{R-}, \tag{3.4.10}$$

where $I_+ \geq 0$, $I_{L-} = 1$, *and* $I_{R-} \geq 0$ *are contributions from the ends that limit to* z_∞ *at the positive, left negative, and right negative ends.*

Proof. The calculation of I_{L-} is similar to the ECH index calculations of Section I.5.7. First observe that $P_{L-,j_0,1/2} = P_{L-,j_0,1}$. By (S), $\mathcal{Z}$ has only one chord $\mathfrak{z}'_1 \to \mathfrak{z}'_2$ corresponding to a sector of type $\mathfrak{S}(\overline{a}_{k,l}, \overline{a}_{k',l'})$. Observe that there are no points of $\mathcal{Z}$ between $\mathfrak{z}'_1$ and $\mathfrak{z}'_2$; otherwise $\mathcal{Z}$ winds more than once around $\mathbb{R}/2\pi\mathbb{Z}$. Hence $P''_{L-,j_0,1/2} = \{\mathfrak{z}'_1\}$, $P'_{L-,j_0,1/2} = \{\mathfrak{z}'_2, \mathfrak{z}''_1, \dots, \mathfrak{z}''_s\}$, $P''_{L-,j_0,1} = \{\mathfrak{z}'_2\}$, and $P'_{L-,j_0,1} = \{\mathfrak{z}'_1, \mathfrak{z}''_1, \dots, \mathfrak{z}''_s\}$, where $P_{L-,j_0,1/2} = P_{L-,j_0,1}$ is written as $\{\mathfrak{z}'_1, \mathfrak{z}'_2, \mathfrak{z}''_1, \dots, \mathfrak{z}''_s\}$ in cyclic order around $\mathbb{R}/2\pi\mathbb{Z}$, and the projection of the left negative end of $\overline{v}_{*_0}^{\circ,1,''}$ that limits to z_∞ intersects $A_\varepsilon^{[1/2,1]}$ along an arc $\mathfrak{c}''_{L-}$ with winding number $w(\mathfrak{c}''_{L-}) = 0$ or 1, depending on whether $\mathfrak{S}(\overline{a}_{k,l}, \overline{a}_{k',l'})$ is a large sector. The left negative ends of $\overline{v}_{*_0}^{\circ,1,'}$ that limit to z_∞ can modified to give a grooming $\mathfrak{c}'_{L-}$ on $A_{\varepsilon/2}^{[1/2,1]}$ such that the winding number $w(\mathfrak{c}'_{L-}) = 0$ or -1 and $\mathfrak{c}'_{L-}$ connects $\mathfrak{z}''_i$ to $\mathfrak{z}''_i$, $i = 1, \dots, s$, by vertical arcs. Then the writhe of $\mathfrak{c}'_{L-} \cup \mathfrak{c}''_{L-}$ is $+1$ and resolving the positive crossing of $\mathfrak{c}'_{L-} \cup \mathfrak{c}''_{L-}$ yields a grooming by vertical arcs from $P_{L-,j_0,1/2}$ to $P_{L-,j_0,1}$. Hence $I_{L-} = 1$.

Finally, since $I(\overline{v}_{*_0}) = I(\overline{v}_{*_0}^{\circ,1}) + I(\overline{v}_{*_0}^{\circ,2})$ and $I(\overline{v}_{*_0}^{\circ,2}) = 0$, we have $I(\overline{v}_{*_0}) = I(\overline{v}_{*_0}^{\circ,1}) = I(\overline{v}_{*_0}^{\circ,1,'}) + I(\overline{v}_{*_0}^{\circ,1,''}) + I_+ + I_{L-} + I_{R-}$. Lemma 3.4.9 implies that $I_+ \geq 0$ and $I_{R-} \geq 0$. $\square$

Remark 3.4.15. In general, each boundary puncture of type (P$_3$) contributes at least $+1$ towards I.

The following is a strengthening of Lemma 3.4.9 in the presence of boundary punctures:

Lemma 3.4.16. *If fiber components are removed from* $\overline{u}_\infty$ *and the only boundary punctures are of type (P*$_3$*), then the ECH index of each level* $\overline{v}_* \neq \overline{v}_+$ *is nonnegative, the only components of* $\overline{u}_\infty$ *which have negative ECH index are branched covers of* σ_∞^+, *and* $I(\overline{v}_{1,j}) \geq I(\overline{v}'_{1,j}) + I(\overline{v}''_{1,j}) + \mathfrak{bp}_{1,j}$ *for* $0 \leq j \leq a$. *Here* $\mathfrak{bp}_*$ *is the number of boundary punctures of type (P*$_3$*) on* $\overline{v}_*$.

Proof. We explain the modifications that need to be made when there are boundary punctures that are removable at z_∞. We assume (S) for simplicity. The cycles $\mathcal{Z}^{(a-j)}$ are defined as before, for $j \geq j_0$. We define $\mathcal{Z}^{(a-j_0,+)} = \mathcal{Z}^{(a-j_0)}$ and $\mathcal{Z}^{(a-j_0,-)}$ as $\mathcal{Z}^{(a-j_0,+)}$ with $\mathfrak{z}'_1 \to \mathfrak{z}'_2$ replaced by $\mathfrak{z}'_2$. Then $\mathcal{Z}^{(a-(j_0-1))}$ is obtained using $\mathcal{Z}^{(a-j_0,-)}$ instead of $\mathcal{Z}^{(a-j_0)}$. Also, $P^\star_{R-,j_0,i}$, $\star = \varnothing, ', ''$, is obtained from

$P^{\star}_{+,j_0,i}$ by replacing $\mathfrak{z}_1'$ by $\mathfrak{z}_2'$. The rest of the argument of Lemma 3.4.9 carries over. ☐

3.4.5. *Case of $\overline{v}_*' \cup \overline{v}_*^\sharp = \varnothing$ for all $\overline{v}_*$.*

Lemma 3.4.17. *If $\overline{u}_\infty \in \partial_{\{+\infty\}}\mathcal{M}$ and $\overline{v}_*' \cup \overline{v}_*^\sharp = \varnothing$ for all levels $\overline{v}_*$ of $\overline{u}_\infty$, then $a = c = 0$; $I(\overline{v}_+) = 0$; $I(\overline{v}_-) = 2$; $\overline{v}_+$ is a W_+-curve; $\overline{v}_-$ is a $\overline{W}_-$-curve; and there may be connectors $\overline{v}_{0,j}$ in between.*

Proof. Suppose that $\overline{u}_\infty \in \partial_{\{+\infty\}}\mathcal{M}$ and $\overline{v}_*' \cup \overline{v}_*^\sharp = \varnothing$ for all levels $\overline{v}_*$ of $\overline{u}_\infty$. Then there is a point $\mathfrak{q} \in int(F_-)$ and a sufficiently small neighborhood $N(\mathfrak{q}) \subset \dot{F}_-$ of $\mathfrak{q}$ such that $\overline{v}_-(\mathfrak{q}) = \overline{\mathfrak{m}}(\infty)$ and $n^*(\overline{v}_-(N(\mathfrak{q}))) \geq m$. By Equation (3.4.1) and Lemma 3.4.5, there are no boundary punctures that are removable at z_∞ and the only possible fiber component passes through $\overline{\mathfrak{m}}(\infty)$. Hence every level $\overline{v}_*$, $* \neq -$, has image inside W', W, or W_+ and $\overline{v}_-$ is a $\overline{W}_-$-curve or a *degenerate* $\overline{W}_-$-curve by the analog of Lemma I.7.6.2.

The ECH index of each level $\neq \overline{v}_+$ which has no fiber component is nonnegative by Lemma 3.4.9. Since $\overline{v}_+$ is a W_+-curve, $I(\overline{v}_+) \geq 0$. On other hand, by the previous paragraph, if there is a fiber component, then it is a component of $\overline{v}_-$. We claim that $I(\overline{v}_-) \geq 2$, with equality if and only if $\overline{v}_-$ is not a degenerate $\overline{W}_-$-curve. Indeed, if $\overline{v}_-$ is not degenerate, then $I(\overline{v}_-) \geq 2$ by the point constraint (this is the only place where we use the fact that $\overline{u}_\infty \in \partial_{\{+\infty\}}\mathcal{M}$) and the ECH index inequality, and if $\overline{v}_-$ is degenerate, then $I(\overline{v}_-) \geq 4$, as computed in the proof of Lemma I.7.6.5.

The lemma then follows from Equation (3.4.2). In particular, degenerate $\overline{W}_-$-curves are not allowed. ☐

3.4.6. *Preliminary restrictions when $\overline{v}_*' \cup \overline{v}_*^\sharp \neq \varnothing$ for some $\overline{v}_*$.* We now consider the case where $\overline{v}_*' \cup \overline{v}_*^\sharp \neq \varnothing$ for some level $\overline{v}_*$.

Lemma 3.4.18. *If $\overline{v}_*' \cup \overline{v}_*^\sharp \neq \varnothing$ for some level $\overline{v}_*$ of $\overline{u}_\infty$, then:*

(1) $p_- = \deg(\overline{v}_-') > 0$;
(2) *$\overline{u}_\infty$ has no boundary puncture of type (P$_1$) or (P$_2$);*
(3) *$\overline{u}_\infty$ has no fiber components and no components of $\overline{v}_*''$ that intersect the interior of a section at infinity;*
(4) *each component of $\overline{v}_{-1,j}^\sharp$, $j = 1, \dots, c$, is a thin strip from z_∞ to some x_i or x_i' with $I = 1$;*
(5) *each component of $\overline{v}_-^\sharp$ is a section of $\overline{W}_- - W_-$ from δ_0 to some x_i or x_i' with $I = 1$;*
(6) *each component of $\overline{v}_{0,j}^\sharp$, $j = 1, \dots, b$, which has a positive end at a multiple of δ_0 is a cylinder from δ_0 to h or e with $I = 1$ or 2;*
(7) *the only boundary punctures are type (P$_3$) punctures of $\overline{v}_{1,j}^\sharp$, $j = 0, \dots, a$.*

Proof. The lemma is a consequence of Equation (3.4.1). Suppose $\overline{v}_*' \cup \overline{v}_*^\sharp \neq \varnothing$ for some level $\overline{v}_*$. Then $\overline{v}_*^\sharp \neq \varnothing$ for some $\overline{v}_*$ and each end of $\overline{v}_*^\sharp$ that limits to some multiple of z_∞ or δ_0 contributes positively to n^* by Lemmas 3.4.1 and 3.4.12.

(1) If $\overline{v}'_- = \varnothing$, then the restriction of $\overline{v}''_-$ to a neighborhood of $\overline{\mathfrak{m}}(+\infty)$ contributes m towards $n^*(\overline{v}_-)$. This contradicts the discussion from the previous paragraph, proving (1).

Since $\overline{v}'_- \neq \varnothing$, some $\overline{v}^\sharp_{0,j}$, $j = 1, \dots, b+1$, or $\overline{v}^\sharp_{1,j}$, $j = 1, \dots, a$, has a negative end at z_∞ or a multiple of δ_0. By Lemmas 3.4.1(1) and 3.4.12, it follows that:

$$\sum_{\overline{v}_* \succ \overline{v}_-} n^*(\overline{v}_*) \geq m - 2g, \tag{3.4.11}$$

where we are counting contributions from the ends and the boundary punctures of type (P$_3$).

(2)–(7) are consequences of Equation (3.4.11). We explain (2) and (3), leaving the rest to the reader. By Lemma 3.4.13, the neighborhoods of the boundary punctures of type (P$_1$) or (P$_2$) contribute at least m towards $n^*(\overline{v}''_*)$ in total. Also, a non-ghost fiber component or a component of $\overline{v}''_*$ that intersects the interior of a section at infinity contributes m towards n^*. They both contradict Equation (3.4.11). □

Lemma 3.4.19. *If $\overline{v}'_* \cup \overline{v}^\sharp_* \neq \varnothing$ for some level $\overline{v}_*$, then the following alternative holds:*

(a) *either some $\overline{v}^\sharp_{0,j_0}$, $j_0 = 1, \dots, b+1$, has a negative end at a multiple of δ_0, in which case $\overline{v}'_{0,j_0} = \varnothing$, $\overline{v}'_* = \overline{v}^\sharp_* = \varnothing$ for all levels $\overline{v}_* \succ \overline{v}_{0,j_0}$, and $\overline{v}'_{0,j} \neq \varnothing$ for all levels $\overline{v}'_- \preceq \overline{v}_{0,j} \prec \overline{v}'_{0,j_0}$; or*

(b) *no $\overline{v}^\sharp_{0,j}$, $j = 1, \dots, b+1$, has a negative end at a multiple of δ_0, in which case $\overline{v}'_* \neq \varnothing$ for all levels $\overline{v}'_- \preceq \overline{v}'_* \preceq \overline{v}'_+$ and $\overline{v}'_{1,j} \cup \overline{v}^\sharp_{1,j} \neq \varnothing$ for some $j > 0$.*

Proof. This follows from Lemmas 3.4.1(1) and 3.4.12 by observing that either case contributes at least $m - 2g$ towards n^* and that it is not possible to have both since $m \gg 2g$. It is also not possible to have multiple occurrences of negative ends of $\overline{v}^\sharp_{0,j}$, $j > 0$, that converge to some multiple of δ_0. □

3.4.7. *List of possibilities when $\overline{v}'_* \cup \overline{v}^\sharp_* \neq \varnothing$ for some $\overline{v}_*$.* We first start with a useful definition.

Definition 3.4.20.

(1) Let $X_1 \cup \dots \cup X_r$ be an r-level building of almost complex manifolds with cylindrical ends, ordered from bottom to top, and let

$$v = v_1 \cup \dots \cup v_r, \quad \operatorname{Im}(v_i) \subset X_i,$$

be a corresponding r-level holomorphic building, where each level has finite energy. If $\mathcal{E}$ is some collection of positive ends of v_r, then the *holomorphic building hanging from $\mathcal{E}$* is the union of irreducible components $\widetilde{v}_{i_0}$ for which there exist irreducible components $\widetilde{v}_i$, $i = i_0 + 1, \dots, r$, where the limit of a positive end of $\widetilde{v}_i$ agrees with the limit of a negative end of $\widetilde{v}_{i+1}$ for all $i = i_0, \dots, r-1$ and a positive end of $\widetilde{v}_r$ is contained in $\mathcal{E}$.

(2) If $X_1 \cup \dots \cup X_{r+1}$ is an $(r+1)$-level building, $v = v_1 \cup \dots \cup v_{r+1}$ is a corresponding $(r+1)$-level holomorphic building and $\mathcal{E}$ is some collection of negative ends of v_{r+1}, then the *holomorphic building hanging from $\mathcal{E}$* is defined similarly.

(3) If $X_1 \cup \dots \cup X_r$ is an r-level building, $v = v_1 \cup \dots \cup v_r$ is a corresponding r-level holomorphic building and $\mathcal{E}$ is some collection of negative ends of v_1, then the *holomorphic building sitting above $\mathcal{E}$* is the union of irreducible components $\widetilde{v}_{i_0}$ for which there exist irreducible components $\widetilde{v}_i$, $i = 1, \dots, i_0 - 1$, where the limit of a negative end of $\widetilde{v}_i$ agrees with the limit of a positive end of $\widetilde{v}_{i-1}$ for all $i = 2, \dots, i_0$ and a negative end of $\widetilde{v}_1$ is contained in $\mathcal{E}$.

(4) If $X_0 \cup \dots \cup X_r$ is an $(r+1)$-level building, $v = v_0 \cup \dots \cup v_r$ is a corresponding $(r+1)$-level holomorphic building and $\mathcal{E}$ is some collection of positive ends of v_0, then the *holomorphic building sitting above $\mathcal{E}$* is defined similarly.

Lemma 3.4.21. *If $\overline{v}'_* \cup \overline{v}^\sharp_* \neq \varnothing$ for some level $\overline{v}_*$, then $\overline{v}'_- \neq \varnothing$, there are no removable punctures, and $\overline{u}_\infty$ contains one of the following subbuildings:*

(1) *A 3-level building consisting of a component of $\overline{v}^\sharp_{0,1}$ with $I = 1$ from γ to $\delta_0\gamma'$; $\overline{v}'_- = \sigma^-_\infty$; and a thin strip.*

(2_i) *A 3-level building consisting of a component of $\overline{v}^\sharp_+$ with $I = i$, $i = 0, 1$, from $\mathbf{y}$ or $\mathbf{y}''$ to $\delta_0\gamma'$; $\overline{v}'_- = \sigma^-_\infty$; and a thin strip.*

(3) *A 4-level building consisting of a component of $\overline{v}^\sharp_+$ with $I = 0$ from $\mathbf{y}$ to $\delta_0^2\gamma'$; $\overline{v}'_{0,1} = \sigma'_\infty$; a component of $\overline{v}^\sharp_{0,1}$ which is an $I = 1$ cylinder from δ_0 to h; $\overline{v}'_- = \sigma^-_\infty$; and a thin strip.*

(4) *A 4-level building consisting of a component of $\overline{v}^\sharp_+$ with $I = 0$ from $\mathbf{y}$ to $\delta_0^2\gamma'$; $\overline{v}'_{0,1}$ with $I = 0$ and $\deg = 2$; $\overline{v}'_- = \sigma^-_\infty$; a component of $\overline{v}^\sharp_-$ which is an $I = 1$ curve from δ_0 to some x_i or x'_i; and a thin strip.*

(5) *A ≥ 3-level building consisting of a component of $\overline{v}^\sharp_+$ with $I = 0$ from $\mathbf{y}$ to $\delta_0^2\gamma'$; $\overline{v}'_{0,1}$ and $\overline{v}'_-$ with $I = 0$ and $\deg = 2$, where $\overline{v}'_{0,1} = \varnothing$ is possible; and two thin strips.*

(6_i) *A ≥ 4-level building consisting of a component of $\overline{v}^\sharp_{1,1}$ with $I = i$, $i = 1, 2$, from $\mathbf{y}$ to $\{z^r_\infty\} \cup \mathbf{y}'$; $0 < p_{0,0} \leq p_{0,1} \leq \dots \leq p_{0,b+1} \leq r$; $I(\overline{v}'_+) = -p_{0,b+1}$; $I(\overline{v}'_{0,j}) = 0$ for $j = 1, \dots, b$; $\overline{v}^\sharp_+$ with $I = 2 - i$ which has no negative ends at a multiple of δ_0; $p_{0,b+1} - p_{0,1}$ cylinders of $\overline{v}^\sharp_{0,j}$, $j \in \{1, \dots, b\}$, with $I = 1$ each from δ_0 to h; $p_{0,1} - p_{0,0}$ components of $\overline{v}^\sharp_-$ with $I = 1$ each from δ_0 to some x_i or x'_i; and $p_{0,0}$ thin strips. The total ECH index of the building hanging from the negative end of $\overline{v}'_+$ is $p_{0,b+1}$.*

Here we are omitting levels which are connectors. If there is more than one thin strip, then the thin strips could be on the same level or on different levels.

See Figure 10.



FIGURE 10. Schematic diagrams for the possible types of degenerations. Here • represents δ_0, ○ represents z_∞, □ represents a possible location of a branch point and × represents some x_i or x_i'. A vertical line indicates the restriction of a trivial cylinder, a dotted vertical line indicates a trivial cylinder, a double vertical line indicates a degree 2 branched cover of a trivial cylinder or a restriction of a trivial cylinder, and a triple vertical line indicates a degree $p\geq 2$ branched cover of a trivial cylinder or a restriction of a trivial cylinder. The labels on the graphs are ECH indices of each component. All the thin strips of $\overline{v}^\sharp_{-1,j}$ are drawn on the same level for convenience.

Proof. The lemma is a consequence of Equations (3.4.1) and (3.4.2). By Lemmas 3.4.16 and 3.4.18(2), (3), all the levels $\overline{v}_*$ besides $\overline{v}_+$ have nonzero ECH index, the only components of $\overline{u}_\infty$ which have negative ECH index are the branched covers of σ^+_∞, and $I(\overline{v}_{1,j})=I(\overline{v}'_{1,j})+I(\overline{v}''_{1,j})+\mathfrak{bp}_{1,j}$ for $0\leq j\leq a$. In particular, all the components of $\overline{u}_\infty$ which are left out of the subbuildings in (1)–(6$_i$) and are not drawn in Figure 10 have $I\geq 0$.

By Lemma 3.4.18(1), $\overline{v}'_-\neq\varnothing$. By Lemma 3.4.18(2), (7), there are no removable punctures in Cases (1)–(5) and no removable punctures of types (P$_1$) and (P$_2$) in Case (6$_i$). Removable punctures of type (P$_3$) in Case (6$_i$) will be eliminated later.

First suppose that some $\overline{v}^\sharp_{0,j_0}$, $j_0\in\{1,\dots,b\}$, has a negative end at δ_0^p for some $p>0$. This is the situation of Lemma 3.4.19(a) with $j_0\in\{1,\dots,b\}$. In this case, all the components and levels have nonnegative ECH index. Each component of $\overline{v}^\sharp_*\prec\overline{v}_{0,j_0}$ described in Lemma 3.4.18(4)–(6) satisfies $I\geq 1$ and there must be $p>0$ such components since $\overline{v}_{0,j_0}$ has multiplicity p at δ_0. We then have the following contributions towards I:

(a) $\sum_{\overline{v}^\sharp_*\prec\overline{v}_{0,j_0}} I(\overline{v}^\sharp_*)\geq p$ and there is at least one thin strip.

(b) $I(\overline{v}^\sharp_{0,j_0})\geq 1$ since $\overline{v}_{0,j_0}$ is nontrivial.

Then $p = 1$ and all the components and levels besides those of (a) and (b) satisfy $I = 0$. We are in Case (1).

Next suppose that $\overline{v}^\sharp_{0,b+1} = \overline{v}^\sharp_+$ has a negative end at δ_0^p. This is the situation of Lemma 3.4.19(a) with $j = b + 1$. Since $\overline{v}'_+ = \varnothing$, all the components and levels satisfy $I \geq 0$. Here $\overline{v}^\sharp_+$ cannot have any multiple of z_∞ at the positive end, since otherwise we contradict Equation (3.4.1). Since $I(\overline{v}^\sharp_+) \geq 0$ and $\sum_{\overline{v}^\sharp_* \prec \overline{v}_+} I(\overline{v}^\sharp_*) \geq p$, either $p = 1$ and we are in Case (2_i) or $p = 2$ and we are in Cases (3)–(5).

Finally suppose that no $\overline{v}^\sharp_{0,j}$, $j = 1, \dots, b+1$, has a negative end at a multiple of δ_0. This is the situation of Lemma 3.4.19(b). We have the following contributions:

(α) $\deg \overline{v}'_+ = p_{0,b+1}$ and $I(\overline{v}'_+) = -p_{0,b+1}$.

(β) By Lemma 3.4.18(4)–(6), the holomorphic building hanging from the negative end of $\overline{v}'_+$ satisfies $\sum_{\overline{v}_* \prec \overline{v}_+} I(\overline{v}'_* \cup \overline{v}^\sharp_*) \geq p_{0,b+1}$, and equality holds if and only if there is no cylinder from δ_0 to e.

(γ) $I = 2$ components of $\overline{v}^\sharp_{0,j}$ from δ_0 to e can be eliminated by observing that it is followed by an $I = -1$ component of $\overline{v}^\flat_-$ from e to some x_i or x'_i, which is a contradiction.[8]

(δ) Some $\overline{v}^\sharp_{1,j_0}$, $j_0 \geq 0$, must have an end at z_∞ (or some boundary puncture of type (P_3) must have a neighborhood) which projects to a large sector of D_{ρ_0}. Since $\overline{v}^\sharp_{1,j_0}$ generically lies on a codimension 1 stratum of some $\operatorname{ind}(\overline{v}^\sharp_{1,j_0})$-dimensional moduli space, the large sector contributes an additional $+1$ to the Fredholm and ECH indices, cf. Lemma I.5.7.18.

(ε) If $\mathfrak{bp}$ is the number of boundary punctures of type (P_3), then the contribution to I is $\mathfrak{bp}$.

(α), (β), and (γ) together give:

$$I(\overline{v}'_+) + \sum_{\overline{v}_* \prec \overline{v}_+} I(\overline{v}'_* \cup \overline{v}^\sharp_*) = 0.$$

Now, $I(\overline{v}''_{1,j}) \geq 1$ for each $j > 0$,[9] (δ) contributes at least 1 to I, and (ε) contributes at least $\mathfrak{bp}$ to I. Hence $\mathfrak{bp} = 0$, $a = 1$, and we are in Case (6_i), $i = 1, 2$. $\square$

3.4.8. *Elimination of some cases when $\overline{v}'_* \cup \overline{v}^\sharp_* \neq \varnothing$ for some $\overline{v}_*$.* The goal of this subsection is to eliminate all but one of the possibilities (namely Case (2_1)) given in Lemma 3.4.21 and to prove Lemma 3.3.2.

Definition 3.4.22. If $\pi : \Sigma \to \Sigma'$ is a branched cover and $\mathcal{B}(\pi)$ is the branch locus of π, then we define

$$\mathfrak{b}(\pi) = \sum_{x \in \mathcal{B}(\pi)} (\deg(x) - 1),$$

---

[8] Strictly speaking, $\overline{J}_-$ is the restriction of a Morse-Bott $\overline{J'}$. To give a proper treatment of regularity, we must perturb $\overline{J'}$ using an arbitrarily small Morse function to obtain $(\overline{J'})^\Diamond$ and then restrict $(\overline{J'})^\Diamond$ to $\overline{W}_-$.

[9] Recall that we are ignoring connectors.

where $\deg(x)$ is the degree of the branch point x. We also write $\mathfrak{b}(\overline{u})=\mathfrak{b}(\overline{\pi}_*\circ\overline{u})$, where $\overline{\pi}_*$ is the projection to the base B, B', etc., and refer to it informally as "the number of branch points of $\overline{u}$."

Let $\mathcal{E}_{-,i}$, $i=1,\dots,q$, be the negative ends of $\cup_{j=1}^{a}\overline{v}^{\sharp}_{1,j}$ that converge to z_∞ and let $\mathcal{E}_{+,i}$, $i=1,\dots,r$, be the positive ends of $\cup_{j=0}^{a-1}\overline{v}^{\sharp}_{1,j}$ that converge to z_∞ as in Lemma 3.4.4.

Lemma 3.4.23. *Suppose we are in Case* (6_i) *of Lemma 3.4.21. Then* $\sum_{\overline{v}'_*}\mathfrak{b}(\overline{v}'_*)$, *including branched points of connector components, is equal to the number of negative ends* $\mathcal{E}_{-,i}$. *Moreover, if* F *is the surface obtained by gluing all the domains of* $\overline{v}'_*$, *then* F *is a connected planar surface whose compactification (after filling in the interior and boundary punctures) is a closed disk.*

Proof. Suppose we are in Case (6_i) of Lemma 3.4.21. We may make the following simplifying assumptions:

1. $\overline{v}_{1,1}$ is a connector;
2. all the ends $\mathcal{E}_{-,i}$ are negative ends of $\overline{v}^{\sharp}_{1,2}$;
3. all the ends $\mathcal{E}_{+,i}$ are positive ends of $\overline{v}^{\sharp}_{+}$;
4. all the branch points of $\overline{v}'_*$ lie on $\overline{v}'_{1,1}$.

(1)–(3) are immediate. Note that in Lemma 3.4.21 we omitted connectors, but in the present analysis we need to keep track of connectors which are branched covers. (4) is obtained by pushing all the branched points from the upper and lower levels to $\overline{v}'_{1,1}$; this operation is possible because $\overline{v}^{\sharp}_* \preceq \overline{v}_+$ does not have a negative end that limits to z_∞ in Case (6_i). This operation does not affect:

- the total Fredholm index $\sum_{\overline{v}'_*}\operatorname{ind}(\overline{v}'_*)$; and
- the topological type of F.

As a consequence of (1)–(4), each component of $\overline{v}'_*\preceq\overline{v}'_+$ is a degree 1 cover of the appropriate σ^*_∞.

We now compute the Fredholm index of $\overline{v}'_{1,1}:\dot{F}_{1,1}\to\overline{W}$ using the Fredholm index formula

$$\operatorname{ind}(\overline{v}'_{1,1})=-\chi(\dot{F}_{1,1})+p_{1,1}+\mu_\tau(\overline{v}'_{1,1})+2c_1((\overline{v}'_{1,1})^*T\overline{S},\tau);$$

see Equation (I.5.7.1). Recall we are writing $p_*=\deg(\overline{v}'_*)$. The groomed multivalued trivialization τ is defined as follows: Let $\mathfrak{S}_{\pm,i}$ be the sector of D_{ρ_0} given by $\pi_{D_{\rho_0}}(\mathcal{E}_{\pm,i})$. If (ii) and its analogs occur in the proof of Lemma 3.4.4, then we add the thin counterclockwise sectors $\mathfrak{S}(\overline{a}_{k_2,l_2},\overline{h}(\overline{a}_{k_2,l_2}))$, etc. to the set $\{\mathfrak{S}_{+,i}\}$. The sets $\{\mathfrak{S}_{+,i}\}$ and $\{\mathfrak{S}_{-,i}\}$ correspond to the data $\overrightarrow{\mathcal{D}}_-$ and $\overrightarrow{\mathcal{D}}_+$ at the negative and positive ends of $\overline{v}'_{1,1}$[10] and we let τ be the induced groomed multivalued trivialization.

---

[10] Note that the plus and minus signs are switched. This is due to the fact that, for example, $\{\mathfrak{S}_{-,i}\}$ gives the data for the negative ends of $\overline{v}_{1,2}$ that limit to z_∞, which agrees with the data for the positive ends of $\overline{v}_{1,1}$ that limit to z_∞.

By a calculation similar to that of Lemma I.5.7.12, $c_1((\overline{v}'_{1,1})^*T\overline{S},\tau)=1$ and the ends of $\overline{v}'_{1,1}$ contribute the following to $\mu_\tau(\overline{v}'_{1,1})$:

- 0 if $\mathfrak{S}_{-,i}$ is a small sector;
- -1 if $\mathfrak{S}_{-,i}$ is a large sector;
- -1 if $\mathfrak{S}_{+,i}$ is a small sector; and
- -2 if $\mathfrak{S}_{+,i}$ is a large sector.

Hence $\mu_\tau(\overline{v}'_{1,1})=-p_{1,1}-1$. Since $\chi(\dot{F}_{1,1})=p_{1,1}-\mathfrak{b}(\overline{v}'_{1,1})$, we obtain:

$$\begin{aligned}\operatorname{ind}(\overline{v}'_{1,1})&=(\mathfrak{b}(\overline{v}'_{1,1})-p_{1,1})+p_{1,1}+(-p_{1,1}-1)+2\\&=\mathfrak{b}(\overline{v}'_{1,1})-(p_{1,1}-1).\end{aligned}\tag{3.4.12}$$

Next we claim that $\partial F_{1,1}$ is connected. Indeed, if $\partial F_{1,1}$ is disconnected, then the method of Lemma 3.4.4 implies that the union of all the $\mathfrak{S}_{\pm,i}$ covers D_{ρ_0} more than once; this contradicts Equation (3.4.1). The claim in turn implies that $\mathfrak{b}(\overline{v}'_{1,1})\geq p_{1,1}-1$, since otherwise $F_{1,1}$ is disconnected and $\partial F_{1,1}$ will have more than one component. Hence $\operatorname{ind}(\overline{v}'_{1,1})\geq 0$; moreover, if $\operatorname{ind}(\overline{v}'_{1,1})>0$, then $\operatorname{ind}(\overline{v}'_{1,1})\geq 2$.

We claim that $\operatorname{ind}(\overline{v}'_{1,1})\geq 2$ is not possible. Indeed, we add up the Fredholm indices of all the remaining components as in the proof of Lemma 3.4.21:

(a) $\operatorname{ind}(\overline{v}^\sharp_{1,2})\geq 1$ and $\operatorname{ind}(\overline{v}'_+)=-p_+$;
(b) $\sum_{\widetilde{v}}\operatorname{ind}(\widetilde{v})\geq p_+$, where the summation is over all components $\widetilde{v}$ of $\overline{u}_\infty$ that are hanging from the negative end of $\overline{v}'_+$;
(c) a large sector of D^2 contributes an additional $+1$ to the Fredholm index; and
(d) all the other components of $\overline{u}_\infty$ have nonnegative Fredholm index.

(a), (b), (d) are clear, and (c) was explained in Lemma 3.4.21. The total of (a)–(d) is ≥ 2, which is an index excess of $+2$, and the claim follows. The claim then implies that $\operatorname{ind}(\overline{v}'_{1,1})=0$, $\mathfrak{b}(\overline{v}'_{1,1})=p_{1,1}-1$, and $F_{1,1}$ is a disk.

Finally, since F is diffeomorphic to a surface obtained by gluing $\dot{F}_{1,1}$ and $p_{1,1}$ surfaces which are diffeomorphic to B_τ or B_+, it must be connected and planar, and its compactification is a closed disk. □

Lemma 3.4.24. *Suppose $m\gg 0$. If $\overline{v}'_*\cup\overline{v}^\sharp_*\neq\varnothing$ for some level $\overline{v}_*$, then Cases (1), (2$_0$), (3)–(6$_i$) of Lemma 3.4.21 are not possible. This leaves us with Case (2$_1$).*

Proof. Arguing by contradiction, suppose there exist sequences $m_i\to\infty$, $\varepsilon_i,\delta_i\to 0$, and $\overline{u}_{ij}\to\overline{u}_{i\infty}$, where $\overline{u}_{ij}\in\mathcal{M}^{(m_i)}$, $\overline{u}_{i\infty}\in\partial_{\{+\infty\}}\mathcal{M}^{(m_i)}$, $\mathcal{M}^{(m_i)}$ is $\mathcal{M}$ with respect to the family $\{\overline{J}^{\Diamond}_\tau(\varepsilon_i,\delta_i,\mathfrak{p}(\tau);m_i)\}$, and $\overline{u}_{i\infty}$ falls into one of Cases (1), (2$_0$), (3)–(6$_i$). We consider a diagonal subsequence $\overline{u}_i:=\overline{u}_{ij(i)}$.

Cases (1) and (2$_0$) are eliminated by an argument similar to that of Case (2) of Theorem I.7.11.1 and Cases (3)–(5) are eliminated by an argument similar to that of Cases (3)–(6) of Theorem I.7.11.1.

Case (6$_i$). Suppose for simplicity that $b=1$, including connectors.

As in the proof of Theorem I.7.11.1, for $i \gg 0$, we consider a truncation $\widetilde{u}_i : \Sigma_i \to \overline{W}_{\tau_i}$ of $\overline{u}_i$ (i.e., a restriction of $\overline{u}_i$ to a neighborhood of $\sigma_\infty^{\tau_i}$) such that there exist real numbers

$$R_{0,i} < R_{1,i} < R_{2,i} < R_{3,i}, \quad R_{0,i} \ll -l(\tau_i), \quad R_{3,i} \gg l(\tau_i)$$

and a map

$$\pi_i : \Sigma_i \to B_{\tau_i} \cap \{R_{0,i} \leq s \leq R_{3,i}\},$$

such that the restriction of π_i to $\pi_i^{-1}(\{R_{j,i} \leq s \leq R_{j+1,i}\})$, $j = 0, 1, 2$, is a degree $p_{0,j}$ branched cover. Here $p_{0,0} \leq p_{0,1} \leq p_{0,2}$. We project $\widetilde{u}_i$ to $D_{\rho_0} \subset \overline{S}$ for $\rho_0 > 0$ small using balanced coordinates and then apply the ansatz from Equation (I.7.9.1) to obtain $w_i : \Sigma_i \to \mathbb{C}$.

Applying the method of Section I.7.9, we rescale w_i by a positive real constant and take the limit $m_i \to \infty$ to obtain a 3-level holomorphic building

$$w_\infty = w_- \cup w_{0,1} \cup w_+.$$

We write $w_* : \Sigma_* \to \mathbb{CP}^1$ for the components of the building w_∞ and $\pi_* : \Sigma_* \to cl(B_*)$ for the corresponding branched covers, where $* = +, -$, or $(0,1)$, and $B_{0,1} = \mathbb{R} \times S^1$. We may also use subscripts $(0,0)$ and $(0,2)$ to mean $-$ and $+$. Note that $\deg \pi_{0,j} = p_{0,j}$. By Lemma 3.4.23, the surface Σ_∞, obtained by gluing the Σ_*, is connected and planar. The next few paragraphs are devoted to the description of w_* and π_*.

Suppose for simplicity that Σ_-, $\Sigma_{0,1}$, and Σ_+ are connected. Starting from the bottom, w_- and π_- satisfy the following:

(i$_-$) $w_-(\partial\Sigma_-) \subset \{\phi = 0, \rho > 0\}$;
(ii$_-$) $\pi_-^{-1}(+\infty)$ is a single point and $w_-(\pi_-^{-1}(+\infty)) = \infty$;
(iii$_-$) $w_-(z_0) = 0$ for one of the points $z_0 \in \pi_-^{-1}(\overline{\mathfrak{m}}^b(\infty))$; and
(iv$_-$) $w_-|_{int(\Sigma_-)}$ is a biholomorphism onto its image.

(i$_-$) and (iii$_-$) are clear, (ii$_-$) follows from the fact that Σ_∞ (and hence Σ_-) is connected and planar, and (iv$_-$) is a consequence of Equation (3.4.1). Let us write $f_* = \pi_* \circ w_*^{-1}$, where defined. Then f_- maps the asymptotic marker $\dot{\mathcal{R}}_\pi(\infty)$ for $\infty \in \mathbb{CP}^1$, corresponding to the radial ray $\mathcal{R}_\pi$, to the asymptotic marker $\dot{\mathcal{L}}_{3/2}(+\infty)$ for $+\infty \in cl(B_-)$, corresponding to the half-line $\mathcal{L}_{3/2}$, by the Involution Lemma I.7.10.3.

We then move up to $w_{0,1}$, which satisfies the following:

(i$_{0,1}$) $w_{0,1}(\pi_{0,1}^{-1}(-\infty)) = \{d_1 = 0, d_2, \dots, d_k\}$, where $d_i \in \mathbb{R}^{\geq 0}$ and $d_i < d_{i+1}$;
(ii$_{0,1}$) $\pi_{0,1}^{-1}(+\infty)$ is a single point and $w_{0,1}(\pi_{0,1}^{-1}(+\infty)) = \infty$;
(iii$_{0,1}$) $w_{0,1}$ is a biholomorphism; and
(iv$_{0,1}$) $f_{0,1} : \mathbb{CP}^1 \to cl(B_{0,1})$ maps $\dot{\mathcal{R}}_\pi(0)$ to $\dot{\mathcal{L}}_{3/2}(-\infty)$.

The placement of the points $d_2, \dots, d_k$ in (i$_{0,1}$) follows from observing that $\overline{v}_-^\sharp$ consists of $I = 1$ components from δ_0 to x_i or x_i'. In particular, the asymptotic eigenfunctions of $\overline{v}_-^\sharp$ at the positive end are close to constant functions with values on $\mathbb{R}^+ \subset \mathbb{C}$. (ii$_{0,1}$) follows from the fact that Σ_∞ is connected and planar and (iii$_{0,1}$) follows from Equation (3.4.1). (iv$_{0,1}$) is a consequence of the fact that

f_- maps $\dot{\mathcal{R}}_\pi(\infty)$ to $\dot{\mathcal{L}}_{3/2}(+\infty)$. Then $f_{0,1}$ maps $\dot{\mathcal{R}}_\pi(\infty)$ to $\dot{\mathcal{L}}_{3/2}(+\infty)$ by the Involution Lemma I.7.10.4.

We now describe the construction of w_+ in some detail. By translating

$$\pi_i^{-1}(B_{\tau_i} \cap \{R_{2,i} \leq s \leq R_{3,i}\})$$

down by T_i, we obtain the branched cover

$$\pi_i^+ : \Sigma_i^+ \to B_+ \cap \{R_{2,i} - T_i \leq s \leq R_{3,i} - T_i\}$$

and the holomorphic map $w_i^+ : \Sigma_i^+ \to \mathbb{C}$. We may assume that $R_{2,i} - T_i \to -\infty$ and $R_{3,i} - T_i \to +\infty$ have been chosen so that there is no sequence of branch points of π_i^+ that limits to $s = \pm\infty$ as $i \to \infty$. Indeed, the branch points that "escape to $s = \pm\infty$" properly belong to a different level. Then w_+ is the limit of w_i^+, after suitably rescaling by positive real constants.

Let $\overrightarrow{\mathcal{D}} = \{(i'_k, j'_k) \to (i_k, j_k)\}_{k=1}^p$ be the data at the positive end of w_+. For $i \gg 0$, the component of $w_i^+|_{(\pi_i^+)^{-1}(\{s=R_{3,i}-T_i\})}$ corresponding to $(i'_k, j'_k) \to (i_k, j_k)$ is arbitrarily close to a normalized asymptotic eigenfunction ϕ_k from $\overline{h}(\overline{a}_{i'_k,j'_k})$ to $\overline{a}_{i_k,j_k}$, after multiplying by some positive real constant. (Here ϕ_k is an eigenfunction of an asymptotic operator and we are not making any *a priori* assumptions on the corresponding eigenvalues.) Now we claim that some ϕ_{k_0} must sweep out a large sector of D^2, since otherwise $\mathrm{Im}(w_i^+)$ cannot pass through the origin, contradicting the "continuity" to $w_{0,1}$. In the limit $i \to \infty$, ϕ_{k_0} will sweep out an angle of 2π.

The top level w_+ satisfies the following:

(i$_+$) $w_+(\pi_+^{-1}(-\infty)) = \{d_1^+ = 0, d_2^+, \dots, d_{k_+}^+\}$, where $d_i^+ \in \mathbb{R}^{\geq 0}$ and $d_i^+ < d_{i+1}^+$;
(ii$_+$) $w_+(\partial\Sigma_+) \subset \{\phi = 0, 0 < \rho \leq \infty\}$;
(iii$_+$) w_+ maps some point of $\pi_+^{-1}(+\infty)$ to ∞;
(iv$_+$) f_+ maps $\dot{\mathcal{R}}_\pi(0)$ to $\dot{\mathcal{L}}_{3/2}(-\infty)$; and
(v$_+$) w_+ is a biholomorphism onto its image.

The placement of the points $d_2^+, \dots, d_{k_+}^+$ in (i$_+$) follows from observing that $\overline{v}_{0,1}^\sharp$ consists of $I = 1$ components from δ_0 to h and that $\mathcal{R}_{\phi_h} \to \mathcal{R}_0$ as $m \to \infty$ by our choice of h from Section 3.1.4. (ii$_+$) and (iii$_+$) are immediate consequences of the construction of w_+ from the previous paragraphs. (iv$_+$) is a consequence of the fact that $f_{0,1}$ maps $\dot{\mathcal{R}}_\pi(\infty)$ to $\dot{\mathcal{L}}_{3/2}(+\infty)$. We now apply the Involution Lemma I.7.10.5 to conclude that $f_+(\infty) = \mathcal{L}_{3/2} \cap \partial B_+$. Note that the Lemma I.7.10.5 applies because the compactification of Σ_+ is a closed disk by Lemma 3.4.23. This contradicts (ii$_+$). We have eliminated Case (6$_i$). $\square$

We are now in a position to prove Lemma 3.3.2.

Proof of Lemma 3.3.2. Suppose $\overline{u}_\infty \in \partial_{\{+\infty\}}\mathcal{M}$. By Lemma 3.4.17, if $\overline{v}'_* \cup \overline{v}_*^\sharp = \varnothing$ for all levels $\overline{v}_*$ of $\overline{u}_\infty$, then $\overline{u}_\infty \in A_1$. If $\overline{v}'_* \cup \overline{v}_*^\sharp \neq \varnothing$ for some level $\overline{v}_*$, then $\overline{u}_\infty$ is as given in Lemma 3.4.21. Now, by Lemma 3.4.24, the only possibility left is Case (2$_1$), which implies that $\overline{u}_\infty \in A_2$. $\square$

3.5. **Degeneration at $-\infty$.** In this subsection we study the limit of holomorphic maps to $\overline{W}_\tau$ as $\tau \to -\infty$, i.e., when $\overline{W}_\tau$ degenerates into $\overline{W}_{-\infty,1} \cup \overline{W}_{-\infty,2}$. This will prove Lemma 3.3.5.

We assume that $m \gg 0$; $\varepsilon, \delta > 0$ are sufficiently small; and $\{\overline{J}_\tau\} \in \overline{\mathcal{I}}^{reg}$ and $\{\overline{J}^\Diamond_\tau(\varepsilon, \delta, \mathfrak{p}(\tau))\}$ satisfy Lemma 3.2.17. Fix $\mathbf{y} \in \mathcal{S}_{\mathbf{a},h(\mathbf{a})}$, $\mathbf{y}' \in \mathcal{S}_{\mathbf{b},h(\mathbf{b})}$ and let

$$\mathcal{M} = \mathcal{M}^{I=2,n^*=m}_{\{\overline{J}^\Diamond_\tau(\varepsilon,\delta,\mathfrak{p}(\tau))\}}(\mathbf{y}, \mathbf{y}'; \overline{\mathfrak{m}}), \quad \mathcal{M}_\tau = \mathcal{M}^{I=2,n^*=m}_{\overline{J}^\Diamond_\tau(\varepsilon,\delta,\mathfrak{p}(\tau))}(\mathbf{y}, \mathbf{y}'; \overline{\mathfrak{m}}(\tau)).$$

We will analyze $\partial_{\{-\infty\}}\mathcal{M}$.

Let $\overline{u}_i$, $i \in \mathbb{N}$, be a sequence of curves in $\mathcal{M}$ such that $\overline{u}_i \in \mathcal{M}_{\tau_i}$ and $\lim_{i\to\infty} \tau_i = -\infty$, and let

$$\begin{aligned} \overline{u}_\infty = \overline{v}_2 \cup (\overline{v}_{L,1} \cup \dots \cup \overline{v}_{L,a}) \cup \overline{v}_1 \cup (\overline{v}_{R,1} \cup \dots \cup \overline{v}_{R,b}) \\ \cup (\overline{v}_{B,1} \cup \dots \cup \overline{v}_{B,c}) \cup (\overline{v}_{T,1} \cup \dots \cup \overline{v}_{T,d}), \end{aligned}$$

be the limit holomorphic building, where $\overline{v}_j$ maps to $\overline{W}_{-\infty,j}$, $j = 1, 2$; $\overline{v}_{L,j}$ and $\overline{v}_{R,j}$ map to $[-2, 2] \times \mathbb{R} \times \overline{S}$; $\overline{v}_{B,j}$ and $\overline{v}_{T,j}$ map to $\mathbb{R} \times [0, 1] \times \overline{S}$. The levels in the first row are arranged in cyclic order from left to right, the levels in the second row are arranged in order from bottom to top, and $\overline{v}_1$ is between $\overline{v}_{B,c}$ and $\overline{v}_{T,1}$. For notational convenience we refer to $\overline{v}_1$ as $\overline{v}_{L,a+1}$, $\overline{v}_{R,0}$, $\overline{v}_{B,c+1}$, or $\overline{v}_{T,0}$, and $\overline{v}_2$ as $\overline{v}_{L,0}$ or $\overline{v}_{R,b+1}$. As usual, we write $p_* = \deg \overline{v}'_*$.

Terminology. Thin counterclockwise sectors in $\mathbb{C}$ from $\overline{b}_{i,j}$ to $\overline{a}_{i,j}$ and from $\overline{h}(\overline{b}_{i,j})$ to $\overline{h}(\overline{a}_{i,j})$ will also be referred to as *thin sectors*.

Outline of proof of Lemma 3.3.5. The initial steps of the proof are similar to those of Section I.7.5–I.7.12 and Section 3.4. However, the authors were unable to prove that $I(\overline{v}_{L,j})$ and $I(\overline{v}_{R,j})$ were nonnegative when $\overline{v}'_1 \neq \varnothing$, since we could not sufficiently control the groomings in order to apply the ECH index inequality (Lemma I.5.7.18). Note that Lemma 3.5.2(4) specifically excludes the case $\overline{v}'_1 \neq \varnothing$. Sections 3.5.3–3.5.6 are intended as a substitute, and involve ideas from tropical geometry (see for example Parker [Pa]) since we are stretching simultaneously in two directions.

3.5.1. *Continuation argument.* We discuss the continuation argument in the current case; this is similar to but more complicated than those of Sections 3.4.1 and 3.4.3. Suppose that $\overline{v}'_* \cup \overline{v}^\sharp_* \neq \varnothing$ for some level $\overline{v}_*$ of $\overline{u}_\infty$. For simplicity we assume that there are no boundary punctures; we leave it to reader to make the appropriate modifications when there are boundary punctures.

Case 1. Suppose that $\overline{v}'_{T,j} \cup \overline{v}^\sharp_{T,j} \neq \varnothing$ for some $j > 0$. We start at a nontrivial negative end $\mathcal{E}_1$ of some $\overline{v}_{T,j_1}$, $j_1 > 0$, limiting to z_∞. We consider the continuation

$$g^1_{j_1-1,1}, \dots, g^1_{1,1}, g_{-a-1,1}, \dots, g_{b+1,1}, g^0_{1,1}, \dots, g^0_{d,1}$$

of the $t = 1$ boundary of $\mathcal{E}_1$ in the direction of $\partial_+ B_\tau$. Here Definition 3.4.3 needs to be adapted in the obvious way to the breaking of a component of $\overline{u}_i|_{\partial \dot{G}_i}$ as $\tau_i \to -\infty$, where $\dot{G}_i$ is the domain of $\overline{u}_i$. The components $g_{-a-1,1}$ and $g_{b+1,1}$

correspond to $\overline{v}_1$ and the components $g_{j,1}$, $j=-a,\dots,b$, correspond to $\overline{v}_{L,j}$, $j=a,a-1,\dots,0$, and $\overline{v}_{R,j}$, $j=b,b-1,\dots,1$, in that order. There are three possibilities:

(i) there is some $g^1_{j,1}$, $0\leq j\leq j_1-1$, which is nontrivial;
(ii) all the $g^1_{j,1}$ are trivial but some $g_{j,1}$ is nontrivial;
(iii) all the $g^1_{j,1}$ and $g_{j,1}$ are trivial.

Cases (i) and (iii) have already been treated in the proof of Lemma 3.4.4. If we are in Case (ii), then the nontrivial component $g_{j,1}$ contains the $s=2$ boundary of a right end $\mathcal{E}_2$ that limits to z_∞. We then consider the continuation of the $s=-2$ boundary of $\mathcal{E}_2$ in the direction of $\partial_- B_\tau$. The details of the continuation are left to the reader, but in the end the sectors $\pi_{D_{\rho_0}}(\mathcal{E}_i)$ will sweep out a neighborhood of z_∞ with the exception of thin sectors.

Case 2. Suppose that $\overline{v}'_{T,j}\cup\overline{v}^\sharp_{T,j}=\varnothing$ for all $j>0$. Then $\overline{v}'_1=\varnothing$. Consider the horizontal levels:

$$\overline{v}_1=\overline{v}_{R,0},\dots,\overline{v}_{R,b},\overline{v}_2,\overline{v}_{L,1}\dots,\overline{v}_{L,a+1}=\overline{v}_1. \tag{3.5.1}$$

Suppose that $\overline{v}'_*\cup\overline{v}^\sharp_*\neq\varnothing$ for some horizontal level $\overline{v}_*$. Let $\overline{v}_*$ be the leftmost level in Equation (3.5.1) such that $\overline{v}^\sharp_*$ has a right end $\mathcal{E}_1$ at z_∞. The sector $\pi_{D_{\rho_0}}(\mathcal{E}_1)$ is not a thin sector and we apply the usual continuation argument to the levels of Equations (3.5.1).

Case 3. The case where $\overline{v}'_{T,j}\cup\overline{v}^\sharp_{T,j}=\varnothing$ for all $j>0$, $\overline{v}'_*\cup\overline{v}^\sharp_*=\varnothing$ for all horizontal levels $\overline{v}_*$, and $\overline{v}'_{B,j}\cup\overline{v}^\sharp_{B,j}\neq\varnothing$ for some $j\leq c$ is treated similarly.

Let $\mathcal{Z}=(\mathfrak{z}_1\to\cdots\to\mathfrak{z}_k\to\mathfrak{z}_1)$ be the cycle constructed using the above continuation argument. Boundary punctures of type (P$_1$), (P$_2$), and (P$_3$) are defined in the same way.

3.5.2. *Some restrictions on $\overline{u}_\infty$.* The following are analogs of Lemmas 3.4.12, 3.4.16 and 3.4.18.

Lemma 3.5.1. *Suppose that $\overline{v}'_*\cup\overline{v}^\sharp_*\neq\varnothing$ for some level $\overline{v}_*$ of $\overline{u}_\infty$. If $\mathcal{E}_i$, $i=1,\dots,q$, are the ends of all the $\overline{v}^\sharp_*$ that converge to z_∞ and $\mathcal{E}'_i$, $i=1,\dots,r$ are the neighborhoods of the punctures of type (P$_3$), then*

$$\sum_{i=1}^q n^*(\mathcal{E}_i)+\sum_{i=1}^r n^*(\mathcal{E}'_i)\geq m-2g.$$

Lemma 3.5.2. *If fiber components are removed from $\overline{u}_\infty$ and the only boundary punctures are of type (P$_3$), then the following hold:*

1. *the ECH index of each $\overline{v}''_*$ is nonnegative;*
2. *the only components of $\overline{u}_\infty$ which have negative ECH index are those of $\overline{v}'_1$, i.e., the branched covers of $\sigma_\infty^{-\infty,1}$;*
3. *the ECH index of each level $\overline{v}_{T,j},\overline{v}_{B,j}\neq\overline{v}_1$ is nonnegative;*
4. *if $\overline{v}'_1=\varnothing$, then the ECH index of each $\overline{v}_{L,j},\overline{v}_{R,j}\neq\overline{v}_1,\overline{v}_2$ is nonnegative;*

(5) *there is an additional contribution of* $\mathfrak{bp}_*$ *towards* I*, where* $\mathfrak{bp}_*$ *is the number of boundary punctures of type (P*$_3$*) on* $\overline{v}_*$.

Proof. (1) and (2) are consequences of the index inequality. (3) and (4) follow from the proof of Lemma 3.4.9. In the proof of (4), the vertical levels $\overline{v}_{1,a},\dots,\overline{v}_{1,0}$ and the arcs $\overline{\mathbf{a}}$, $\overline{h}(\overline{\mathbf{a}})$ are replaced by the horizontal levels given by Equation (3.5.1) and the arcs $\overline{\mathbf{b}}$, $\overline{\mathbf{a}}$. (5) is argued in the same way as Lemma 3.4.16. $\square$

Lemma 3.5.3. *If* $\overline{u}_\infty\in\partial_{\{-\infty\}}\mathcal{M}$ *and* $\overline{v}'_*\cup\overline{v}^\sharp_*=\varnothing$ *for all levels* $\overline{v}_*$ *of* $\overline{u}_\infty$*, then* $\overline{u}_\infty$ *satisfies the following:* $a=b=c=d=0$*;* $I(\overline{v}_1)=\operatorname{ind}(\overline{v}_1)=0$ *and* $I(\overline{v}_2)=\operatorname{ind}(\overline{v}_2)=2$*; and* $\overline{v}_1$ *is a* $W_{-\infty,1}$*-curve and* $\overline{v}_2$ *is a* $\overline{W}_{-\infty,2}$*-curve.*

Proof. Similar to that of Lemma 3.4.17 and is omitted. $\square$

The following is the analog of Lemma 3.4.18.

Lemma 3.5.4. *If* $\overline{v}'_*\cup\overline{v}^\sharp_*\neq\varnothing$ *for some level* $\overline{v}_*$ *of* $\overline{u}_\infty$*, then:*

1. $p_2=\deg(\overline{v}'_2)>0$*;*
2. *there is no boundary puncture of type (P*$_1$*) or (P*$_2$*), i.e., the only boundary punctures are of type (P*$_3$*);*
3. $\overline{u}_\infty$ *has no fiber components and no components* $\overline{v}''_*$ *that intersect the interior of a section at infinity;*
4. *each of* $\overline{v}_{L,j}$*,* $j=1,\dots,a$*, and* $\overline{v}_{R,j}$*,* $j=1,\dots,b+1$*, consists of thin strips and trivial strips; in particular,* $\overline{v}_{L,j}$*,* $j=1,\dots,a$*, and* $\overline{v}_{R,j}$*,* $j=1,\dots,b+1$ *have no punctures that are removable at* z_∞*.*

Proof. (1), (2), (3) We have the following contributions towards n^*: (1) the restriction of $\overline{v}''_2$ to a neighborhood of $\overline{\mathfrak{m}}(-\infty)$ contributes m if $\overline{v}'_2=\varnothing$; (2) boundary punctures of type (P$_1$) or (P$_2$) contribute at least m in total; and (3) a fiber component or a component of $\overline{v}''_*$ that intersects a section at infinity contributes at least m. All cases contradict Lemma 3.5.1.

(4) Arguing by contradiction, suppose that some level $\overline{v}_{R,j_0}$, $1\leq j_0\leq b$, has a component which is not a thin strip or a trivial strip. (The case of $\overline{v}_{L,j_0}$, $0\leq j_0\leq a$, only differs in notation.) Here $\overline{v}_{R,j_0}$ may have a boundary puncture of type (P$_3$).

We claim that the following hold:

(a) $n^*(\overline{v}_{R,j_0})=m$ and $n^{*,alt}(\overline{v}_{R,j_0})\geq m-2g$;
(b) $\overline{v}^\sharp_{R,j_0}\neq\varnothing$ and some end or boundary puncture of type (P$_3$) of $\overline{v}^\sharp_{R,j_0}$ corresponds to a large sector;
(c) $\overline{v}'_1=\varnothing$ and $\overline{v}^\sharp_{T,j}=\varnothing$ for all $j>0$;
(d) no left end of $\overline{v}_{R,j_0}$ limits to a multiple of z_∞; in particular $\overline{v}'_{R,j_0}=\varnothing$;
(e) there are no boundary punctures of type (P$_3$);
(f) the ECH index of each $\overline{v}_{L,j}$, $1\leq j\leq a$, and $\overline{v}_{R,j}$, $1\leq j\leq b+1$ is nonnegative and $I(\overline{v}_{R,j_0})\geq 2$.

Recall the definition of $n^{*,alt}$ from Section 3.2.3.

We first prove (a). Consider the projection

$$\pi_{\overline{S}}:[-2,2]\times\mathbb{R}\times\overline{S}\to\overline{S}.$$

Since we are only dealing with compactness issues, we may assume without loss of generality that $\overline{J}^{\diamondsuit}_{-\infty,2}(\varepsilon,\delta,\mathfrak{p}(-\infty))$ is a product complex structure and the projection $\pi_{\overline{S}}$ is holomorphic. Since $\overline{v}_{R,j_0}$ has a component which is not a thin strip or a trivial strip, $\operatorname{Im}(\pi_{\overline{S}}\circ\overline{v}_{R,j_0})$ must contain the complement of all the thin strips between the $\overline{b}_i$ and the $\overline{a}_i$. This proves (a).

We now prove (b)–(f). (b) follows from (a) by excluding some possibilities using (2), (3). If (c) does not hold, then there is some negative end $\mathcal{E}$ of $\cup_{j=1}^{a}\overline{v}^{\sharp}_{T,j}$ that limits to z_∞ and $n^*(\mathcal{E})\geq k_0-1\gg 2g$. This is a contradiction of (a). If (d) does not hold, i.e., a left end of $\overline{v}_{R,j_0}$ limits to a multiple of z_∞, then in view of (c) there is some right end $\mathcal{E}$ of $\overline{v}^{\sharp}_{*}$ to the left of $\overline{v}_{R,j_0}$ that limits to z_∞ and satisfies $n^*(\mathcal{E})\geq k_0-1\gg 2g$. This is a contradiction of (a). (e) is a consequence of (d). Finally we prove (f). By Lemma 3.5.2(4) and (c), the ECH indices of $\overline{v}_{L,j}$, $j=1,\dots,a$, and $\overline{v}_{R,j}$, $j=1,\dots,b+1$, are nonnegative. Since $\overline{v}^{\sharp}_{R,j_0}$ has a large sector, its ECH index is increased by one. Hence $I(\overline{v}_{R,j_0})\geq 2$.

By (d), the right end of $\overline{v}_{R,j_0}$ limits to a multiple of z_∞. In view of (a), each component of $\overline{v}^{\sharp}_{*}\neq\overline{v}_1$ to the right of $\overline{v}_{R,j_0}$ must be a thin strip. On the other hand, since the ECH index of each thin strip is 1, Equation (3.4.2), Lemma 3.5.2, and (f) together imply that no component of $\overline{v}_{*}\neq\overline{v}_1$ to the right of $\overline{v}_{R,j_0}$ can be a thin strip. Hence $\overline{v}^{\sharp}_{1}$ has a left end that limits to z_∞. Now, by (a), the projection to $\overline{S}$ of the union of all the ends of $\overline{v}^{\sharp}_{1}$ limiting to z_∞ is a union of thin wedges of type $\mathfrak{S}(\overline{b}_{i,j},\overline{a}_{i,j})$. Such a curve $\overline{v}^{\sharp}_{1}$, however, does not exist (cf. Figure 6).

Since we have a contradiction, (4) follows. $\square$

3.5.3. *Truncations.* In the rest of this section (until the proof of Lemma 3.5.15), we consider the case where $\overline{v}'_{*}\cup\overline{v}^{\sharp}_{*}\neq\varnothing$ for some $\overline{v}_{*}$. We have $\overline{v}'_{2}\neq\varnothing$ in view of Lemma 3.5.4(1).

In the next few subsections we consider the case $\overline{v}'_1=\varnothing$. By Lemma 3.5.4(4), $\overline{u}_\infty$ has no boundary punctures that are removable at z_∞.

Definition 3.5.5. Let X be set and $\varepsilon>0$ be a positive real number. Then two functions $f,g:X\to\mathbb{C}^\times$ are *ε-approximate* if

$$|f(x)-g(x)|<\varepsilon\cdot|g(x)|\quad\text{and}\quad|f(x)-g(x)|<\varepsilon\cdot|f(x)|$$

for all $x\in X$.

We now define the following sequence of truncations, analogous to those which appear in the proof of Theorem I.7.11.1,

Definition 3.5.6. For $i\gg 0$ and $\varepsilon>0$ small, a restriction of $\overline{u}_i$ to a neighborhood of $\sigma^{\tau_i}_{\infty}$ (where τ_i is chosen appropriately) is called a *truncation* $\widetilde{u}_i:\Sigma_i\to\overline{W}_{\tau_i}$ of $\overline{u}_i$ if the following hold:

(T1) $\widetilde{u}_i(\Sigma_i)$ is contained in a 2ε-neighborhood of $\sigma^{\tau_i}_{\infty}$;
(T2) if $\overline{u}_i(x)$ is contained in an $\frac{\varepsilon}{2}$-neighborhood of $\sigma^{\tau_i}_{\infty}$, then $x\in\Sigma_i$;

(T3) there exist constants $r(\tau_i) - 2 > R_i^{(1)} > \cdots > R_i^{(\iota+1)} > 2$ and a decomposition $\Sigma_i = \Sigma_i^{(1)} \cup \cdots \cup \Sigma_i^{(\iota)}$ such that

$$\Sigma_i^{(j)} = (\pi_{B_{\tau_i}} \circ \widetilde{u}_i)^{-1}([-2,2] \times [R_i^{(j+1)}, R_i^{(j)}]) \cap \Sigma_i$$

and

$$\pi_{B_{\tau_i}} \circ \widetilde{u}_i : \Sigma_i^{(j)} \to [-2,2] \times [R_i^{(j+1)}, R_i^{(j)}]$$

is a branched cover with possibly empty branch locus;

(T4) each component c of $\pi_{D_{\rho_0}} \circ \widetilde{u}_i|_{\partial_v \Sigma_i}$ is ε-approximate to a positive multiple of a normalized eigenfunction of $\overline{v}^{\sharp}_{L,j}$, $j = 0, \dots, a$, or $\overline{v}^{\sharp}_{R,j}$, $j = 0, \dots, b$; moreover, *the positive multiple is independent of* i *and the component* c.

Here

$$\pi_{D_{\rho_0}} : \pi^{-1}_{B_{\tau_i}}([-2,2] \times [R_i^{(\iota+1)}, R_i^{(1)}]) \cap \{\rho \leq \rho_0\} \to D_{\rho_0}$$

is obtained by projecting out the ∂_s- and $\overline{R}_{\tau_i}$-directions and

$$\partial_v \Sigma_i = \partial \Sigma_i - (\pi_{B_{\tau_i}} \circ \widetilde{u}_i)^{-1}(\{s = \pm 2\})$$
$$\partial_h \Sigma_i = \partial \Sigma_i \cap (\pi_{B_{\tau_i}} \circ \widetilde{u}_i)^{-1}(\{s = \pm 2\}).$$

Given a truncation $\widetilde{u}_i : \Sigma_i \to \overline{W}_{\tau_i}$, let $\widetilde{\pi}_i$ be the map obtained by postcomposing

$$\pi_{B_{\tau_i}} \circ \widetilde{u}_i : \Sigma_i \to [-2,2] \times [R_i^{(\iota+1)}, R_i^{(1)}]$$

with a $-\frac{r(\tau_i)+1}{2}$-translation in the t-direction, and let $\widetilde{w}_i = \pi_{D_{\rho_0}} \circ \widetilde{u}_i|_{\Sigma_i}$. Also let

$$R_i^+ = R_i^{(1)} - \frac{r(\tau_i)+1}{2}, \quad R_i^- = R_i^{(\iota+1)} - \frac{r(\tau_i)+1}{2}.$$

Notation. When we want to distinguish the t-coordinates for $B_{-\infty,1}$ and $B_{-\infty,2}$, we write t_i for the t-coordinate for $B_{-\infty,i}$.

Lemma 3.5.7. *If* $i \gg 0$, *then* Σ_i *is a disk with* $\geq 2p_2$ *boundary punctures.*

Proof. This is a consequence of (T4) and the following:

(i) $n^*(\widetilde{u}_i|_{\Sigma_i}) = m$;
(ii) $\widetilde{w}_i(z_0) = 0$ for some $z_0 \in (\pi_{B_{\tau_i}} \circ \widetilde{u}_i)^{-1}(\overline{\mathfrak{m}}^b(\tau_i))$;
(iii) $\widetilde{w}_i$ maps each component of $\widetilde{\pi}_i^{-1}(\{s = 2\})$ to a different $\mathcal{R}_{\phi(\overline{a}_*)}$ and each component of $\widetilde{\pi}_i^{-1}(\{s = -2\})$ to a different component of $\mathcal{R}_{\phi(\overline{b}_*)}$;
(iv) $\widetilde{w}_i|_{int(\Sigma_i)}$ is a biholomorphism onto its image.

Here $\phi(\overline{a}_*)$ is the ϕ-coordinate for $\overline{a}_*$, etc. □

3.5.4. *Large-scale behavior of* Ξ_i. When taking the limits of $\widetilde{\pi}_i$ and $\widetilde{w}_i$, we are simultaneously stretching in two directions t_2 and $\log \rho$. In this subsection we study the large-scale behavior of the map

$$\Xi_i = (t_2 \circ \widetilde{\pi}_i, \log \rho \circ \widetilde{w}_i) : \Sigma_i \to [R_i^-, R_i^+] \times [-\infty, \infty)$$

as $i \to \infty$, where we use coordinates (x', y') on $[R_i^-, R_i^+] \times [-\infty, \infty)$.[11] The goal is to construct the "tropical curve"

$$\overline{\Xi}_i : \Gamma_i \to [-1,1] \times [0, d_i],$$

with $d_i \geq 1$, which approximates Ξ_i when viewed from "far away"; see Figure 3.5.10. Here Γ_i is a finite graph whose topological type is independent of $i \gg 0$. The analysis is of the same type as that of Parker [Pa].

Step 1. We start with the following lemma, which is a consequence of Gromov compactness and which describes the behavior of $\widetilde{w}_i$ and $\widetilde{\pi}_i$ for $i \gg 0$.

Lemma 3.5.8. *Given $\varepsilon > 0$ sufficiently small, there exist constants $i_0 > 0$, $L > 0$, and $\kappa, \kappa' \in \mathbb{Z}^+$ such that for each $i \geq i_0$ there exist:*

- *disjoint compact subsets $K_{i1}, \dots, K_{i\kappa} \subset [-2,2] \times [R_i^-, R_i^+]$ of the form $[-2,2] \times [\tau_{ij} - L, \tau_{ij} + L]$ for suitable τ_{ij}; and*
- *components $C_{i1}, \dots, C_{i\kappa'}$ such that $\Sigma_i - \widetilde{\pi}_i^{-1}(\cup_j K_{ij}) = \sqcup_{j=1}^{\kappa'} C_{ij}$;*

such that:

1. *$\widetilde{\pi}_i$ has no branch points outside $K_{i1}, \dots, K_{i\kappa}$;*
2. *K_{ij} is disjoint from $[-2,2] \times \{R_i^{\pm}\}$;*
3. *for each i, $\cup_{j=1}^{\kappa} K_{ij} \supset [-2,2] \times \{0\}$; and*
4. *$\widetilde{w}_i|_{C_{ij}}$ is ε-approximate to a multiple of $e^{\lambda_{ij}(t-is)}$, where $\lambda_{ij} \in \mathbb{R} - \{0\}$ is $\pm\frac{\pi}{2}$ times the angle of a sector of type $\mathfrak{S}(\overline{b}_{k,l}, \overline{a}_{k',l'})$.*

Observe that there is an upper bound on the number of branch points of $\widetilde{\pi}_i$ which is independent of $i \gg 0$ and $m \gg 0$. This is a consequence of the ECH compactness theorem (cf. Section I.3.4).

Corollary 3.5.9. *Given $\varepsilon > 0$ sufficiently small, there exists $i_0 > 0$ such that, for each $i \geq i_0$, the map Ξ_i satisfies the following:*

1. *$\Xi_i(C_{ij})$ is close to a line segment $\{y' = \lambda_{ij} x' + \beta_{ij} \mid x' \in t_2 \circ \widetilde{\pi}_i(C_{ij})\}$, where β_{ij} is a constant;*
2. *$\Xi_i(\pi_i^{-1}(K_{ij})) \subset t_2(K_{ij}) \times \log \rho(\widetilde{w}_i \circ \widetilde{\pi}_i^{-1}(K_{ij}))$; and*
3. *if $K_{ij_0} \supset [-2,2] \times \{0\}$, then $(0, -\infty) \in \Xi_i(\pi_i^{-1}(K_{ij_0}))$.*
4. *there exists $d'_i \in \mathbb{R}$ such that $y' \circ \Xi_i(\mathfrak{b}) \approx d'_i$ for each component $\mathfrak{b}$ of $\partial_v \Sigma_i$ and $\max_{\mathfrak{b}} y' \circ \Xi_i(\mathfrak{b}) = d'_i$.*

Proof. (1)–(3) immediately follow from Lemma 3.5.8. (4) is a consequence of (T4) in Definition 3.5.6. □

Step 2. We now apply *interval contractions* to construct the tropical curve $\overline{\Xi}_i$.

Definition 3.5.10 (Interval contraction map)**.** Let $I_1, \dots, I_k$ be disjoint closed intervals of $[a,b]$. An *interval contraction map $f : [a,b] \to [c,d]$ which contracts $I_1, \dots, I_k$* is a piecewise linear, nondecreasing, continuous map such that

---

[11] We already have coordinates $(t_2, \log \rho)$, but the expressions are cleaner if we use $(x' = t_2, y' = \log \rho)$.

(1) $f(a) = c$, $f(b) = d$;
(2) f is constant on each I_i; and
(3) f' is constant on $[a, b] - \cup_i I_i$.

We can similarly define an interval contraction map $f : [-\infty, b] \to [c, d]$, provided some I_i is equal to $[-\infty, b']$ for some b'.

We define two interval contraction maps: $p_{i1} : [R_i^-, R_i^+] \to [-1, 1]$ which contracts $t_2(K_{ij})$, $j = 1, \dots, \kappa$, and $p_{i2} : [-\infty, d_i'] \to [0, d_i]$, which contracts $\log \rho(\widetilde{w}_i \circ \widetilde{\pi}_i^{-1}(K_{ij}))$, $j = 1, \dots, \kappa$. Here $d_i \geq 1$ will be chosen later so that Lemma 3.5.11(4) holds. In particular, $p_{i1}(0) = 0$ and $p_{i2}(-\infty) = 0$ by Corollary 3.5.9(2) and (3).

We then define $\Xi_i' := (p_{i1}, p_{i2}) \circ \Xi_i$. By the above definitions of p_{i1} and p_{i2}, Ξ_i' maps K_{ij} to points and components of $\Sigma_i - \widetilde{\pi}_i^{-1}(\cup_j K_{ij})$ to approximate straight line segments.

Lemma 3.5.11. *Given $\varepsilon > 0$ sufficiently small, there exists $i_0 > 0$ such that, for each $i \geq i_0$, the map Ξ_i' is C^0-close to a map which factors through*

$$\overline{\Xi}_i : \Gamma_i \to [-1, 1] \times [0, d_i],$$

where $\Gamma_i = (V_{\Gamma_i}, E_{\Gamma_i})$ is a finite graph, V_{Γ_i} is the set of vertices, and E_{Γ_i} is the set of edges, and the following hold:

(1) *the set V_{Γ_i} admits a decomposition $V_{\Gamma_i,i} \sqcup V_{\Gamma_i,e}$, where $V_{\Gamma_i,i}$ is in one-to-one correspondence with the set of components of $\widetilde{\pi}_i^{-1}(K_{ij})$ and $V_{\Gamma_i,e}$ is in one-to-one correspondence with the set of components of $\partial_v \Sigma_i$;*
(2) *the set E_{Γ_i} is in one-to-one correspondence with the set $\{C_{i1}, \dots, C_{i\kappa'}\}$ and each $e \in E_{\Gamma_i}$ is mapped to a straight line segment;*
(3) *if $\lambda_i : E_{\Gamma_i} \to \mathbb{R}$ maps $e \mapsto \lambda_{ij}$, where e corresponds to C_{ij}, and $\lambda_i' : E_{\Gamma_i} \to \mathbb{R}$ maps e to the slope of $\overline{\Xi}_i(e)$, then λ_i / λ_i' is ε-approximate to a constant function for small $\varepsilon > 0$;*
(4) *$d_i \geq 1$ is chosen so that $\max\limits_{e \in E_{\Gamma_i}} |\lambda_i'(e)| = 1$;*
(5) *if $E_{\Gamma_i}' \subset E_{\Gamma_i}$ is the set of edges such that $|\lambda_i'(e)| < 1 - \delta$ for some small $\delta > 0$ and $e \in E_{\Gamma_i}'$, then $|\lambda_i'(e)| \leq K/m$ for some constant $K > 0$ which is independent of $m \gg 0$ and i.*
(6) *$y \circ \overline{\Xi}_i(p) = d_i$ for all $p \in V_{\Gamma_i,e}$ and $\# V_{\Gamma_i,e}$ is even;*
(7) *$\overline{\Xi}_i(p) = (\pm 1, d_i)$ for all $p \in V_{\Gamma_i,e}'$, where $V_{\Gamma_i,e}' \subset V_{\Gamma_i,e}$ is the subset of vertices which correspond to ends of $\overline{v}_1^\sharp$;*
(8) *each vertex of $V_{\Gamma_i,i}$, has the same number ≥ 1 of edges to its left (i.e., smaller x-coordinate) and to its right (i.e., larger x-coordinate);*
(9) *there is a vertex $q_0 \in V_{\Gamma_i,i}$ such that $\overline{\Xi}_i(q_0) = (0, 0)$.*

Note that $V_{\Gamma_i,i}$ are the interior vertices and $V_{\Gamma_i,e}$ are the exterior vertices.

Proof. (1), (2), the second assertion of (6), (8), and (9) follow from the construction. (4) is a normalization. The first assertion of (6) follows from perturbing $\overline{\Xi}_i$ and using Corollary 3.5.9(4).

(7) follows from Corollary 3.5.9(4), together with the observation that, once $\varepsilon > 0$ is fixed, the distance between $t_2 \circ \widetilde{\pi}_i(\mathfrak{b})$ and $t_2 \circ \widetilde{\pi}_i(\mathfrak{b}')$ is bounded above by a constant which is independent of i, whenever $\mathfrak{b}, \mathfrak{b}'$ are components of $\partial_v \Sigma_i$ corresponding to left (resp. right) ends of $\overline{v}_1^\sharp$. As $i \to \infty$, this distance becomes negligible since $\overline{\Xi}_i$ is the result of rescaling to a bounded size (i.e., there exists $d > 1$ such that $[-1,1] \times [0, d_i] \subset [-1,1] \times [0, d]$ for all i).

(3) follows from Lemma 3.5.8(4), Definition 3.5.10(3), and the observation that $|R_i^+| - |R_i^-|$ is bounded above a constant which is independent of i. (5) follows from (3). $\square$

See Figure 11 for an example. Let (x, y) be the coordinates of $[-1,1] \times [0, d_i]$. The following lemma is immediate from the construction.



FIGURE 11. The top figure represents $[-2,2] \times [R_i^-, R_i^+]$, where × indicates the location of $\overline{\mathfrak{m}}^b(\tau_i)$ after translation and the shaded regions are K_{ij}. The bottom is a schematic diagram for the image of Ξ_i and becomes the image of $\overline{\Xi}_i$ after applying interval contractions. The dots along the line $y' = d'_i$ correspond to the endpoints of $V_{\Gamma_i,e}$ and the shaded regions are $t_2(K_{ij}) \times \log \rho(\widetilde{w}_i \circ \widetilde{\pi}_i^{-1}(K_{ij}))$. The edges with the largest slope (in absolute value) correspond to $|\lambda'_i| \approx 1$; the remaining edges have much smaller slope when $m \gg 0$.

Lemma 3.5.12. *After passing to a subsequence, we may assume that the following do not depend on the choice of* $i \gg 0$*:*

- *the graph* Γ_i*, the function* λ_i*, and the set* E'_{Γ_i}*;*
- *given* $p, q \in V_{\Gamma_i}$*, whether* $x \circ \overline{\Xi}_i(p) \geq x \circ \overline{\Xi}_i(q)$ *and whether* $y \circ \overline{\Xi}_i(p) \geq y \circ \overline{\Xi}_i(q)$.

In view of Lemma 3.5.12, we may write $\Gamma = (V_\Gamma, E_\Gamma)$ for $\Gamma_i = (V_{\Gamma_i}, E_{\Gamma_i})$.

Finally, we orient the edges $p \overset{e}{\to} q \in E_\Gamma$ so that $y \circ \overline{\overline{\Xi}}_i(q) - y \circ \overline{\overline{\Xi}}_i(p) > 0$. Given $p, q \in V_\Gamma$, we write $p \succeq q$ (resp. $p \simeq q$) to mean $y \circ \overline{\overline{\Xi}}_i(p) - y \circ \overline{\overline{\Xi}}_i(q) \geq 0$ (resp. $= 0$) for all $i \gg 0$.

3.5.5. *The case* $\overline{v}'_1 = \varnothing, \overline{v}'_2 \neq \varnothing$. In this subsection we consider the case $\overline{v}'_1 = \varnothing$, $\overline{v}'_2 \neq \varnothing$. See Figure 12(1).



(1) (2) (3)

FIGURE 12. Schematic diagrams for the possible types of degenerations corresponding to Lemmas 3.5.14 and 3.5.15. Here $\circ$ represents z_∞ or $z_\infty^\#$, $\square$ represents a branch point, and $\triangle$ represents an end with a large sector. Double dotted lines indicate multiple covers of σ_∞^*. The labels on the graphs indicate the components and their ECH indices. If there is more than one $\square$ or $\triangle$ in a diagram, then we interpret it as one of the possible locations for $\square$ or $\triangle$.

We start with the following useful lemma:

Lemma 3.5.13 (Comparison Lemma). *Suppose* $m \gg 0$, $i = i(m) \gg 0$, *and* $\varepsilon = \varepsilon(m) > 0$ *is small. Let* $q \overset{e}{\to} q'$ *be an edge of* Γ *such that* $\lambda'_i(e) \simeq 1$ *and let*

$$\delta' = (p_1 \overset{f_1}{\to} \cdots \overset{f_k}{\to} p_{k+1})$$

be an oriented path of Γ *from* p_1 *to* p_{k+1} *such that* $p_1 \preceq q \preceq q' \preceq p_{k+1}$ *and* $0 < \lambda'_i(f_j) \leq K/m$ *for* $j = 1, \dots, k$. *Then*

$$|x \circ \overline{\Xi}_i(q') - x \circ \overline{\Xi}_i(q)| \leq K'/m,$$

where K' *is independent of* m *and* i.

Note that $0 < \lambda'_i(f_j) \leq K/m$ whenever $\lambda'_i(f_j) \not\simeq 1$ by Lemma 3.5.11(5).

Proof. The horizontal variation

$$\sum_{j=1}^{k} |x \circ \overline{\Xi}_i(p_{j+1}) - x \circ \overline{\Xi}_i(p_j)|$$

is bounded above by 2 times the maximal covering degree of $\overline{v}'_{L,j}$, $j = 1, \dots, a$, and $\overline{v}'_{R,j}$, $j = 1, \dots, b+1$, which we denote by $\mathcal{K}$. (Here the 2 comes from the width of the interval $[-1,1]$.) On the other hand, since $p_1 \preceq q \preceq q' \preceq p_{k+1}$, we have:

$$\begin{aligned} |x \circ \overline{\Xi}_i(q') - x \circ \overline{\Xi}_i(q)| &= |y \circ \overline{\Xi}_i(q') - y \circ \overline{\Xi}_i(q)| \\ &\leq \sum_{j=1}^{k} \lambda'_i(f_j) |x \circ \overline{\Xi}_i(p_{j+1}) - x \circ \overline{\Xi}_i(p_j)| \\ &\leq \left(\max_{j=1,\dots,k} \lambda'_i(f_j) \right) (2\mathcal{K}) \leq K'/m, \end{aligned}$$

where $K' = 2\mathcal{K}K$. ☐

Given $(x_0, y_0) \in [-1,1] \times [0, d_i]$, let

$$l_{(x_0,y_0)} = \{x = x_0, 0 \leq y \leq y_0\}$$

be a line segment oriented in the positive y-direction. Next orient Γ so that the oriented edges of $\overline{\Xi}_i(\Gamma)$ are pointing in the positive x-direction; we denote the oriented graph by $\widetilde{\Gamma}$ to distinguish this orientation from the orientation of Γ defined previously.

Consider the *weight function*

$$\mathcal{W}_i : ([-1,1] \times [0, d_i]) - \overline{\Xi}_i(\Gamma) \to \mathbb{Z}^{\geq 0},$$

which is defined as follows:

(1) $\mathcal{W}_i$ is locally constant;

(2) $\mathcal{W}_i(x,y)$ is the signed intersection number $\langle \overline{\Xi}_i(\widetilde{\Gamma}), l_{(x,y)} \rangle$ for generic (x,y), with respect to the orientation (∂_x, ∂_y) for $[-1,1] \times [0, d_i]$.

The function $\mathcal{W}_i$ is well-defined by Lemma 3.5.11(7).

Lemma 3.5.14. *If $m \gg 0$, then there is no $\overline{u}_\infty \in \partial_{\{-\infty\}}\mathcal{M}$ such that $\overline{v}'_1 = \varnothing$ and $\overline{v}'_2 \neq \varnothing$.*

Proof. The proof we give is not the most efficient, but carries over more easily to other situations. We choose $m \gg 0$ so that $0 < K'/m \ll 1$

Let $\delta = (q_0 \overset{e_0}{\to} \cdots \overset{e_{l-1}}{\to} q_l)$ be an oriented path from q_0 to $q_l \in V_{\Gamma,e}$, such that $\lambda'_i(e_j) \approx 1$ for each $i \gg 0$ and each edge e_j, $j = 0, \dots, l-1$. We may assume that $x \circ \overline{\Xi}_i(q_l) = \pm 1$ in view of Lemma 3.5.4(4), since the only end $\mathcal{E}$ of $\overline{v}^\sharp_{L,j}$, $j = 1, \dots, a+1$, or $\overline{v}^\sharp_{R,j}$, $j = 1, \dots, b+1$, whose $\pi_{D_{\rho_0}}$-projection is a sector with angle $> \pi$ is an end of $\overline{v}^\sharp_1 = \overline{v}^\sharp_{L,a+1}$.

We claim that, for each $l_0 = 0, \dots, l-1$, there is an oriented path

$$\delta' = (p_1 \overset{f_1}{\to} \cdots \overset{f_k}{\to} p_{k+1})$$

such that $p_1 \preceq q_{l_0} \preceq q_{l_0+1} \preceq p_{k+1}$ and $0 < \lambda'_i(f_j) < K'/m$ for $j = 1, \dots, k$. Arguing by contradiction, if the claim does not hold, then $\overline{\Xi}_i(\Gamma) \cap \{y = y_0\}$ consists of only one point $(x_0, y_0) \in \overline{\Xi}_i(e_{l_0}) \cap \{y = y_0\}$ for any constant y_0 in the interval $(y \circ \overline{\Xi}_i(q_{l_0}), y \circ \overline{\Xi}_i(q_{l_0+1}))$. This means that there is an integer κ such that $\mathcal{W}_i(x, y_0) = \kappa$ for $-1 \leq x < x_0$ and $\mathcal{W}_i(x, y_0) = \kappa \pm 1$ for $1 \geq x > x_0$. This contradicts $\mathcal{W}_i(-1, y_0) = \mathcal{W}_i(1, y_0) = 0$, which is due to the fact that all the exterior vertices $p \in V_{\Gamma_i,e}$ satisfy $y \circ \overline{\Xi}_i(p) = d_i$ by Lemma 3.5.11(6).

The claim, together with the Comparison Lemma, implies that

$$|x \circ \overline{\Xi}_i(q_{k+1}) - x \circ \overline{\Xi}_i(q_k)| \leq K'/m$$

for each $k = 0, \dots, l-1$. This in turn gives $|x \circ \overline{\Xi}_i(q_l) - x \circ \overline{\Xi}_i(q_0)| \leq lK'/m$, which contradicts Lemma 3.5.11(7). $\square$

3.5.6. *The case $\overline{v}'_1 \neq \varnothing$, $\overline{v}'_2 \neq \varnothing$.* In this subsection we consider the case $\overline{v}'_1 \neq \varnothing$, $\overline{v}'_2 \neq \varnothing$. For simplicity assume that there are no punctures of type (P_3). Some of the possibilities are given by Figure 12(2) and (3).

We outline the necessary modifications in the current case:

(1) We consider the truncation $\widetilde{u}_i : \Sigma_i \to \overline{W}_{\tau_i}$ of $\overline{u}_i$ so that (T1) and (T2) as well as the analogs of (T3) and (T4) of Definition 3.5.6 hold. We then restrict Σ_i to $|s| \leq L$, while keeping the same notation. In other words, we throw away the portion of Σ_i that maps to the positive and negative ends of $\overline{W}_{\tau_i}$.

(2) To the list of compact subsets K_{ij} of Lemma 3.5.8 (viewed as subsets of B_{τ_i} instead of $[-2, 2] \times [R_i^-, R_i^+]$), we add the compact subset K_{ij} of the following type, which we call "type $\overline{v}_1$":

$$K_{ij} = B_{\tau_i} \cap \{|s| \leq L, |t_1 - 1/2| \leq L\}.$$

(3) After suitable translations, interval contractions, and rescalings, we obtain the "tropical curves"

$$\overline{\Xi}_i : \Gamma \to [-1, 1]/(-1 \sim 1) \times [0, d_i],$$

where the equivalence relation $\sim$ comes from contracting K_{ij} of type $\overline{v}_1$ to a point and the graph $\Gamma = (V_\Gamma, E_\Gamma)$ is independent of i.

(4) Lemma 3.5.11(1) is modified as follows: The set V_Γ of vertices admits the decomposition $V_{\Gamma,i} \sqcup V_{\Gamma,e}$, where $V_{\Gamma,i}$ is in one-to-one correspondence with the set of components of $\widetilde{\pi}_i^{-1}(K_{ij})$ and $V_{\Gamma,e}$ is in one-to-one correspondence with the set of left and right ends of $\overline{v}^\sharp_{L,j}$, $j = 1, \dots, a+1$, and $\overline{v}^\sharp_{R,j}$, $j = 1, \dots, b+1$, that limit to z_∞. There is a subset $V_{\Gamma,i'} \subset V_{\Gamma,i}$ consisting of vertices which are not initial points of any oriented edge; $V_{\Gamma,i'}$ is a subset of points obtained by contracting K_{ij} of type $\overline{v}_1$ to a point.

(5) Lemma 3.5.11(2)–(5), (8), (9) still hold, but (6) is modified as follows: $y \circ \overline{\Xi}_i$ attains its maximum d_i at all $p \in V_{\Gamma,e}$ and $\# V_{\Gamma,e}$ is even, but $V_{\Gamma,e}$ may be empty. In that case, $y \circ \overline{\Xi}_i$ attains its maximum at some point of $V_{\Gamma,i'}$.

(6) The weight function $\mathcal{W}_i$ is now a function

$$\mathcal{W}_i : (([-1, 1]/\sim) \times [0, d_i]) - \overline{\Xi}_i(\Gamma) \to \mathbb{Z}^{\geq 0},$$

i.e., it is periodic in the x-direction.

Lemma 3.5.15. *If $m \gg 0$, then there is no $\overline{u}_\infty \in \partial_{\{-\infty\}}\mathcal{M}$ such that $\overline{v}'_1 \neq \varnothing$ and $\overline{v}'_2 \neq \varnothing$.*

Proof. The proof is similar to that of Lemma 3.5.14 and uses the Comparison Lemma.

Suppose that there are no punctures of type (P$_3$). Let $\delta = (q_0 \overset{e_0}{\to} \cdots \overset{e_{l-1}}{\to} q_l)$ be a maximal oriented path which starts from q_0, has $\lambda_i(e) \approx 1$ for each edge, and ends at some q_l with $x \circ \overline{\Xi}_i(q_l) = \pm 1$. Then, for each $l_0 = 0, \dots, l-1$, there is an oriented path

$$\delta' = (p_1 \overset{f_1}{\to} \cdots \overset{f_k}{\to} p_{k+1})$$

such that $p_1 \preceq q_{l_0} \preceq q_{l_0+1} \preceq p_{k+1}$ and $0 < \lambda_i'(f_j) \leq K'/m$ for $j = 1, \dots, k$. Otherwise, using the same notation as that of Lemma 3.5.14, there is a point (x_0, y_0) and an integer κ such that $\mathcal{W}_i(x, y_0) = \kappa$ for $-1 \leq x < x_0$ and $\mathcal{W}_i(x, y_0) = \kappa \pm 1$ for $x_0 \leq x \leq 1$, which is impossible by the x-periodicity.

Suppose there are punctures of type (P$_3$). Then there are no punctures of type (P$_3$) on $\overline{v}_{L,j}$, $j = 1, \dots, a$, and $\overline{v}_{R,j}$, $j = 1, \dots, b+1$, by Lemma 3.5.4(4). Then the maximal oriented path δ from the previous paragraph has $y \circ \overline{\Xi}_i(q_l)$ which is much larger than $y \circ \overline{\Xi}_i$ of the endpoint of another path. This is a contradiction. □

Proof of Lemma 3.3.5. Suppose $\overline{u}_\infty \in \partial_{\{-\infty\}}\mathcal{M}$. If $\overline{v}_*' \cup \overline{v}_*^\sharp = \varnothing$ for all levels $\overline{v}_*$ of $\overline{u}_\infty$, then, by Lemma 3.5.3, $\overline{u}_\infty$ is a 2-level building $\overline{v}_1 \cup \overline{v}_2$, where $\overline{v}_1$ is a $W_{-\infty,1}$-curve with $I = 0$ and $\overline{v}_2$ is a $\overline{W}_{-\infty,2}$-curve with $I = 2$ which passes through $\overline{\mathfrak{m}}(-\infty)$. By Lemma 3.2.10, $\mathbf{y}_2$ and $\mathbf{y}_4$ satisfy the conditions of A_3.

On the other hand, it is not possible that $\overline{v}_*' \cup \overline{v}_*^\sharp \neq \varnothing$ for some level $\overline{v}_*$ by Lemmas 3.5.14 and 3.5.15. □

3.6. **Breaking in the middle.**

In this subsection we study the limit of holomorphic maps to $\overline{W}_\tau$ as $\tau \to T'$ for some $T' \in (-\infty, \infty)$. This will prove Lemma 3.3.8.

We assume that $m \gg 0$; $\varepsilon, \delta > 0$ are sufficiently small; and $\{\overline{J}_\tau\} \in \overline{\mathcal{I}}^{reg}$ and $\{\overline{J}_\tau^\Diamond(\varepsilon, \delta, \mathfrak{p}(\tau))\}$ satisfy Lemma 3.2.17. Fix $\mathbf{y} \in \mathcal{S}_{\mathbf{a},h(\mathbf{a})}$, $\mathbf{y}' \in \mathcal{S}_{\mathbf{b},h(\mathbf{b})}$ and let

$$\mathcal{M} = \mathcal{M}^{I=2,n^*=m}_{\{\overline{J}_\tau^\Diamond(\varepsilon,\delta,\mathfrak{p}(\tau))\}}(\mathbf{y}, \mathbf{y}'; \overline{\mathfrak{m}}), \quad \mathcal{M}_\tau = \mathcal{M}^{I=2,n^*=m}_{\overline{J}_\tau^\Diamond(\varepsilon,\delta,\mathfrak{p}(\tau))}(\mathbf{y}, \mathbf{y}'; \overline{\mathfrak{m}}(\tau)).$$

We will analyze $\partial_{(-\infty,\infty)}\mathcal{M}$.

Let $\overline{u}_i$, $i \in \mathbb{N}$, be a sequence of curves in $\mathcal{M}$ such that $\overline{u}_i \in \mathcal{M}_{\tau_i}$ and $\lim_{i\to\infty} \tau_i = T'$, and such that its limit

$$\overline{u}_\infty = (\overline{v}_{-1,1} \cup \cdots \cup \overline{v}_{-1,c}) \cup \overline{v}_0 \cup (\overline{v}_{1,1} \cup \cdots \cup \overline{v}_{1,a})$$

is a holomorphic building in $\partial_{\{T'\}}\mathcal{M}$, ordered from bottom to top, where $\overline{v}_{-1,j}$, $j = 1, \dots, c$, and $\overline{v}_{1,j}$, $j = 1, \dots, a$, map to $\overline{W}$ and $\overline{v}_0$ maps to $\overline{W}_{T'}$. Sometimes we will refer to $\overline{v}_0$ as $\overline{v}_{-1,c+1}$ or $\overline{v}_{1,0}$.

The following are analogs of Lemmas 3.4.16–3.4.18, stated without proof.

Lemma 3.6.1. *If fiber components are removed from $\overline{u}_\infty$ and the only boundary punctures are of type (P$_3$), then the ECH index of each level $\overline{v}_* \neq \overline{v}_0$ is nonnegative, $I(\overline{v}_{1,j}) \geq I(\overline{v}_{1,j}') + I(\overline{v}_{1,j}'') + \mathfrak{bp}_{1,j}$ for $0 \leq j \leq a$, and the only components of $\overline{u}_\infty$ which have negative ECH index are the following:*

1. *branched covers of $\sigma_\infty^{T'}$; and*

(2) *at most one component* $\widetilde{v}$ *of* $\overline{v}''_0$ *with* $I(\widetilde{v}) = -1$.

Here (2) occurs when $T' \in \mathcal{T}_1$ *and* $\widetilde{v} \in \mathcal{M}^{\dagger,s,irr,\mathrm{ind}=-1}_{\overline{J}^{\Diamond}_{T'}(\varepsilon,\delta,\mathfrak{p}(T'))}(\mathbf{z},\mathbf{z}')$, *as described in Lemma 3.2.17(1).*

Lemma 3.6.2. *If* $\overline{u}_\infty \in \partial_{(-\infty,\infty)}\mathcal{M}$ *and* $\overline{v}'_* \cup \overline{v}^\sharp_* = \varnothing$ *for all levels* $\overline{v}_*$ *of* $\overline{u}_\infty$, *then* $\overline{u}_\infty$ *is one of the following:*

(1) $a = 0$, $c = 1$; $\overline{v}_0$ *is a* $\overline{W}_{T'}$*-curve with* $I = 1$ *which passes through* $\overline{\mathfrak{m}}(T')$; *and* $\overline{v}_{-1,1}$ *is a* W*-curve with* $I = 1$; *or*

(2) $a = 1$, $c = 0$; $\overline{v}_{1,1}$ *is a* W*-curve with* $I = 1$; *and* $\overline{v}_0$ *is a* $\overline{W}_{T'}$*-curve with* $I = 1$ *which passes through* $\overline{\mathfrak{m}}(T')$.

Here either $T' \in \mathcal{T}_2$ *and there is a component of* $\overline{v}_0$ *which is in*

$$\mathcal{M}^{\dagger,s,irr,\mathrm{ind}=1,n^*=m}_{\overline{J}^{\Diamond}_{T'}(\varepsilon,\delta,\mathfrak{p}(T'))}(\mathbf{z},\mathbf{z}',\overline{\mathfrak{m}}(T'))$$

from Lemma 3.2.17(2), for some $\mathbf{z}$, $\mathbf{z}'$; *or* $T' \in \mathcal{T}_1$ *and there is a component of* $\overline{v}_0$ *which does not pass through* $\overline{\mathfrak{m}}(T')$ *but is in*

$$\mathcal{M}^{\dagger,s,irr,\mathrm{ind}=-1,n^*=0}_{\overline{J}^{\Diamond}_{T'}(\varepsilon,\delta,\mathfrak{p}(T'))}(\mathbf{z},\mathbf{z}')$$

from Lemma 3.2.17(1).

Lemma 3.6.3. *If* $\overline{v}'_* \cup \overline{v}^\sharp_* \neq \varnothing$ *for some level* $\overline{v}_*$ *of* $\overline{u}_\infty$, *then:*

(1) $p_0 = \deg(\overline{v}'_0) > 0$;

(2) $\overline{u}_\infty$ *has no boundary puncture of type (*P_1*) or (*P_2*);*

(3) $\overline{u}_\infty$ *has no fiber components and no components of* $\overline{v}''_*$ *that intersect the interior of a section at infinity;*

(4) *each component of* $\overline{v}^\sharp_{-1,j}$, $j = 1, \dots, c$, *is a thin strip from* z_∞ *to some* x_i *or* x'_i *with* $I = 1$;

(5) *the only boundary punctures are type (*P_3*) punctures of* $\overline{v}^\sharp_{1,j}$, $j = 0, \dots, a$.

The following is the analog of Lemma 3.4.21, stated without proof.

Lemma 3.6.4. *If* $\overline{v}'_* \cup \overline{v}^\sharp_* \neq \varnothing$ *for some level* $\overline{v}_*$, *then* $\overline{u}_\infty$ *contains a subbuilding consisting of* $\overline{v}^\sharp_{1,a}$ *with* $I \geq 1$; $\overline{v}'_{1,j}$, $1 \leq j < a$, *which branch cover* σ_∞; $\overline{v}'_0$ *with* $I = -p$ *which is a degree* p *branched cover of* $\sigma^{T'}_\infty$; $\cup^c_{j=1}\overline{v}^\sharp_{-1,j}$ *which is a union of* p *thin strips; and possibly the following:*

- $\overline{v}^\sharp_0$ *with positive ends at multiples of* z_∞ *and no negative ends at multiples of* z_∞; *and*
- $\overline{v}^\sharp_{1,j}$, $1 \leq j < a$, *with* $I(\overline{v}^\sharp_{1,j}) \geq 1$ *and (positive or negative) ends at multiples of* z_∞.

Here at most one component of $\overline{v}''_0$ *satisfies* $I = -1$ *and the remaining components of* $\overline{v}''_*$ *satisfy* $I \geq 0$. *Also, there may be boundary punctures of type (*P_3*) with a total contribution of* $\mathfrak{bp}$ *towards* I.

See Figure 13 for some possibilities. Observe that in the current case there is at most one component of $\overline{v}''_0$ with $I = -1$ whereas there are none in Lemma 3.4.21.



FIGURE 13. Schematic diagrams for some possible types of degenerations. Here ○ represents z_∞, □ represents one or more branch points and × represents some x_i or x_i'. A vertical line indicates a trivial cylinder or a restriction of a trivial cylinder, and a triple vertical line indicates a degree p branched cover of a trivial cylinder or a restriction of a trivial cylinder. The labels on the graphs are ECH indices of each component.

Lemma 3.6.5. *For each interval $[-T, T]$, there exists $m \gg 0$ such that there is no sequence of curves $\overline{u}_i \in \mathcal{M}_{\tau_i}$, $\tau_i \to T' \in [-T, T]$, that limits to $\overline{u}_\infty$ for which $\overline{v}'_* \cup \overline{v}^\sharp_* \neq \varnothing$ for some level $\overline{v}_*$.*

Proof. This is similar to Case (6_i) of Lemma 3.4.24.

First suppose that there are no boundary punctures of type (P_3) on $\overline{v}_0$. Applying the usual rescaling argument with $m \to \infty$, we obtain $w_0 : \Sigma_0 \to \mathbb{CP}^1$ and a branched cover $\pi_0 : \Sigma_0 \to cl(B_{T'})$ such that:

(i) $w_0(\partial\Sigma_0) \subset \{\phi = 0, \rho > 0\} \cup \{\infty\}$;
(ii) $w_0(z_0) = \infty$ for some point $z_0 \in \pi_0^{-1}(+\infty)$;
(iii) $w_0(z_1) = 0$ for some point $z_1 \in \pi_0^{-1}(\overline{\mathfrak{m}}^b(T'))$;
(iv) $w_0|_{int(\Sigma_0)}$ is a biholomorphism onto its image.

Let us write $f_0 = \pi_0 \circ w_0^{-1}$ where defined. We now apply the Involution Lemma I.7.10.6. Using the notation of Lemma I.7.10.6, let $\overline{\Sigma}_1$ be the compact Riemann surface with boundary whose interior is biholomorphic to $w_0(int(\Sigma_0))$ and let $\overline{\Sigma}_2 = cl(B_{T'})$. We then extend f_0 to a holomorphic map $\overline{\Sigma}_1 \to \overline{\Sigma}_2$. By the Involution Lemma I.7.10.6 and (i), (iii) and (iv), f_0 maps the point on $\overline{\Sigma}_1$ which corresponds to $\infty \in \mathbb{CP}^1$ to $\mathcal{L}_{(r(T')+1)/2} \cap \partial cl(B_{T'})$. This contradicts (ii).

Next suppose that there are boundary punctures of type (P_3) on $\overline{v}_0$. By an ECH index count, there can be at most one boundary puncture; let $\mathfrak{q} \in cl(B_{T'})$ be the image of the boundary puncture under the projection to $cl(B_{T'})$. The rescaling argument gives $w_0 : \Sigma_0 \to \mathbb{CP}^1$ and $\pi_0 : \Sigma_0 \to cl(B_{T'})$ such that (i), (iii), (iv) hold and (ii) is replaced by (ii') $w_0(z_0)=\infty$ for some point $z_0\in\pi_0^{-1}(\mathfrak{q})$.

By the Involution Lemmas, $f_0=\pi_0\circ w_0^{-1}$ maps the point on $\overline{\Sigma}_1$ which corresponds to $\infty\in\mathbb{CP}^1$ to $\mathcal{L}_{(r(T')+1)/2}\cap\partial cl(B_{T'})$. In other words, the boundary puncture must lie on $\mathcal{L}_{(r(T')+1)/2}\cap\partial cl(B_{T'})$; this is *effectively* a codimension one condition (this is discussed in Remark 3.6.6 below). In total, the puncture contributes $+2$ towards I, the large sector contributes $+1$, and some $\overline{v}_{1,j}$, $j>0$, contributes $+1$, for a total of $I=4$, a contradiction. ☐

Remark 3.6.6. Let $\widetilde{v}$ be a component of $\overline{v}_0''\subset\overline{u}_\infty$ with a boundary puncture $\mathfrak{r}$. We justify the assertion that requiring the boundary puncture $\mathfrak{r}$ to map to a specific point in $\partial B_\tau\times\{z_\infty\}$ is *effectively* a codimension one condition. Recall that the almost complex structure $\overline{J}_\tau^\Diamond(\varepsilon,\delta,\mathfrak{p}(\tau))$ that we are using is a generic almost complex structure in a neighborhood of $K_{\mathfrak{p}(\tau),\delta}$. If $\widetilde{v}$ passes through $K_{\mathfrak{p}(\tau),\delta}$, then the neighborhood of $\widetilde{v}$ in the relevant moduli space is transversely cut out. In particular, mapping $\mathfrak{r}$ to a specific point in $\partial B_\tau\times\{z_\infty\}$ is a codimension one condition.

On the other hand, suppose that $\widetilde{v}(\varepsilon,\delta)$ (this means that $\widetilde{v}$ is dependent on ε and δ) does not intersect $K_{\mathfrak{p}(\tau),\delta}$ for all sufficiently small $\varepsilon,\delta>0$. Then a sequence of $\overline{u}_\infty(\varepsilon,\delta)$ converges to a building $\overline{u}_\infty^{bis}$ with $\deg(\overline{v}_1^{bis})'>\deg\overline{v}_1'$. Here $(\overline{v}_1^{bis})'$ is the part of $\overline{v}_1^{bis}$ that branch covers the section at infinity. Since the degree of the branched cover increases, we inductively reduce to a case which has been eliminated already.

In the rest of the paper, we often say "*effectively* a codimension l constraint" to indicate an analogous situation, where the details are left to the reader.

We are now in a position to prove Lemma 3.3.8.

Proof of Lemma 3.3.8. Suppose $\overline{u}_\infty\in\partial_{(-\infty,+\infty)}\mathcal{M}$. By Lemma 3.6.2, if $\overline{v}_*'\cup\overline{v}_*^\sharp=\varnothing$ for all levels $\overline{v}_*$ of $\overline{u}_\infty$, then $\overline{u}_\infty\in A_4$ or A_5. By Lemma 3.6.5, for any $T>0$, there exists $m\gg 0$ such that if $\overline{u}_\infty\in\partial_{[-T,T]}\mathcal{M}$, then $\overline{v}_*'\cup\overline{v}_*^\sharp=\varnothing$ for all $\overline{v}_*$.

It remains to consider the case where there exist sequences $m_i\to\infty$, $\varepsilon_i,\delta_i\to 0$, $T_i\to\infty$, and $\overline{u}_{ij}\to\overline{u}_{i\infty}$, where $\overline{u}_{ij}\in\mathcal{M}^{(m_i)}$, $\overline{u}_{i\infty}\in\partial_{\{\pm T_i\}}\mathcal{M}^{(m_i)}$, $\mathcal{M}^{(m_i)}$ is $\mathcal{M}$ with respect to the family $\{\overline{J}_\tau^\Diamond(\varepsilon_i,\delta_i,\mathfrak{p}(\tau);m_i)\}$, and $\overline{u}_{i\infty}$ satisfies $\overline{v}_*'\cup\overline{v}_*^\sharp\neq\varnothing$ for some $*$. We then take a diagonal subsequence $\overline{u}_{ij(i)}$. The proofs of Lemmas 3.4.24 and 3.5.15 carry over to give a contradiction. ☐

3.7. **Degeneration at $+\infty$, part II.** In Sections 3.7–3.9 we study the limit of holomorphic maps to $\overline{W}_\tau$ whose positive end is of the form $\mathbf{z}=\{z_{\infty,i}\}_{i\in\mathcal{I}}\cup\mathbf{y}$, i.e., we are in Step 4 of the proof of Theorem 3.3.1.

Let us write

$$\mathcal{M}:=\mathcal{M}^{I=2,n^*\le m+|\mathcal{I}|}_{\{\overline{J}_\tau^\Diamond(\varepsilon,\delta,\mathfrak{p}(\tau))\}}(\mathbf{z},\mathbf{y}';\overline{\mathfrak{m}}),\ \text{and}$$

$$\mathcal{M}_\tau:=\mathcal{M}^{I=2,n^*\le m+|\mathcal{I}|}_{\overline{J}_\tau^\Diamond(\varepsilon,\delta,\mathfrak{p}(\tau))}(\mathbf{z},\mathbf{y}';\overline{\mathfrak{m}}),$$

where $|\mathcal{I}|\geq 1$.

Let $\overline{u}_\infty \in \partial_{+\infty}\mathcal{M}$ be the limit of $\overline{u}_i \in \mathcal{M}_{\tau_i}$, where $\tau_i \to +\infty$. We use the notation from Section 3.4 for the levels and components of $\overline{u}_\infty$ and write $p_* = \deg(\overline{v}'_*)$ as before.

We now have two constraints

$$n^*(\overline{u}_i) = \sum_{\overline{v}_*} n^*(\overline{v}_*) = m + |\mathcal{I}|; \tag{3.7.1}$$

$$I(\overline{u}_i) = \sum_{\overline{v}_*} I(\overline{v}_*) = 2, \tag{3.7.2}$$

where the summations are over all the levels $\overline{v}_*$ of $\overline{u}_\infty$.

Outline of proof of Lemma 3.3.10. The proof is similar to that of Lemma 3.3.2, with the following differences: The proof of Lemma 3.7.10, which is the analog of Lemma 3.4.18, is more involved and the proof of Lemma 3.7.4, which is the analog of Lemma 3.4.9, uses a slightly more complicated notion of an "almost alternating" pair $(P_{*,0}, P_{*,1})$.

3.7.1. *Continuation argument.* First observe that Lemma 3.4.1 carries over verbatim. The analog of Lemmas 3.4.4 and 3.4.12 hinge on the continuation argument: Let $\mathcal{E}_{-,i}$, $i = 1, \dots, q$, be the negative ends of $\cup_{j=1}^{a} \overline{v}^\sharp_{1,j}$ that converge to z_∞ and let $\mathcal{E}_{+,i}$, $i = 1, \dots, r$, be the positive ends of $\cup_{j=0}^{a-1} \overline{v}^\sharp_{1,j}$ that converge to z_∞. Suppose that $q \neq 0$ (i.e., some $\mathcal{E}_{-,i}$ exists) or not all $\mathcal{E}_{+,i}$ project to thin sectors. Also for simplicity we assume that there are no punctures of type (P_3). By assumption, we may start with an end $\mathcal{E}_{-,i}$ or $\mathcal{E}_{+,i}$ that projects to a non-thin sector. The continuation argument (i.e., the proof of Lemma 3.4.4) carries over with one modification: When we are considering the continuation

$$g^0_{j_2+1,2}, \dots, g^0_{a,2}$$

of $g^0_{j_2,2}$, it is possible that $g^0_{j,2}$ is trivial for all $j_2 + 1 \le j \le a$. In other words, there is no nontrivial negative end $\mathcal{E}_{-,2}$ such that

$$\pi_{D_{\rho_0}}(\mathcal{E}_{-,2}) = \mathfrak{S}(\overline{h}(\overline{a}_{k_3,l_3}), \overline{a}_{k_4,l_4})$$

for some (k_4, l_4). This happens when $(k_3, l_3) \to (k_3, l_3)$ belongs to the data $\overrightarrow{\mathcal{D}}$ at the positive end of $\overline{v}'_{1,a}$. In this case we set $j_3 = a$ and $\overline{a}_{k_3,l_3} = \overline{a}_{k_4,l_4}$ and skip $\mathfrak{S}(\overline{h}(\overline{a}_{k_3,l_3}), \overline{a}_{k_3,l_3})$. The rest of the argument is the same.

3.7.2. *Bounds on ECH indices.* The goal of this subsection is to show the nonnegativity of $I(\overline{v}_*)$ except when $\overline{v}_* = \overline{v}_+$, *under the assumption that there are no boundary punctures that are removable at* z_∞.

Let $A_\varepsilon = \partial D_\varepsilon \times [0,1]$ for $0 < \varepsilon < \rho_0$ small and let $\pi_{[0,1]\times\overline{S}}$ be the projection of $\overline{W}$ or the positive end of $\overline{W}_\tau$ to $[0,1] \times \overline{S}$. Let $\mathfrak{c}'$ be the grooming on A_ε corresponding to the data $\overrightarrow{\mathcal{D}}'_{+,a}$ at z_∞ for the positive end of $\overline{v}'_{1,a}$, such that the winding number $w(\mathfrak{c}')$ is zero. Here the data $\overrightarrow{\mathcal{D}}'_{+,a}$ satisfies $(\mathcal{D}'_{+,a})^{to} = (\mathcal{D}'_{+,a})^{from}$. Let $\mathfrak{c}'' = \pi_{[0,1]\times\overline{S}}(\cup_i \mathcal{E}_{-,i}) \cap A_\varepsilon$. By Section 3.7.1, $\mathfrak{c}''$ is groomed and the sets of initial and terminal points of $\mathfrak{c}''$ alternate along $(0,2\pi)$. We then define P_0 and P_1 as the initial and terminal points of $\mathfrak{c}'\cup\mathfrak{c}''$.

Remark 3.7.1. It is possible for initial/terminal points of $\mathfrak{c}'$ to also appear as initial/terminal points of $\mathfrak{c}''$.

Definition 3.7.2. A pair of points $(\mathfrak{q}_0,\mathfrak{q}_1)\subset\partial D_\varepsilon$ is a *thin pair* if $\mathfrak{q}_0=\overline{h}(\overline{a}_{i,j})\cap\partial D_\varepsilon$ and $\mathfrak{q}_1=\overline{a}_{i,j}\cap\partial D_\varepsilon$ for the same i,j. A pair $(P_{*,0},P_{*,1})$ consisting of disjoint finite subsets of $(0,2\pi)$ with the same cardinality is *almost alternating along* $(0,2\pi)$ if each $P_{*,i}$, $i=0,1$, admits a splitting $P_{*,i}=Q_{*,i}\sqcup R_{*,i}$ such that the pair $(R_{*,0},R_{*,1})$ is alternating along $(0,2\pi)$ and there is a partition of $Q_{*,0}\cup Q_{*,1}$ into thin pairs $(\mathfrak{q}_0,\mathfrak{q}_1)$, $\mathfrak{q}_i\in Q_{*,i}$.

By definition, (P_0,P_1) is almost alternating along $(0,2\pi)$. See Figure 14 for an example.

Section 3.7.1 gives the main cycle

$$\mathcal{Z}_{\text{main}}=(\mathfrak{z}_0\to\mathfrak{z}_1\to\cdots\to\mathfrak{z}_{k-1}\to\mathfrak{z}_0),$$

where $\{\mathfrak{z}_1,\dots,\mathfrak{z}_k\}$ can be decomposed into an almost alternating pair and the cycle winds around $\mathbb{R}/2\pi\mathbb{Z}$ once. If we apply the continuation method to the positive end of $\overline{v}'_{1,a}$, then we also obtain a union $\mathcal{Z}_{\text{aux}}$ of auxiliary cycles of the form $(\mathfrak{z}_0\to\mathfrak{z}_1\to\mathfrak{z}_0)$, where $(\mathfrak{z}_0,\mathfrak{z}_1)$ is a thin pair, and the chords are short chords from $\mathfrak{z}_i$ to $\mathfrak{z}_{1-i}$, $i=0,1$. In view of Remark 3.7.1, the points of $\mathcal{Z}_{\text{aux}}$ can also used in $\mathcal{Z}_{\text{main}}$. The sets of initial and terminal points of $\mathcal{Z}_{\text{main}}\cup\mathcal{Z}_{\text{aux}}$ are P_0 and P_1.



FIGURE 14. The rectangular box, with the sides identified, is A_ε. The grooming $\mathfrak{c}'$ connects Q_0 to Q_1 and the grooming $\mathfrak{c}''$ connects R_0 to R_1. The vertical lines corresponding to Q_1 (resp. Q_0) are the intersections $(\overline{a}_{i,j}\cap\partial D_\varepsilon)\times[0,1]$ (resp. $(\overline{h}(\overline{a}_{i,j})\cap\partial D_\varepsilon)\times[0,1]$). The dotted lines, together with some solid lines, give the main cycle $\mathcal{Z}_{\text{main}}$. There are also two auxiliary cycles of $\mathcal{Z}_{\text{aux}}$ corresponding to the middle and right arcs of $\mathfrak{c}'$.

Let $\overrightarrow{\mathcal{D}}_{\pm,j}$ be the data at z_∞ for the $\pm$ end of $\overline{v}_{1,j}$ and let $P_{\pm,j,0}$ and $P_{\pm,j,1}$ be the initial and terminal points on A_ε determined by $\overrightarrow{\mathcal{D}}_{\pm,j}$. Then we write

$$P_{\pm,j,i}=P'_{\pm,j,i}\cup P''_{\pm,j,i},$$

where $P'_{\pm,j,i}$ corresponds to $\overline{v}'_{1,j}$ and $P''_{\pm,j,i}$ corresponds to $\overline{v}''_{1,j}$. For convenience we write $P_{-,a+1,i}=P'_{-,a+1,i}$, $i=0,1$, for the endpoints of $\mathfrak{c}'$. Also let $Q^\star_*$ and $R^\star_*$ be the thin and alternating parts of $P^\star_*$ as given in Definition 3.7.2.

The following lemmas are analogs of Lemma 3.4.8 and 3.4.9:

Lemma 3.7.3. *If $\overline{u}_\infty$ has no boundary puncture which is removable at z_∞, then, for each $* = (\pm, j)$, $P_{*,0} \subset P_0$ and $P_{*,1} \subset P_1$ and the pair $(P_{*,0}, P_{*,1})$ is almost alternating along $(0, 2\pi)$.*

Proof. Similar to but slightly more complicated than that of Lemma 3.4.8. We set $P_i^{(0)} = P_i$ and $\mathcal{Z}_*^{(0)} = \mathcal{Z}_*$, where $* =$ main or aux. Then $P_{-,a+1,i} = P'_{-,a+1,i} \subset P_i^{(0)}$ and the pair $(P_{-,a+1,0}, P_{-,a+1,1})$ is almost alternating. In $(j0)$–$(j4)$ we replace "alternating" by "almost alternating", j_0 by $a+1$, and $(j2)$ and $(j4)$ by:

- $(j2)$ there is a partition of $P_{-,a-j+1,0} \cup P_{-,a-j+1,1}$ into pairs $\{\mathfrak{p}_0, \mathfrak{p}_1\}$, $\mathfrak{p}_i \in P_{-,a-j+1,i}$, such that $\mathfrak{p}_0 \lessdot_{R_0^{(j)} \cup R_1^{(j)}} \mathfrak{p}_1$ or $\{\mathfrak{p}_0, \mathfrak{p}_1\}$ is a thin pair of $Q_0^{(j)} \cup Q_1^{(j)}$; in particular, the points of $P_{-,a-j+1,0}$ and $P_{-,a-j+1,1}$ almost alternate along $(0, 2\pi)$;
- $(j4)$ there is a partition of $P'_{+,a-j,0} \cup P'_{+,a-j,1}$ into pairs $\{\mathfrak{p}_0, \mathfrak{p}_1\}$, $\mathfrak{p}_i \in P'_{+,a-j,i}$, such that $\mathfrak{p}_0 \lessdot_{R_0^{(j+1)} \cup R_1^{(j+1)}} \mathfrak{p}_1$ or $\{\mathfrak{p}_0, \mathfrak{p}_1\}$ is a thin pair of $Q_0^{(j+1)} \cup Q_1^{(j+1)}$; in particular, the points of $P'_{+,a-j,0}$ and $P'_{+,a-j,1}$ almost alternate along $(0, 2\pi)$.

There is a partition of $P''_{+,a-j,0} \cup P''_{+,a-j,1}$ into pairs $\{\mathfrak{p}_0, \mathfrak{p}_1\}$, $\mathfrak{p}_i \in P''_{+,a-j,i}$, such that $\mathfrak{p}_1 \lessdot_{P_0^{(j)} \cup P_1^{(j)}} \mathfrak{p}_0$. We then inductively define $P_i^{(j)}$ and $\mathcal{Z}_*^{(j)}$ as follows: For each pair $\{\mathfrak{p}_0, \mathfrak{p}_1\}$ of $P''_{+,a-j,0} \cup P''_{+,a-j,1}$, if $(\mathfrak{p}_1 \to \mathfrak{p}_0)$ is a chord of $\mathcal{Z}_{\text{aux}}^{(j-1)}$, then we remove $(\mathfrak{p}_1 \to \mathfrak{p}_0 \to \mathfrak{p}_1)$ from $\mathcal{Z}_{\text{aux}}^{(j-1)}$; otherwise, in $\mathcal{Z}_{\text{main}}^{(j-1)}$, we replace $\mathfrak{q} \to \mathfrak{p}_1 \to \mathfrak{p}_0 \to \mathfrak{q}'$ by $\mathfrak{q} \to \mathfrak{q}'$, given by concatenation. Then $P_0^{(j)}$ and $P_1^{(j)}$ are the sets of endpoints of $\mathcal{Z}_{\text{main}}^{(j)} \cup \mathcal{Z}_{\text{aux}}^{(j)}$.

The verification of $(j0)$–$(j4)$ is left to the reader. □

Lemma 3.7.4. *If fiber components are removed from $\overline{u}_\infty$ and there are no boundary punctures that are removable at z_∞, then the ECH index of each level $\overline{v}_* \neq \overline{v}_+$ is nonnegative, the only components of $\overline{u}_\infty$ which have negative ECH index are branched covers of σ_∞^+, and $I(\overline{v}_{1,j}) \geq I(\overline{v}'_{1,j}) + I(\overline{v}''_{1,j})$ for $0 \leq j \leq a$.*

Before embarking on the proof of Lemma 3.7.4 we encourage the reader to review Section I.5.7.10 and in particular Lemmas I.5.7.19 and I.5.7.20.

Proof. Similar to the proof of Lemma 3.4.9. The levels $\overline{v}_{0,j}$, $1 \leq j \leq b$, satisfy $I(\overline{v}_{0,j}) \geq 0$ by [HT1, Proposition 7.15(a)].

In order to treat the case of $\overline{v}_{1,j}$, $0 \leq j \leq a$, we use Lemma 3.7.3 and Lemmas I.5.7.19 and I.5.7.20 to compute $I(\overline{v}_{1,j})$. Let

$$\pi_{[0,1]\times\overline{S}} : \check{\overline{W}} = [-1,1] \times [0,1] \times \overline{S} \to [0,1] \times \overline{S}$$

be the projection along $[-1,1]$.

We claim that there exist representatives $\check{C}_1, \check{C}_2, \check{C}_3 \subset \check{\overline{W}}$ such that $\check{C}_1 \cup \check{C}_2$ is a representative of $\overline{v}'_{1,j}$, $\check{C}_3$ is a representative of $\overline{v}''_{1,j}$, and the following hold:

(1) the components of $\pi_{[0,1]\times\overline{S}}(\check{C}_1|_{s=\pm1})$ corresponding to z_∞ are contained in $A_{\varepsilon/2}$ and their union $\mathfrak{c}_1^\pm$ is groomed;
(2) the components of $\pi_{[0,1]\times\overline{S}}(\check{C}_i|_{s=\pm1})$, $i=2,3$, corresponding to z_∞ are contained in A_ε; if we write $\mathfrak{c}_i^\pm$ for their unions, then $\mathfrak{c}_2^+\cup\mathfrak{c}_3^+$ and $\mathfrak{c}_2^-\cup\mathfrak{c}_3^-$ are groomed;
(3) $\mathfrak{c}_1^+=\mathfrak{c}_1^-$ and they both correspond to data $\overrightarrow{\mathcal{D}}$ where $\mathcal{D}^{to}=\mathcal{D}^{from}$;
(4) $w(\mathfrak{c}_2^+\cup\mathfrak{c}_3^+)=0$ or -1 and $w(\mathfrak{c}_2^-\cup\mathfrak{c}_3^-)=0$ or 1;
(5) $\mathfrak{c}_2^+\cup\mathfrak{c}_3^+$ satisfies (G_4') from Section I.5.7.10 and $\mathfrak{c}_2^-\cup\mathfrak{c}_3^-$ satisfies (G_4) from Section I.5.7.10.

Let $\check{C}_3$ be a representative of $\overline{v}''_{1,j}$ such that the first sentence of (2) holds for $i=3$ and $\mathfrak{c}_3^\pm=\pi_{[0,1]\times\overline{S}}(\cup_k\mathcal{E}_{\pm,k})\cap A_\varepsilon$, where $\mathcal{E}_{\pm,k}$ are the $\pm$ ends of $\overline{v}''_{1,j}$ that limit to z_∞. The representative $\check{C}_1\cup\check{C}_2$ of $\overline{v}'_{1,j}$ is chosen as follows: Let $\check{C}_1=[-1,1]\times\mathfrak{c}_1^+$ so that (1) and (3) hold and the endpoints of $\mathfrak{c}_1^+=\mathfrak{c}_1^-$ are partitioned into thin pairs of $(P'_{+,j,0},P'_{+,j,1})$. Let $\check{C}_2\subset[-1,1]\times[-1,1]\times D_\varepsilon$ be a disk such that the first sentence of (2) holds for $i=2$, $\check{C}_2|_{s=\pm1}=\mathfrak{c}_2^\pm$, $\mathfrak{c}_2^\pm$ are groomed, $w(\mathfrak{c}_2^+)=0$ or -1, $w(\mathfrak{c}_2^-)=0$ or 1, and the endpoints of $\mathfrak{c}_2^\pm$ are alternating along $(0,2\pi)$. This is possible by $(j4)$ in the proof of Lemma 3.7.3. The second sentence of (2), as well as (4) and (5), follow from the proof of Lemma 3.7.3.

We now prove that $I(\overline{v}_{1,j})\geq 0$ for $j>0$; the verification of $I(\overline{v}_{1,j})\geq I(\overline{v}'_{1,j})+I(\overline{v}''_{1,j})$ for $0\leq j\leq a$ is similar and is left to the reader. First observe that $I(\check{C}_1)=0$ and $I(\check{C}_2)=0$, where the latter follows from the proof of Lemma I.5.7.12. Also $I(\check{C}_3)\geq 0$ by the ECH index inequality. Next observe that $\langle\check{C}_1,\check{C}_2\rangle=-l$, where l is the degree of $\check{C}_1$. Hence the contribution of $\check{C}_1\cap\check{C}_2$ to $I(\overline{v}_{1,j})$ is $-2l$. We now apply Lemmas I.5.7.19 and I.5.7.20, where we split into $\check{C}_1$ and $\check{C}_2\cup\check{C}_3$ instead of $\overline{v}'_{1,j}$ and $\overline{v}''_{1,j}$. Observe that $q_2=0$ (resp. $q_2=1$) in Lemma I.5.7.19 corresponds to $q_2=-1$ (resp. $q_2=0$) in Lemma I.5.7.20. One can verify that the extra contributions to I from the rightmost terms of Equations (I.5.7.9) and (I.5.7.10) add up to at least $2l$. Hence $I(\overline{v}_{1,j})\geq 0$ for $j>0$.

The case of $\overline{v}_{-1,j}$, $1\leq j\leq c+1$, is easier and will be omitted. □

3.7.3. *Boundary punctures.* In this subsection we describe the necessary modifications when $\overline{u}_\infty$ has boundary punctures that are removable at z_∞. We use the notation from Section 3.4.3, with some modifications: The continuation method gives rise to a main cycle $\mathcal{Z}_{\text{main}}$ which winds around $\mathbb{R}/2\pi\mathbb{Z}$ once and a union $\mathcal{Z}_{\text{aux}}$ of auxiliary cycles. Also, in Definition 3.4.10 we replace $\mathcal{Z}$ by $\mathcal{Z}_{\text{main}}$.

Lemma 3.7.5. *If fiber components are removed from $\overline{u}_\infty$, then the ECH index of each level $\overline{v}_*\neq\overline{v}_+$ is nonnegative, the only components of $\overline{u}_\infty$ which have negative ECH index are branched covers of σ_∞^+, and*

$$I(\overline{v}_{1,j})\geq I(\overline{v}'_{1,j})+I(\overline{v}''_{1,j})+\max(\mathfrak{bp}_{1,j},\mathfrak{bp}'_{1,j}) \tag{3.7.3}$$

for $0\leq j\leq a$, where $\mathfrak{bp}_*$ is the number of boundary punctures of type (P_3) on $\overline{v}_*$ and $\mathfrak{bp}'_*=2p_*$ if there is a boundary puncture of type (P_2) on $\overline{v}_*$ and $\mathfrak{bp}'_*=0$ otherwise. Here $p_*=\deg(\overline{v}'_*)$.

According to the lemma, each boundary puncture contributes at least 1 towards I; the proof will give a better lower bound.

Proof. We assume (S) from Section 3.4.3 for simplicity. Let $\mathfrak{r}$ be the boundary puncture on $\overline{v}_{*_0}$ with $*_0 = (1, j_0)$, $j_0 > 0$.

Suppose $\mathfrak{r}$ is of type (P$_3$). We compute the extra contribution I_{L-} to $I(\overline{v}^{\circ,1}_{*_0})$ that comes from the left negative end at z_∞ as in Lemma 3.4.14. First observe that $P_{L-,j_0,1/2} = P_{L-,j_0,1}$. By (S), $\mathcal{Z}_{\text{main}}$ has only one chord $\mathfrak{z}'_1 \to \mathfrak{z}'_s$ corresponding to a sector of type $\mathfrak{S}(\overline{a}_{k,l}, \overline{a}_{k',l'})$. Hence we have:

$$P''_{L-,j_0,1/2} = \{\mathfrak{z}'_1\}, \quad P'_{L-,j_0,1/2} = \{\mathfrak{z}'_2, \dots, \mathfrak{z}'_s, \mathfrak{z}''_1, \dots, \mathfrak{z}''_t\},$$
$$P''_{L-,j_0,1} = \{\mathfrak{z}'_s\}, \quad P'_{L-,j_0,1} = \{\mathfrak{z}'_1, \dots, \mathfrak{z}'_{s-1}, \mathfrak{z}''_1, \dots, \mathfrak{z}''_t\},$$

where $P_{L-,j_0,1/2} = P_{L-,j_0,1}$ is written as $\{\mathfrak{z}'_1, \dots, \mathfrak{z}'_s, \mathfrak{z}''_1, \dots, \mathfrak{z}''_t\}$ in cyclic order around $\mathbb{R}/2\pi\mathbb{Z}$. Note that $\mathfrak{z}'_2, \dots, \mathfrak{z}'_{s-1}$ are points of $\mathcal{Z}_{\text{aux}}$ but not $\mathcal{Z}_{\text{main}}$; otherwise $\mathcal{Z}_{\text{main}}$ winds more than once around $\mathbb{R}/2\pi\mathbb{Z}$. The projection of the left negative end of $\overline{v}^{\circ,1,''}_{*_0}$ that limits to z_∞ intersects $A^{[1/2,1]}_{\varepsilon}$ along an arc $\mathfrak{c}''_{L-}$ with winding number $w(\mathfrak{c}''_{L-}) = 0$ or 1, depending on whether $\mathfrak{S}(\overline{a}_{k,l}, \overline{a}_{k',l'})$ is a large sector. The left negative ends of $\overline{v}^{\circ,1,'}_{*_0}$ that limit to z_∞ can modified to give a grooming $\mathfrak{c}'_{L-}$ on $A^{[1/2,1]}_{\varepsilon/2}$ such that the winding number $w(\mathfrak{c}'_{L-}) = 0$ or -1 and $\mathfrak{c}'_{L-}$ connects $\mathfrak{z}''_i$ to $\mathfrak{z}''_i$, $i = 1, \dots, t$, by vertical arcs. Hence it suffices to assume that $t = 0$. The writhe of $\mathfrak{c}'_{L-} \cup \mathfrak{c}''_{L-}$ is equal to the number $\#\mathfrak{c}'_{L-}$ of components of $\mathfrak{c}'_{L-}$; see Figure 15. Resolving the (positive) crossings of $\mathfrak{c}'_{L-} \cup \mathfrak{c}''_{L-}$ yields a grooming with vertical arcs from $\mathfrak{z}'_i$ to $\mathfrak{z}'_i$. Hence the left negative end at z_∞ contributes an extra $\#\mathfrak{c}'_{L-}$ towards I.


$\mathfrak{z}'_1$ $\mathfrak{z}'_2$ $\mathfrak{z}'_3$ $\cdots$ $\mathfrak{z}'_s$
=
=
$\mathfrak{z}'_1$ $\mathfrak{z}'_2$ $\mathfrak{z}'_3$ $\cdots$ $\mathfrak{z}'_s$

FIGURE 15

The cycles $\mathcal{Z}^{(a-j)}_{\text{main}}$ and $\mathcal{Z}^{(a-j)}_{\text{aux}}$ are defined as before, for $j \geq j_0$. We define $\mathcal{Z}^{(a-j_0,+)}_{*} = \mathcal{Z}^{(a-j_0)}_{*}$, $\mathcal{Z}^{(a-j_0,-)}_{\text{aux}} = \mathcal{Z}^{(a-j_0,+)}_{\text{aux}}$, and $\mathcal{Z}^{(a-j_0,-)}_{\text{main}}$ as $\mathcal{Z}^{(a-j_0,+)}_{\text{main}}$ with $\mathfrak{z}'_1 \to \mathfrak{z}'_s$ replaced by $\mathfrak{z}'_s$. Also, $P^{\star}_{R-,j_0,i}$, $\star = \varnothing, ', ''$, is obtained from $P^{\star}_{+,j_0,i}$ by replacing $\mathfrak{z}'_1$ by $\mathfrak{z}'_s$. The rest of the arguments of Lemmas 3.7.3 and 3.7.4 carry over.

Suppose $\mathfrak{r}$ is of type (P$_2$). In this case (S) implies that $\mathfrak{z}'_1 = \mathfrak{z}'_s$ and $\mathcal{Z}_{\text{main}} = (\mathfrak{z}'_1 \to \mathfrak{z}'_1)$. Then:

$$P''_{L-,j_0,1/2} = \{\mathfrak{z}'_1\}, \quad P'_{L-,j_0,1/2} = \{\mathfrak{z}''_1, \dots, \mathfrak{z}''_{p_{*_0}}\},$$
$$P''_{L-,j_0,1} = \{\mathfrak{z}'_1\}, \quad P'_{L-,j_0,1} = \{\mathfrak{z}''_1, \dots, \mathfrak{z}''_{p_{*_0}}\}.$$

Also $\mathfrak{c}''_{L-}$ is an arc from $\mathfrak{z}'_1$ to itself with $w(\mathfrak{c}''_{L-}) = 1$ and $\mathfrak{c}'_{L-}$ consists of p_{*0} vertical arcs from $\mathfrak{z}''_i$ to itself. The extra contribution to I from the negative end at z_∞ is $2p_{*0}$: in order to groom $\mathfrak{c}'_{L-} \cup \mathfrak{c}''_{L-}$ (so that the result $\mathfrak{c}$ satisfies $w(\mathfrak{c}) = 0$), we switch the (positive) crossings of $\mathfrak{c}'_{L-} \cup \mathfrak{c}''_{L-}$ while keeping the same endpoints. $\square$

The following is a consequence of Section 3.7.1:

Lemma 3.7.6. *Suppose $\overline{v}'_{1,j} \cup \overline{v}^\sharp_{1,j} \neq \varnothing$ for some $j > 0$. Let $\mathcal{E}_{-,i}$, $i = 1, \dots, q$, be the negative ends of $\cup_{j=1}^a \overline{v}^\sharp_{1,j}$ that converge to z_∞, let $\mathcal{E}_{+,i}$, $i = 1, \dots, r$, be the positive ends of $\cup_{j=0}^{a-1} \overline{v}^\sharp_{1,j}$ that converge to z_∞, and let $\mathcal{E}'_i$, $i = 1, \dots, s$, be the neighborhoods of the punctures of type (P$_3$). Then*

$$n^+(\mathcal{E}_{-,i}) \geq k_0 - 1 \gg 2g \tag{3.7.4}$$

for each i, where the constant k_0 is as given in Section I.5.2.2. If $q \neq 0$ (i.e., some $\mathcal{E}_{-,i}$ exists) or not all $\mathcal{E}_{+,i}$ project to thin sectors, then

$$n^+((\cup_{i=1}^q \mathcal{E}_{-,i}) \cup (\cup_{i=1}^r \mathcal{E}_{+,i}) \cup (\cup_{i=1}^s \mathcal{E}'_i)) \geq m - p_+, \tag{3.7.5}$$

where $p_+ = \deg(\overline{v}'_+)$. Moreover,

$$D_{\rho_0} - (\cup_{i=1}^q \pi_{D_{\rho_0}}(\mathcal{E}_{-,i})) \cup (\cup_{i=1}^r \pi_{D_{\rho_0}}(\mathcal{E}_{+,i})) \cup (\cup_{i=1}^s \pi_{D_{\rho_0}}(\mathcal{E}'_i))$$

consists of at most p_+ thin sectors.

The following is the analog of Lemma 3.4.5:

Lemma 3.7.7. *If $\mathfrak{p}_1, \dots, \mathfrak{p}_s$ are the punctures of type (P$_1$) and (P$_2$) and $N(\mathfrak{p}_i) \subset F_*$ is a small neighborhood of $\mathfrak{p}_i$, then $\sum_{i=1}^s n^*(\overline{v}_*(N(\mathfrak{p}_i))) \geq m$.*

3.7.4. *Higher order asymptotic eigenfunctions.* Fix $m \gg 0$ and let $\overline{u}_i \to \overline{u}_\infty \in \partial_{\{+\infty\}}\mathcal{M}$. In this subsection we use higher order asymptotic expansions to give restrictions on $\overline{u}_\infty$. The reader is referred to Siefring [Si] for a discussion of higher order asymptotic expansions near an end.

Lemma 3.7.8. *The is no $\overline{u}_\infty \in \partial_{\{+\infty\}}\mathcal{M}$ such that $\deg(\overline{v}'_*) = p_+ \geq 1$, $\overline{v}^\sharp_* = \varnothing$, and $\overline{v}'_* \cap \overline{v}''_* = \varnothing$ for all $\overline{v}_* \succeq \overline{v}_+$ and $\overline{v}^\sharp_{0,b}$ is a union of p_+ cylinders from δ_0 to h or e.*

Proof. We restrict $\overline{u}_i$ to a neighborhood of $\sigma_\infty^{\tau_i}$ and further restrict it to the components that are close to $\overline{v}'_+$. After projecting to D_{ρ_0} using balanced coordinates, applying the ansatz given by Equation (I.7.9.1), rescaling, and taking the SFT limit, we obtain a holomorphic map $w_+ : \Sigma_+ \to \mathbb{CP}^1$ together with the branched cover $\pi_+ : \Sigma_+ \to cl(B_+)$. By considerations of n^*, w_+ maps each component of $\partial\Sigma_+$ to a distinct $\mathfrak{S}(\overline{a}_{i,j}, \overline{h}(\overline{a}_{i,j}))$. Hence we may assume without loss of generality that $p_+ = 1$ and $\Sigma_+ = B_+$. We identify $cl(B_+)$ with the closed unit disk $\overline{\mathbb{D}} = \{|z| \leq 1\}$ by sending $+\infty$ to 1 and $-\infty$ to 0.

The Lagrangian boundary condition for $\overline{u}_i$, when projected to D_{ρ_0} using balanced coordinates, descends to the boundary condition $w_+(e^{i\theta}) \in \mathcal{R}_{\varphi(e^{i\theta})}$, where the map $\varphi : \partial\mathbb{D} - \{1\} \to S^1 = \mathbb{R}/2\pi\mathbb{Z}$ satisfies the following:

- $\lim_{\theta\to 0^+}\varphi(e^{i\theta})=\phi(\overline{a}_{i_0,j_0})$ and $\lim_{\theta\to 0^-}\varphi(e^{i\theta})=\phi(\overline{h}(\overline{a}_{i_0,j_0}))$ for some (i_0,j_0);
- $\varphi(e^{i\theta})\in(0,2\pi)$ is a nondecreasing function of $\theta\in(0,2\pi)$; and
- $\varphi\circ i_0=i_1\circ\varphi$, where $i_0:\partial\mathbb{D}-\{1\}\stackrel{\sim}{\to}\partial\mathbb{D}-\{1\}$ is the reflection across the x-axis and $i_1:S^1\stackrel{\sim}{\to}S^1$ is the reflection across $\mathcal{R}_{\phi_0}$ where $\phi_0=(\phi(\overline{a}_{i_0,j_0})+\phi(\overline{h}(\overline{a}_{i_0,j_0})))/2$.

Lemma 3.7.9. *The holomorphic map* $w_+:\overline{\mathbb{D}}\to\mathbb{CP}^1$ *satisfies the following:*

(i) $w_+(1)=0$ *and* $w_+(e^{i\theta})\in\mathcal{R}_{\varphi(e^{i\theta})}$ *for* $e^{i\theta}\neq 1$;
(ii) $w_+(0)\in\mathcal{R}_{\phi_0}$ *and* w_+ *maps* $\mathcal{R}_0\cup\mathcal{R}_\pi$ *to* $\mathcal{R}_{\phi_0}$;
(iii) $\operatorname{Im}(w_+)\subset\mathfrak{S}(\overline{a}_{i_0,j_0},\overline{h}(\overline{a}_{i_0,j_0}))$;
(iv) w_+ *is a biholomorphism onto its image.*

Moreover, w_+ *is the unique holomorphic map* $\overline{\mathbb{D}}\to\mathbb{CP}^1$ *which satisfies (i), (iii) and (iv), up to multiplication by a positive real constant.*

Proof. (i), (iii), and (iv) are immediate from the construction. To prove the uniqueness statement, suppose there exists $\widetilde{w}_+$ satisfying (i), (iii) and (iv). Then we consider $\frac{w_+}{\widetilde{w}_+}$ as in Section I.7.10. It is easy to see that $\frac{w_+}{\widetilde{w}_+}$ is a positive real constant on $\partial\mathbb{D}$ and has no zeros or poles on $\mathbb{D}$, hence is a constant map. (ii) then follows from the uniqueness, Observation I.7.10.1, and the fact that i_0 and i_1 extend to involutions of $\mathbb{D}$ and $\mathbb{CP}^1$. Here we are viewing S^1 as the unit circle in $\mathbb{CP}^1$. $\square$

The proof of Lemma 3.7.8 now follows from Lemma 3.7.9(ii). We first eliminate the case where $\overline{v}^\sharp_{0,b}$ is a cylinder from δ_0 to h. This is due to the choice of h from Section 3.1.4: the asymptotic eigenfunction of $\overline{v}^\sharp_{0,b}$ at the positive end δ_0 is a constant $c\in\mathcal{R}_{\phi=-2\pi/m}$. This contradicts the first statement of Lemma 3.7.9(ii).

Next we eliminate the case where $\overline{v}^\sharp_{0,b}$ is a cylinder from δ_0 to e. This is done by looking at the next higher order term in the asymptotic expansion at the positive end that limits to δ_0. If we project the positive end to D_{ρ_0} using balanced coordinates and apply the ansatz, then the asymptotic expansion is of the form $c_0+c_1e^{-s-\pi it}$ as $s\to+\infty$. The first statement of Lemma 3.7.9(ii) determines c_0 up to a positive real constant and the second statement of Lemma 3.7.9(ii) determines c_1 up to a positive real constant. In other words, generically Lemma 3.7.9(ii) imposes two independent conditions; the proof is similar to that of [HT2, Section 3]. Hence $\overline{v}^\sharp_{0,b}$ must have $I\geq 3$, a contradiction. $\square$

3.7.5. *Preliminary restrictions.*

Lemma 3.7.10. *If* $m\gg 0$ *and* $\overline{u}_\infty\in\partial_{\{+\infty\}}\mathcal{M}$, *then the following hold:*

(1) $\overline{u}_\infty$ *has no fiber components;*
(2) *there is no level* $\overline{v}_*$ *such that* $\overline{v}'_*\cap\overline{v}''_*\neq\varnothing$ *and* $\overline{v}'_*\cap\overline{v}''_*\subset int(\overline{v}'_*)$; *and*
(3) $\overline{u}_\infty$ *has no boundary puncture which is removable at* z_∞.

Proof. First observe that (1)–(3) are mutually exclusive by considerations of n^*.

(1) Arguing by contradiction, suppose that $\overline{u}_\infty$ has a fiber component $\widetilde{v}:\dot{F}\to\overline{W}_*$. The fiber component $\widetilde{v}$ satisfies $n^*(\widetilde{v})\geq m$.

We first claim the following:

(a) The only ends of $\overline{v}^\sharp_*$ that limit to multiples of z_∞ or δ_0 are positive ends.

(b) All the positive ends $\mathcal{E}_+$ of $\overline{v}^\sharp_{0,j}$, $j = 0, \dots, b$, that limit to δ_0^r satisfy $n^*(\mathcal{E}_+) = r$.

(c) All the positive ends of $\overline{v}^\sharp_{1,j}$, $j = 0, \dots, a$, and $\overline{v}^\sharp_{-1,j}$, $j = 1, \dots, c$, that limit to z_∞ project to thin sectors under $\pi_{D_{\rho_0}}$.

(d) $p_{*_1} \leq p_{*_2}$ if $\overline{v}_{*_1} \preceq \overline{v}_{*_2}$.

(a) Arguing by contradiction, let $\mathcal{E}_-$ be a negative end of some $\overline{v}^\sharp_*$ that limits to a multiple of z_∞ or δ_0. Then $n^*(\mathcal{E}_-) \gg 2g$ by Lemmas 3.4.1(1) and 3.7.6. Since $n^*(\widetilde{v}) \geq m$, such a negative end $\mathcal{E}_-$ cannot exist by Equation (3.7.1). The arguments for (b) and (c) are similar and (d) is a consequence of (a)–(c).

Next we have the following contributions towards I:

(β_1) $I(\overline{v}'_+) = -p_{0,b+1}$.

(β_2) $\cup_{j=1}^b \overline{v}^\sharp_{0,j}$ is a union of $p_{0,b+1} - p_{0,1}$ cylinders from δ_0 to h or e and

$$\sum_{j=1}^{b} I(\overline{v}^\sharp_{0,j}) \geq p_{0,b+1} - p_{0,1}.$$

(β_3) $\overline{v}^\sharp_-$ is a union of $p_{0,1} - p_{0,0}$ sections from δ_0 to x_i or x'_i and

$$I(\overline{v}^\sharp_-) = p_{0,1} - p_{0,0}.$$

(β_4) $\cup_{j=1}^c \overline{v}^\sharp_{-1,j}$ is a union of $p_{0,0}$ trivial strips and

$$\sum_{j=1}^{c} I(\overline{v}^\sharp_{-1,j}) = p_{0,0}.$$

(β_5) If the fiber $\widetilde{v}$ is a component of $\overline{v}_*$, then $\widetilde{v}$ and the intersection $\widetilde{v} \cap (\overline{v}_* - \widetilde{v})$ contribute $2g + 2 \geq 4$ towards I.

(β_6) After removing the fiber component $\widetilde{v}$, all the levels $\neq \overline{v}_+$ have nonnegative ECH index, $I(\overline{v}''_+) \geq 0$, and $I(\overline{v}_+) \geq I(\overline{v}'_+) + I(\overline{v}''_+)$.

(β_6) follows from Lemma 3.7.4 and (β_1)–(β_4) are clear.

We now prove (β_5). Suppose that $\widetilde{v}$ is a component of $\overline{v}_{*_0}$. If $\overline{v}''_{*_0}$ has a boundary puncture which is removable at z_∞, then a neighborhood of the puncture contributes $n^* \gg 2g$. Since $n^*(\widetilde{v}) \geq m$, we have a contradiction of Equation (3.7.1). Hence all the non-fiber components of $\overline{v}_{*_0}$ that pass through z_∞ are components of $\overline{v}'_{*_0}$. When $\widetilde{v}$ is an interior fiber, (β_5) follows from Equation (I.7.6.6). Next suppose that $\widetilde{v}$ is a boundary fiber, i.e., $\widetilde{v}(\widetilde{F}) \subset \{p\} \times \overline{S}$, where $\widetilde{F}$ is the domain of $\widetilde{v}$ and $p \in \partial B_{*_0}$. If $(p, z_\infty) \notin \widetilde{v}(\partial\widetilde{F})$, i.e., $\widetilde{v}(\partial\widetilde{F})$ consists of $2g$ slits along $\widehat{\mathbf{a}}$, for example, then $\overline{v}'_{*_0} = \varnothing$ and the Fredholm index of $\widetilde{v}$ is:

$$\mathrm{ind}(\widetilde{v}) = -\chi(\widetilde{F}) + \mu_\tau(\partial\widetilde{F}) + 2c_1(\widetilde{v}^* T\overline{S}) = -(2-4g) + 0 + 2(2-2g) = 2,$$

where τ is some trivialization of $T\overline{S}$ which agrees with $T\widehat{\mathbf{a}}$. Since there are $2g$ points that are pinched, $\widetilde{v}$ and the pinched points contribute a total of $2g+2$ towards I. Now suppose that $(p, z_\infty) \in \widetilde{v}(\partial\widetilde{F})$. For simplicity we assume that $\overline{v}''_{*_0} = \varnothing$

and that $*_0 = (1,j)$ with $j \geq 1$. By considerations of n^*, all the components of the data $\overrightarrow{\mathcal{D}}_\pm$ are of the form $(i,j) \to (i,j)$. Let τ_0 be a groomed multivalued trivialization which is compatible with $\overrightarrow{\mathcal{D}}_\pm$, and suppose that the corresponding groomings $\mathfrak{c}_\pm \subset A_\varepsilon = [0,1] \times \partial D_\varepsilon$ satisfy $\mathfrak{c}_+ = \mathfrak{c}_-$. In order to construct a τ_0-trivial representative $\check{C}$ of $\widetilde{v} \cup \overline{v}'_{*_0}$, we consider $\overline{u}_i$ which is close to breaking and truncate it so that the truncation is close to $\widetilde{v} \cup \overline{v}'_{*_0}$. Note that the boundary of $\overline{u}_i$ does not intersect the section at infinity. By applying Steps 2 and 3 of Section I.5.7.5 to the truncation, we obtain $[\check{C}] \in \pi_2(\mathbf{z}_+, \mathbf{z}_-, \tau_0)$, where $\mathbf{z}_\pm = \{z_\infty^{2g}(\overrightarrow{\mathcal{D}}_\pm)\}$. We easily verify that $[\check{D}] = [\check{C}] - [\overline{S}] = [[-1,1] \times \mathfrak{c}]$, where $[\overline{S}]$ is the homology class of an interior fiber. Hence $[\check{C}] = [[-1,1] \times \mathfrak{c}] + [\overline{S}] \in \pi_2(\mathbf{z}_+, \mathbf{z}_-, \tau_0)$ and (β_5) follows from the calculation in the case of a closed $\widetilde{v}$.

Summing (β_1)–(β_6), the total ECH index is ≥ 4, which contradicts Equation (3.7.2). Hence (1) follows.

(2) Arguing by contradiction, suppose there exist sequences $m_l \to \infty$ and $\overline{u}_{li} \to \overline{u}_{l\infty}$, where $\overline{u}_{li} \in \mathcal{M}^{(m_l)}$, $\overline{u}_{l\infty} \in \partial_{\{+\infty\}}\mathcal{M}^{(m_l)}$, and $\mathcal{M}^{(m_l)}$ is $\mathcal{M}$ with respect to m_l, such that each $\overline{u}_{l\infty}$ has some $\overline{v}_{l,*_0}$ such that $\overline{v}'_{l,*_0} \cap \overline{v}''_{l,*_0} \neq \varnothing$. Unless indicated otherwise, we fix $l \gg 0$ and suppress l from the notation.

The same argument as in (1) implies that (a)–(d) hold. Since $n^*(\overline{u}_i) = m + |\mathcal{I}| \leq m + 2g$, there is only one intersection point of $\overline{v}'_{*_0}$ and $\overline{v}''_{*_0}$, which we denote by $\mathfrak{r} = (\mathfrak{r}^b, z_\infty)$.

(2A) Suppose that $*_0 = +$. As in the proof of (1), the holomorphic building hanging from the positive end of $\overline{v}'_+$ has total ECH index ≥ 0 — this is obtained by adding all the ECH index contributions from (β_1)–(β_4). We also have (β'_5) instead of (β_5):

- (β'_5) $\overline{v}'_{*_0} \cap \overline{v}''_{*_0}$ contributes $2 \cdot m(\mathfrak{r})$ towards I, where $m(\mathfrak{r}) \geq 1$ is the multiplicity of $\mathfrak{r}$.

Hence $\sum_{\overline{v}_*} I(\overline{v}_*) \geq 2 \cdot m(\mathfrak{r})$. This implies that $m(\mathfrak{r}) = 1$ and $p_+ = \deg \overline{v}'_+ = 1$. The sum of the ECH indices from (β_1)–(β_4), (β'_5), (β_6), and $I(\overline{v}''_+) \geq 0$ is at least $+2$. We also have $p_+ = 1$.

We claim that $p_- \geq 1$ by considerations of n^*. Indeed, if $\overline{v}'_- = \varnothing$, then there exist a point $\mathfrak{q} \in int(F_-)$ and a sufficiently small neighborhood $N(\mathfrak{q}) \subset \dot{F}_-$ of $\mathfrak{q}$ such that $\overline{v}_-(\mathfrak{q}) = \overline{\mathfrak{m}}(+\infty)$ and $n^*(\overline{v}_-(N(\mathfrak{q}))) \geq m$. This is a contradiction.

Since $p_+ = 1$ and $p_- \geq 1$, it follows that $p_{0,j} = 1$ for all $j = 0, \dots, b+1$ by (d). For the purposes of applying the rescaling argument, we may assume that $b = 0$. By restricting $\overline{u}_{li}$ to a neighborhood of $\sigma_\infty^{\tau_{li}}$, taking the limit of a diagonal subsequence $\overline{u}_{li(l)}$, and applying the rescaling argument, we obtain a 2-level holomorphic building $w_+ \cup w_-$, where $w_\pm : cl(B_\pm) \to \mathbb{CP}^1$ satisfy the following:

- (i) $w_-(\partial B_-) \subset \{\phi = 0, \rho > 0\}$;
- (ii) $w_-(\overline{\mathfrak{m}}^b(+\infty)) = 0$ and $w_-(+\infty) = \infty$;
- (iii) $w_+(\partial B_+) \subset \{\phi = 0, \rho > 0\}$;
- (iv) $w_+(-\infty) = 0$ and $w_+(\mathfrak{r}^b) = \infty$;

(v) $w_\pm|_{int(B_\pm)}$ is a biholomorphism onto its image.

Then, by the Involution Lemma I.7.10.3, w_- maps the marker $\dot{\mathcal{L}}_{3/2}(+\infty)$ to $\dot{\mathcal{R}}_\pi(\infty)$ and w_+ maps the ray $\mathcal{L}_{3/2}$ to the ray $\mathcal{R}_\pi$. This *effectively* (cf. Remark 3.6.6) gives rise to two constraints for $\overline{v}''_+$ and hence $I(\overline{v}''_+) \geq 2$. Hence the total ECH index is at least 4, a contradiction.

(2B) Suppose that $*_0 = (1, j_0)$ for some $1 \leq j_0 \leq a$. Summing (β_1)–(β_4), (β_5'), (β_6), and $I(\overline{v}''_{*_0}) \geq 1$, the total ECH index is at least 3, a contradiction. However, we want to do slightly better so that the argument carries over in the case of Lemma 3.9.2: The rescaling argument with $m_l \to \infty$ gives $w_+ : cl(B_+) \to \mathbb{CP}^1$ satisfying the following:

(i) $w_+(\partial B_+) \subset \{\phi = 0, \rho > 0\}$;
(ii) $w_+(-\infty) = 0$ and $w_+(+\infty) = \infty$;
(iii) w_+ maps the marker $\dot{\mathcal{L}}_{3/2}(-\infty)$ to $\dot{\mathcal{R}}_\pi(0)$;
(iv) $w_+|_{int(B_+)}$ is a biholomorphism onto its image.

By the Involution Lemmas, w_+^{-1} maps the component of $\{\mathrm{Im}(z) = 0\}$ passing through 0 to $\mathcal{L}_{3/2}$. This contradicts (ii).

(2C) Suppose that $*_0 = (0, j_0)$ for some $1 \leq j_0 \leq b$. We replace (β_2) by:

(β_2') If $j = j_0$, then each positive end of $\overline{v}^\sharp_{0,j_0}$ that limits to δ_0 and is not the end of a cylinder from δ_0 to h or e contributes $+2$ towards I.
(β_2'') All the other components of $\overline{v}^\sharp_{0,j}$, $1 \leq j \leq b$, are cylinders from δ_0 to h or e with $I = 1$ or 2.

(β_2'') is clear and (β_2') is a consequence of the argument of Lemma 3.7.8. Summing (β_1), (β_2'), (β_2''), (β_3), (β_4), (β_5'), (β_6), and $I(\overline{v}''_{*_0}) \geq 1$, the total ECH index is at least 3, a contradiction.

(2D) Suppose that $*_0 = -$. In a manner similar to (2C) we replace (β_3) by:

(β_3') Each positive end of $\overline{v}^\sharp_-$ that limits to δ_0 and is not the end of a section from δ_0 to x_i or x_i' contributes $+2$ towards I.
(β_3'') All the other components of $\overline{v}^\sharp_-$ are sections from δ_0 to x_i or x_i' with $I = 1$.

Summing (β_1), (β_2), (β_3'), (β_3''), (β_4), (β_5'), (β_6), and $I(\overline{v}''_-) \geq 0$, the total ECH index is at least 2.

Suppose that $\mathfrak{r}^b \neq \overline{\mathfrak{m}}^b(+\infty)$. Then the rescaling argument gives $w_- : cl(B_-) \to \mathbb{CP}^1$ which satisfies the following:

(i) $w_-(\partial B_-) \subset \{\phi = 0, \rho > 0\}$;
(ii) $w_-(\overline{\mathfrak{m}}^b(+\infty)) = 0$ and $w_-(\mathfrak{r}^b) = \infty$;
(iii) $w_-(+\infty) \subset \{\phi = 0, \rho > 0\}$; and
(iv) $w_-|_{int(B_-)}$ is a biholomorphism onto its image.

Here (iii) follows from considering the continuation of w_{-,m_l} to upper levels w_{*,m_l} and π_{*,m_l} (here w_{*,m_l} is the limit of $\overline{u}_{li}$ for $m_l \gg 0$, without taking $m_l \to \infty$) and the fact that $w_{+,m_l} : \Sigma_+ \to \mathbb{CP}^1$ must map $\partial\Sigma_+$ to a thin sector near $\{\phi = 0, \rho > 0\}$.

By the Involution Lemma I.7.10.3 and (i)–(iv), $\mathfrak{r}^b$ must lie on $\mathcal{L}_{1/2}\cup\mathcal{L}_{3/2}$ and the map w_-^{-1} maps the marker $\dot{\mathcal{R}}_\pi(\infty)$ to the marker $\dot{\mathcal{L}}_{1/2}$ or $\dot{\mathcal{L}}_{3/2}$ at $\mathfrak{r}^b$, as appropriate. This *effectively* gives rise to two constraints for $\overline{v}''_-$. Hence $I(\overline{v}''_-)\geq 2$ and the total ECH index is ≥ 4, a contradiction.

Now suppose that $\mathfrak{r}^b=\overline{\mathfrak{m}}^b(+\infty)$. Since passing through $\overline{\mathfrak{m}}^b(+\infty)$ is *effectively* a codimension 2 condition, we have $I(\overline{v}''_-)\geq 2$. The total ECH index is ≥ 4, a contradiction.

(2E) Suppose that $*_0=(-1,j_0)$ for some $1\leq j_0\leq c$. Then (β_1)–(β_3) hold and the rescaling argument with $m_l\to\infty$ gives $w_-:\Sigma_-\to\mathbb{CP}^1$ and $\pi_-:\Sigma_-\to cl(B_-)$ such that:

(i) $w_-(\partial\Sigma_-)\subset\{\phi=0,\rho>0\}$;

(ii) $w_-(z_0)=0$ for some $z_0\in\pi_-^{-1}(\overline{\mathfrak{m}}^b(+\infty))$;

(iii) $w_-(z_1)=\infty$ for some $z_1\in\pi_-^{-1}(-\infty)$;

(iv) $w_-(\pi_-^{-1}(+\infty))\subset\{\phi=0,\rho>0\}$;

(v) $w_-|_{int(\Sigma_-)}$ is a biholomorphism onto its image.

Here (iv) follows from the considerations of Lemma 3.7.8. By the Involution Lemmas, $\pi_-\circ w_-^{-1}$, where defined, maps the component of $\{\mathrm{Im}(z)=0\}$ passing through 0 to $\mathcal{L}_{3/2}$. This contradicts (iii).

(3) Suppose there is a boundary puncture $\mathfrak{r}\in\partial F_{*_0}$ which is removable at z_∞. Then the neighborhood of $\overline{v}''_{*_0}(\mathfrak{r})$ contributes at least k_0 towards n^* by Lemma 3.4.5. We remark that a priori there could be more than one boundary puncture.

(3A) Suppose that $\overline{v}'_-=\varnothing$. Then there is a point $\mathfrak{q}\in int(F_-)$ and a sufficiently small neighborhood $N(\mathfrak{q})\subset\dot{F}_-$ of $\mathfrak{q}$ such that $\overline{v}_-(\mathfrak{q})=\overline{\mathfrak{m}}(+\infty)$ and $n^*(\overline{v}_-(N(\mathfrak{q})))\geq m$. This is a contradiction.

(3B) Suppose that $\overline{v}'_-\neq\varnothing$ and there exists $*_0$ such that $\overline{v}'_{*_0}=\varnothing$, i.e., $\mathfrak{r}$ is of type (P$_1$). Then there exist restrictions $\widetilde{u}_{i,-}$ and $\widetilde{u}_{i,*_0}$ of $\overline{u}_i$ such that $\widetilde{u}_{i,-}$ is close to $\overline{v}'_-$ after translation and passes through $\overline{\mathfrak{m}}(\tau_i)$, $\widetilde{u}_{i,*_0}$ is close to $\overline{v}''_{*_0}$ after translation and nontrivially intersects $\sigma_\infty^{\tau_i}$, and the images of $\widetilde{u}_{i,-}$ and $\widetilde{u}_{i,*_0}$ are disjoint. This is a contradiction since we have an excess of n^*.

(3C) Suppose that $\overline{v}'_-\neq\varnothing$, $\overline{v}'_{*_0}\neq\varnothing$, and $*_0=(1,j)$, $j=0,\dots,a$.

We claim that (a)–(d) from the proof of (1) hold, where we are only considering levels $\overline{v}_*\prec\overline{v}_+$. Indeed, if (a) or (b) does not hold, then the sum of n^* of the ends is $\geq m-2g$ by Lemmas 3.4.1 and 3.7.6, a contradiction. (The situation of $\overline{v}^\sharp_{-1,j}$ is not explicitly covered by Lemma 3.7.6, but the same proof works.) If (c) does not hold, then (a) does not hold, which again gives us a contradiction. (d) is a consequence of (a)–(c). The claim implies that (β_1)–(β_4) hold.

Next we claim that there must be a boundary puncture on $\overline{v}_+$. Arguing by contradiction, a rescaling argument similar to (2B) gives $w_+:\Sigma_+\to\mathbb{CP}^1$ and $\pi_+:\Sigma_+\to cl(B_+)$ such that:

(i) $w_+(\partial\Sigma_+)\subset\{\phi=0,\rho>0\}$;

(ii) $w_+(z_0)=0$ for some $z_0\in\pi_+^{-1}(-\infty)$;

(iii) $w_+(\pi_+^{-1}(-\infty)) \subset \{\phi = 0, \rho > 0\}$;
(iv) $w_+(z_1) = \infty$ for some $z_1 \in \pi_+^{-1}(+\infty)$;
(v) $w_+|_{int(\Sigma_+)}$ is a biholomorphism onto its image.

By the Involution Lemmas, $\pi_+ \circ w_+^{-1}$, where defined, maps the component of $\{\mathrm{Im}(z) = 0\}$ passing through 0 to $\mathcal{L}_{3/2}$, a contradiction. Hence there must be a boundary puncture on $\overline{v}_+$. A similar argument implies that a boundary puncture must lie on $\mathcal{L}_{3/2}$ and must project to a large sector.

Suppose that $\mathfrak{r}$ is of type (P$_2$). Then $\mathfrak{r}$ contributes at least $2p_{*_0} \geq 2$ towards I by Lemma 3.7.5. In addition, the constraint of lying on $\mathcal{L}_{3/2}$ *effectively* contributes $+1$ towards I and the large sector condition contributes $+1$ towards I. These add up to at least 4, a contradiction.

Suppose that $\mathfrak{r}$ is of type (P$_3$). Then $\mathfrak{r}$ contributes at least $+1$ towards I by Lemma 3.7.5 and the total contribution is at least 3, a contradiction. For the purposes of Lemma 3.9.2, we want to improve it by 1. In the current situation we may reduce to:

- ignoring connectors, $a = 0$ and there is no negative end of $\overline{v}^\sharp_{1,j}$ that limits to z_∞;
- $\mathfrak{r}$ is the unique puncture, is of type (P$_3$), is on $\overline{v}''_+$, and corresponds to a large sector;
- $P_{+,0,1}$ corresponds to $\{\overline{a}_{k_1,l_1}, \dots, \overline{a}_{k_s,l_s}\}$, arranged in cyclic order around $\mathbb{R}/2\pi\mathbb{Z}$, and $P_{+,0,0}$ corresponds to $\{\overline{h}(\overline{a}_{k_1,l_1}), \dots, \overline{h}(\overline{a}_{k_s,l_s})\}$;
- $P'_{+,0,1}$ corresponds to $\{\overline{a}_{k_2,l_2}\}$ and $P'_{+,0,0}$ corresponds to $\{\overline{h}(\overline{a}_{k_1,l_1})\}$;
- $P''_{+,0,1}$ corresponds to $\{\overline{a}_{k_1,l_1}, \dots, \overline{a}_{k_s,l_s}\} - \{\overline{a}_{k_2,l_2}\}$ and $P''_{+,0,0}$ corresponds to $\{\overline{h}(\overline{a}_{k_2,l_2}), \dots, \overline{h}(\overline{a}_{k_s,l_s})\}$.

Let $\mathfrak{c}' \subset A_{\varepsilon/2}$ be the grooming corresponding to $\overline{v}'_+$ with $w(\mathfrak{c}') = 0$ and let $\mathfrak{c}'' \subset A_\varepsilon$ be the intersection of the projection of the positive end of $\overline{v}''_+$ with A_ε; one verifies that $w(\mathfrak{c}'') = 0$. The extra contribution to I coming from the positive end of $\overline{v}_+$ is due to minus the writhe of $\mathfrak{c}' \cup \mathfrak{c}''$. Since the writhe is negative, the extra contribution is at least $+1$. Hence the total ECH index is at least 4.

(3D) Suppose that $\overline{v}'_- \neq \varnothing$, $\overline{v}'_{*_0} \neq \varnothing$, and $*_0 = (-1, j)$, $j = 1, \dots, c$.
We claim that we can replace (β_4) by:
(β'_4) $\sum_{j=1}^{c} I(\overline{v}^\sharp_{-1,j}) \geq p_{0,0}$.

Without loss of generality assume that $c = 1$ and $p_{0,0} = 2$. Applying the rescaling argument without taking $m \to \infty$, we obtain $w_- : \Sigma_- \to \mathbb{CP}^1$ and $\pi_- : \Sigma_- \to cl(B_-)$ such that:

(i) $w_-(z_0) = 0$ for some $z_0 \in \pi_-^{-1}(\overline{\mathfrak{m}}^b(+\infty))$;
(ii) w_- maps the negative ends of Σ_- to sectors $\mathfrak{S}(\overline{b}_{k,l}, \overline{h}(\overline{b}_{k',l'}))$.

We now make crucial use of the fact that the sectors $\mathfrak{S}(\overline{b}_{k,l}, \overline{h}(\overline{b}_{k',l'}))$ corresponding to the negative ends of Σ_- have different angles, provided the sectors are not both thin sectors; this follows from the definition of $\overline{\mathbf{a}}$ from Section I.5.2.2. Since each thin sector corresponds to a thin strip of $\overline{v}^\sharp_{-1,1}$ which contributes $I = 1$, let us assume that there are no thin sectors in (ii). We use coordinates $(s, t) \in (-\infty, c] \times$

$[0,1]$ for the two negative ends $\mathcal{E}_{-,1}$, $\mathcal{E}_{-,2}$ of Σ_- and write $w_-^i := w_-|_{\mathcal{E}_{-,i}}$. Then $w_-^i(s,t) \approx c_i e^{\lambda_i(s+it)}$, where $\lambda_1 > \lambda_2 > 0$ without loss of generality; $c_1, c_2 \neq 0$; and the approximation gets better as $s \to -\infty$. On the other hand, suppose that $\operatorname{ind}(\overline{v}^\sharp_{-1,1}) = 1$. Let $\mathcal{E}_{+,i}$, $i = 1,2$, be the positive ends of $\overline{v}^\sharp_{-1,1}$ that limit to z_∞. Here $\mathcal{E}_{+,i}$ is to be paired with $\mathcal{E}_{-,i}$. We use coordinates $(s',t') \in [c',\infty) \times [0,1]$ for the positive ends $\mathcal{E}_{+,i}$. Then $\mathcal{E}_{+,1}$ approaches z_∞ much faster than $\mathcal{E}_{+,2}$ does, as $s' \to +\infty$. This is inconsistent with our description of w_-, when we change coordinates $s' = s + c'$, where $c' \gg 0$. Hence each positive end of $\overline{v}^\sharp_{-1,1}$ that limits to z_∞ contributes $+1$ towards I.

The rest of the argument is similar to that of (3C). □

3.7.6. *The case $\overline{v}'_- \neq \varnothing$.*

Lemma 3.7.11. *If $m \gg 0$, $\overline{u}_\infty \in \partial_{\{+\infty\}}\mathcal{M}$, and $\overline{v}'_- \neq \varnothing$, then no negative end of $\overline{v}^\sharp_{0,j}$, $1 \leq j \leq b$, is asymptotic to a multiple of δ_0.*

Proof. Arguing by contradiction, suppose there exists some $\overline{v}^\sharp_{0,j_0}$, $1 \leq j_0 \leq b$, which has a negative end asymptotic to a multiple of δ_0, say $\delta_0^{q_{0,j_0}}$. Since the negative end of $\overline{v}^\sharp_{0,j_0}$ contributes at least $m - q_{0,j_0}$ towards n^*, the following hold:

(i) at most one $\overline{v}^\sharp_{0,j}$, $j > 0$, has a negative end asymptotic to a multiple of δ_0;
(ii) only one negative end of $\overline{v}^\sharp_{0,j_0}$ limits to a multiple of δ_0;
(iii) $p_{0,0} \leq p_{0,1} \leq \dots \leq p_{0,j_0-1} \leq p_{0,j_0} + q_{0,j_0}$ and $p_{0,j_0} \leq p_{0,j_0+1} \leq \dots \leq p_{0,b+1}$;
(iv) if $\mathcal{E}$ is a positive end of $\overline{v}^\sharp_{0,j}$, $j \geq 0$, that limits to δ_0^r, then $n^*(\mathcal{E}) = r$; and
(v) $\cup_{j=1}^c \overline{v}^\sharp_{-1,j}$ is a union of $p_{0,0} > 0$ thin strips.

By Lemma 3.7.5, all the components besides $\overline{v}'_+$ have nonnegative ECH index. We have contributions (β_1), (β_3), (β_4), (β_6) as in Lemma 3.7.10, as well as the following variant of (β_2):

(β_2') For $1 \leq j < j_0$, $\overline{v}^\sharp_{0,j}$ is a union of cylinders from δ_0 to h or e and

$$\sum_{j=1}^{j_0-1} I(\overline{v}^\sharp_{0,j}) \geq p_{0,j_0} + q_{0,j_0} - p_{0,1}.$$

(β_2'') $\overline{v}^\sharp_{0,j} = \varnothing$ for $j_0 + 1 \leq j \leq b$ and $p_{0,j_0+1} = \dots = p_{0,b+1}$.
(β_2''') $I(\overline{v}^\sharp_{0,j_0}) \geq 2(p_{0,j_0+1} - p_{0,j_0}) + 1$.

We prove (β_2''): First note that $n^*(\overline{v}^\sharp_{0,j_0}) \geq m - q_{0,j_0}$. This implies that $\overline{v}^\sharp_{0,j}$, $j_0 + 1 \leq j \leq b$, must be a union of cylinders from δ_0 to h or e. On the other hand, such cylinders are eliminated by Lemma 3.7.8. Similarly, (β_2''') follows from the proof of Lemma 3.7.8, which implies that each positive end of $\overline{v}^\sharp_{0,j_0}$ that limits to δ_0 additionally contributes two constraints. (β_2') is clear.

Since $\overline{v}_{0,j_0}$ does not satisfy the partition condition at $\delta_0^{p_{0,j_0}+q_{0,j_0}}$, a straightforward calculation which takes into account the braiding near $\delta_0^{p_{0,j_0}+q_{0,j_0}}$ gives:

$$I(\overline{v}_{0,j_0}) \geq I(\overline{v}'_{0,j_0}) + I(\overline{v}''_{0,j_0}) + 2p_{0,j_0}. \tag{3.7.6}$$

Summing the contributions of (β_1), (β'_2)–(β'''_2), (β_3), (β_4), and Equation (3.7.6), we obtain:

$$\sum_{\overline{v}_*} I(\overline{v}_*) \geq q_{0,j_0} + p_{0,j_0+1} + p_{0,j_0} + 1. \tag{3.7.7}$$

Here $q_{0,j_0} \geq 1$ by assumption. If $p_{0,j_0} > 0$ or $p_{0,j_0+1} > 0$, then $\sum_{\overline{v}_*} I(\overline{v}_*) \geq 3$, a contradiction. On the other hand, if $p_{0,j_0} = \cdots = p_{0,b+1} = 0$, then $I(\overline{v}^\sharp_{0,j_0}) \geq 2$ by the usual rescaling argument (cf. Cases (3)–(6) of Theorem I.7.11.1). In this case the right-hand side of Equation (3.7.7) can be increased by one and $\sum_{\overline{v}_*} I(\overline{v}_*) \geq 3$, a contradiction. ☐

Lemma 3.7.12. *If $m \gg 0$, $\overline{u}_\infty \in \partial_{\{+\infty\}}\mathcal{M}$, $\overline{v}'_- \neq \varnothing$, and $\overline{v}^\sharp_+$ has a negative end asymptotic to a multiple of δ_0, then $\overline{u}_\infty \in A'_2$.*

Here A'_2 is as given in Step 4 of the proof of Theorem 3.3.1.

Proof. This is similar to Lemma 3.7.12. The contributions of (β_1), (β'_2)–(β''_2), (β_3), (β_4), and Equation (3.7.6) add up to $q_+ + p_+$.

If $p_+ > 0$, then $q_+ + p_+ \geq 2$. By the usual rescaling argument, the negative end of $\overline{v}^\sharp_+$ that limits to δ_0 additionally contributes $+1$ to I. Hence $\sum_{\overline{v}_*} I(\overline{v}_*) \geq 3$, a contradiction.

If $p_+ = 0$, then $q_+ + p_+ \geq 1$. The negative end of $\overline{v}^\sharp_+$ that limits to δ_0 again contributes $+1$ to I. Hence if $\sum_{\overline{v}_*} I(\overline{v}_*) = 2$, then $q_+ = 1$. This implies that $\overline{u}_\infty \in A'_2$. ☐

Lemma 3.7.13. *If $m \gg 0$, $\overline{u}_\infty \in \partial_{\{+\infty\}}\mathcal{M}$, and $\overline{v}'_- \neq \varnothing$, then $\overline{u}_\infty \in A'_2$.*

Proof. Consider $\overline{u}_\infty \in \partial_{\{+\infty\}}\mathcal{M}$ such that $\overline{v}'_- \neq \varnothing$. By Lemma 3.7.10, there is no boundary puncture which is removable at z_∞, no fiber component and no $\overline{v}''_*$ such that $\overline{v}'_* \cap \overline{v}''_* \neq \varnothing$. By Lemmas 3.7.11 and 3.7.12, if $\overline{v}^\sharp_{0,j_0}$, $1 \leq j_0 \leq b+1$, has a negative end asymptotic to a multiple of δ_0, then $\overline{u}_\infty \in A'_2$.

It remains to consider three possibilities, all of which satisfy $p_{0,0} \leq \cdots \leq p_{0,b+1}$:

(1) Some $\overline{v}^\sharp_{-1,j_0}$, $j_0 = 1, \dots, c+1$, has a positive or negative end $\mathcal{E}$ at z_∞ such that $n^*(\mathcal{E}) > m/2$.
(2) Some $\overline{v}^\sharp_{1,j_0}$, $j_0 = 0, \dots, a$, has a positive or negative end $\mathcal{E}$ at z_∞ such that $n^*(\mathcal{E}) > m/2$.
(3) Some $\overline{v}^\sharp_{0,j_0}$, $j_0 = 0, \dots, b$, has a positive end $\mathcal{E}$ at a multiple of δ_0 such that $n^*(\mathcal{E}) > m/2$.

It is clear that (1), (2), and (3) are mutually exclusive.

(1) is similar to (2E) of Lemma 3.7.10 and (2) is similar to Case (6_i) of Lemma 3.4.24.

(3) In this case, (a)–(c) in the proof of Lemma 3.7.10 hold, with the exception of the level $\overline{v}_{0,j_0}$. Observe that $\overline{v}'_{0,j_0} \neq \varnothing$ and $\overline{v}^\sharp_{0,j_0}$ consists of:

- c_1 cylinders from δ_0 to e or h which contribute $I = 1$ or 2 each; and
- a curve $\overline{v}^{\sharp_2}_{0,j_0}$ with one positive end $\mathcal{E}$ which is asymptotic to $\delta_0^{c_2}$ and satisfies $n^*(\mathcal{E}) = m + c_2$ and a union $\mathcal{E}'$ of other positive ends which in total is asymptotic to $\delta_0^{c_3}$ and satisfies $n^*(\mathcal{E}') = c_3$.

Without loss of generality assume that there are no positive ends of $\overline{v}^\sharp_{1,j}$ for $j > j_0$ and that all the branch points of $\overline{v}'_{1,j}$ for $j > j_0$ are in $\overline{v}'_{1,j_0+1}$. We apply the rescaling argument with $m \to \infty$ to obtain $w_{0,j_0+1} : \Sigma_{0,j_0+1} \to \mathbb{CP}^1$ and $\pi_{0,j_0+1} : \Sigma_{0,j_0+1} \to cl(B_{0,j_0+1})$ with $B_{0,j_0+1} \simeq B'$ and Σ_{0,j_0+1} connected, such that the following hold:

(i) $w_{0,j_0+1}(z_0) = 0$ for some $z_0 \in \pi^{-1}_{0,j_0+1}(-\infty)$;
(ii) $w_{0,j_0+1}(z_1) = \infty$ for some $z_1 \neq z_0 \in \pi^{-1}_{0,j_0+1}(-\infty)$;
(iii) w_{0,j_0+1} is a biholomorphism; and
(iv) in particular, $w_{0,j_0+1}(z) \neq 0, \infty$ for $z \in \pi^{-1}_{0,j_0+1}(\pm\infty) - \{z_0, z_1\}$.

An argument similar to that of (3D) in the proof of Lemma 3.7.10 implies that $c_3 = 0$, since otherwise (ii) and (iv) contradict the relative asymptotic decay rates of $\mathcal{E}$ and $\mathcal{E}'$ (i.e., $\mathcal{E}$ approaches δ_0 much faster as $s \to \infty$).

We now compute the ECH index $I(\overline{v}^{\sharp_2}_{0,j_0})$. We use the formula:

$$\operatorname{ind}(\overline{u}) + (\widetilde{\mu}_\tau(\overline{u}) - \mu_\tau(\overline{u}) - w_\tau(\overline{u})) \leq I(\overline{u}), \tag{3.7.8}$$

where the notation is as in Section I.3.4 and the equation follows from the adjunction inequality and [Hu2, Lemma 4.20]. The contributions towards $\widetilde{\mu}_\tau(\overline{v}^{\sharp_2}_{0,j_0})$, $\mu_\tau(\overline{v}^{\sharp_2}_{0,j_0})$, and $w_\tau(\overline{v}^{\sharp_2}_{0,j_0})$ coming from the positive end $\delta_0^{c_2}$ are c_2, 1, and $1 - c_2$, respectively. Moreover, $\operatorname{ind}(\overline{v}^{\sharp_2}_{0,j_0}) \geq 3$ since the vanishing of the leading asymptotic eigenfunction corresponding to the end $\mathcal{E}$ is a codimension two condition. Hence,

$$I(\overline{v}^{\sharp_2}_{0,j_0}) \geq 3 + (c_2 - 1 - (1 - c_2)) = 1 + 2c_2.$$

Next, since $p_{0,j_0} = \deg \overline{v}'_{0,j_0} > 0$, we have

$$\begin{aligned} I(\overline{v}'_{0,j_0} \cup \overline{v}^{\sharp_2}_{0,j_0}) &\geq I(\overline{v}'_{0,j_0}) + I(\overline{v}^{\sharp_2}_{0,j_0}) + 2p_{0,j_0} \\ &\geq 0 + (1 + 2c_2) + 2p_{0,j_0} \geq 5. \end{aligned}$$

Hence the total ECH index of all the levels is ≥ 5, a contradiction. □

3.7.7. *The case $\overline{v}'_- = \varnothing$.*

Lemma 3.7.14. *If $m \gg 0$, then there is no $\overline{u}_\infty \in \partial_{\{+\infty\}}\mathcal{M}$ such that $\overline{v}'_- = \varnothing$.*

Proof. Arguing by contradiction, suppose there exists $\overline{u}_\infty \in \partial_{\{+\infty\}}\mathcal{M}$ such that $\overline{v}'_- = \varnothing$. By Lemma 3.7.10, there is no boundary puncture which is removable at z_∞, no fiber component, and no $\overline{v}''_*$ such that $\overline{v}'_* \cap \overline{v}''_* \neq \varnothing$.

The point constraint gives $n^*(\overline{v}''_-) \geq m$ and $I(\overline{v}''_-) \geq 2$. This immediately implies that (a)–(d) in the proof of Lemma 3.7.10 hold. In particular, (β_2) from Lemma 3.7.10 holds.

If $p_{0,b+1} > 0$, then the proof of Lemma 3.7.8 gives us a contradiction. Hence $p_{0,b+1} = 0$. If a positive end of $\overline{v}''_+$ limits to z_∞, then a component $\widetilde{v}$ of $\overline{v}''_+$ must be a section of $\overline{W}_+$ from z_∞ to h or e. However, by the choice of h from Section 3.1.4 and the proof of Lemma I.6.6.5, sections of $\overline{W}_+$ from z_∞ to h cannot exist, leaving us with the possibility that $\widetilde{v}$ is a section from z_∞ to e. Hence $I(\widetilde{v}) \geq 1$ and the total ECH index is ≥ 3, a contradiction. On the other hand, if $\overline{v}''_+ = \overline{v}^\flat_+$, then $I(\overline{v}_{1,j}) \geq 1$ for some $j > 0$, which is again a contradiction. □

Proof of Lemma 3.3.10. Lemma 3.3.10 follows from Lemmas 3.7.13 and 3.7.14. □

3.8. **Degeneration at $-\infty$, part II.** Let $\overline{u}_\infty \in \partial_{\{-\infty\}}\mathcal{M}$ be the limit of $\overline{u}_i \in \mathcal{M}_{\tau_i}$, where $\tau_i \to -\infty$.

Lemma 3.8.1. *If fiber components are removed from $\overline{u}_\infty$ and $\overline{v}'_1 = \varnothing$, then the ECH index of each level is nonnegative.*

Proof. The proof is similar to that of Lemma 3.5.2 and uses the considerations of Lemma 3.7.5. □

3.8.1. *The case $\overline{v}'_2 = \varnothing$.*

Lemma 3.8.2. *If $\overline{u}_\infty \in \partial_{\{-\infty\}}\mathcal{M}$ and $\overline{v}'_2 = \varnothing$, then:*

1. *$\overline{u}_\infty$ is a 2-level building consisting of $\overline{v}_1$ with $I(\overline{v}_1) = 0$, $\overline{v}_2$ with $I(\overline{v}_2) = 2$, and $\overline{v}'_j = \varnothing$ for $j = 1, 2$;*
2. *the left and right ends of $\overline{v}_1$ (= left and right ends of $\overline{v}_2$) do not limit to z_∞;*
3. *$\overline{u}_\infty$ has no fiber component;*
4. *$\overline{u}_\infty$ has no boundary puncture which is removable at z_∞.*

Proof. Suppose that $\overline{v}'_2 = \varnothing$. First observe that

$$n^*(\overline{v}_2(N(\mathfrak{q}))) \geq m, \tag{3.8.1}$$

where $\mathfrak{q} \in \dot{F}_2$ is the point such that $\overline{v}_2(\mathfrak{q}) = \overline{\mathfrak{m}}(-\infty)$.

We also have $I(\overline{v}_2) \geq 2$: If $\overline{u}_\infty$ has a fiber component $\widetilde{v}$, then $\widetilde{v}$ must pass through $\overline{\mathfrak{m}}(-\infty)$ and $I(\overline{v}_2) \geq 2g + 2 \geq 4$ by an argument similar to that of Lemma I.7.6.5. On the other hand, if $\overline{u}_\infty$ does not have a fiber component, then $I(\overline{v}_2) \geq 2$, since passing through $\overline{\mathfrak{m}}(-\infty)$ is a generic condition.

(1) We claim that $\overline{v}'_1 = \varnothing$. Arguing by contradiction, if $\overline{v}'_1 \neq \varnothing$, then at least one of $\overline{v}^\sharp_{L,j}$, $j = 0, \dots, a$, has a right end $\mathcal{E}_R$ that limits to z_∞. On the other hand, the end $\mathcal{E}_R$ satisfies

$$n^*(\mathcal{E}_R) \geq k_0 - 1 \gg 2g, \tag{3.8.2}$$

which contradicts Equation (3.8.1). This proves the claim.

The claim and Lemma 3.8.1 imply that each level of $\overline{u}_\infty$ has nonnegative ECH index. Hence $I(\overline{v}_2) = 2$, $I(\overline{v}_1) = 0$, and $a = b = c = d = 0$ since a level $\overline{v}_{L,j}$, $j = 1, \dots, a$, is not a connector if and only if $I(\overline{v}_{L,j}) > 0$ (and the same holds for $\overline{v}_{R,j}$, $j = 1, \dots, b$, $\overline{v}_{B,j}$, $j = 1, \dots, c$, and $\overline{v}_{T,j}$, $j = 1, \dots, d$).

(2) If $\overline{v}_2$ has a right end $\mathcal{E}_R$ that limits to z_∞, then $\mathcal{E}_R$ satisfies Equation (3.8.2), and we have a contradiction of Equation (3.8.1). If $\overline{v}_2$ has a left end that limits to z_∞, then $\overline{v}_1$ has a right end $\mathcal{E}_R$ that limits to z_∞ and satisfies Equation (3.8.2) (since $\overline{v}'_1=\varnothing$), which is again a contradiction.

(3) Since $I(\overline{v}_2)=2$, we cannot have a fiber component.

(4) A boundary puncture which is removable at z_∞ contradicts Equation (3.8.1). □

3.8.2. *The case* $\overline{v}'_1=\varnothing, \overline{v}'_2\neq\varnothing$.

Lemma 3.8.3. *If $m\gg 0$, then there is no $\overline{u}_\infty\in\partial_{\{-\infty\}}\mathcal{M}$ such that $\overline{v}'_1=\varnothing$ and $\overline{v}'_2\neq\varnothing$.*

Proof. Arguing by contradiction, suppose there exists $\overline{u}_\infty\in\partial_{\{-\infty\}}\mathcal{M}$ such that $\overline{v}'_1=\varnothing$ and $\overline{v}'_2\neq\varnothing$. The analog of Lemma 3.5.1 holds, i.e.,

$$\sum_{i=1}^{q} n^*(\mathcal{E}_i)+\sum_{i=1}^{r} n^*(\mathcal{E}'_i)=m,\quad \sum_{i=1}^{q} n^{*,alt}(\mathcal{E}_i)+\sum_{i=1}^{r} n^{*,alt}(\mathcal{E}'_i)\geq m-2g,$$

where $\mathcal{E}_i$, $i=1,\dots,q$, are the ends that limit to z_∞ and $\mathcal{E}'_i$, $i=1,\dots,r$, are the neighborhoods of punctures of type (P$_3$) of all the $\overline{v}^\sharp_{L,j}$, $j=0,\dots,a$, and $\overline{v}^\sharp_{R,j}$, $j=0,\dots,b$. The conclusion of Lemma 3.5.4 then holds by an almost identical argument. The proof of Lemma 3.5.14 then carries over without modification. □

3.8.3. *The case* $\overline{v}'_1\neq\varnothing, \overline{v}'_2\neq\varnothing$.

Lemma 3.8.4. *Suppose $\overline{v}'_1\neq\varnothing$ and $\overline{v}'_2\neq\varnothing$. Let $\mathcal{E}_i$, $i=1,\dots,q$, be the ends of all the $\overline{v}^\sharp_*$ that limit to z_∞ and let $\mathcal{E}'_i$, $i=1,\dots,r$, be the neighborhoods of the punctures of type (P$_3$). Then one of the following holds:*

(a) $\sum_{i=1}^{q} n^*(\mathcal{E}_i)+\sum_{i=1}^{r} n^*(\mathcal{E}'_i)\geq m-2g$.

(b) *Each end $\mathcal{E}_i$ is an end of $\overline{v}^\sharp_{B,j}$, $j=1,\dots,c$, or $\overline{v}^\sharp_{T,j}$, $j=0,\dots,d$, that projects to a thin sector of type $\mathfrak{S}(\overline{a}_{i',j'},\overline{h}(\overline{a}_{i',j'}))$ or $\mathfrak{S}(\overline{b}_{i',j'},\overline{h}(\overline{b}_{i',j'}))$. In particular, $\mathcal{E}_i$ is not a left or right end of $\overline{v}^\sharp_*$, where $*=(L,j)$, $j=0,\dots,a$, or (R,j), $j=0,\dots,b$.*

Proof. We apply the continuation argument. If some $\mathcal{E}_i$, $i=1,\dots,q$, does not project to a thin sector (of type $\mathfrak{S}(\overline{a}_{i',j'},\overline{h}(\overline{a}_{i',j'}))$, $\mathfrak{S}(\overline{b}_{i',j'},\overline{h}(\overline{b}_{i',j'}))$, or $\mathfrak{S}(\overline{b}_{i',j'},\overline{a}_{i',j'})$), then the sectors $\pi_{D_{\rho_0}}(\mathcal{E}_i)$, $i=1,\dots,q$, and $\pi_{D_{\rho_0}}(\mathcal{E}'_i)$, $i=1,\dots,r$, will sweep out a neighborhood of z_∞ with the exception of some thin sectors, implying (a).

On the other hand, suppose that all the ends $\mathcal{E}_i$, $i=1,\dots,r$, project to thin sectors. We claim that there are no thin sectors of type $\mathfrak{S}(\overline{b}_{i',j'},\overline{a}_{i',j'})$. Indeed, the number of left and right ends of all the $\overline{v}^\sharp_*$ that limit to z_∞ must be equal and the right ends cannot map to thin sectors. This gives (b). □

Lemma 3.8.5. *If $\overline{u}_\infty\in\partial_{\{-\infty\}}\mathcal{M}$, $\overline{v}'_1\neq\varnothing$, and $\overline{v}'_2\neq\varnothing$, then:*

(1) *$\overline{u}_\infty$ has no fiber components;*

(2) *if $\overline{v}''_*$ satisfies $\overline{v}'_*\cap\overline{v}''_*\neq\varnothing$ and $\overline{v}'_*\cap\overline{v}''_*\subset int(\overline{v}'_*)$, then $*=1$ and $p_*=1$;*

(3) *$\overline{u}_\infty$ has no boundary puncture of type (P$_1$) or (P$_2$).*

Proof. This is similar to the proof of Lemma 3.7.10 and uses Equations (3.7.2) and (3.7.1).

First observe that if $\overline{u}_\infty$ has a fiber component, a boundary puncture of type (P$_1$) or (P$_2$), or some $\overline{v}'_*\cap\overline{v}''_*\neq\varnothing$ with $\overline{v}'_*\cap\overline{v}''_*\subset int(\overline{v}'_*)$, then we are in the situation of Lemma 3.8.4(b). It is not hard to verify that $I(\overline{v}_*)\geq 0$ for all $*$ with the exception of $\overline{v}_1$.

(1) Suppose $\overline{u}_\infty$ has a fiber component $\widetilde{v}$. Then we have the following contributions towards I:

(β_1) $I(\overline{v}'_1)=-p_1=-\deg(\overline{v}'_1)$.
(β_2) $\sum_{j=1}^{c} I(\overline{v}^{\sharp}_{B,j})=p_1$.
(β_3) If the fiber $\widetilde{v}$ is a component of $\overline{v}_*$, then $\widetilde{v}$ and the intersection $\widetilde{v}\cap(\overline{v}_*-\widetilde{v})$ contribute $2g+2\geq 4$ towards I.

The argument is similar to that of Lemma 3.7.10(1). This gives a total of $I>2$, which is a contradiction.

(2) Suppose $\overline{v}'_{*_0}\cap\overline{v}''_{*_0}\neq\varnothing$ and $\overline{v}'_{*_0}\cap\overline{v}''_{*_0}\subset int(\overline{v}'_{*_0})$. If $*_0\neq 1$, then $I(\overline{v}''_{*_0})\geq 1$ and the intersection points contribute at least $2p_{*_0}\geq 2$. If $*_0=1$ and $p_{*_0}>1$, then $I(\overline{v}''_{*_0})\geq 0$ and the intersection points contribute at least $2p_{*_0}\geq 4$. In view of the ECH contributions from (β_1) and (β_2), we have a contradiction.

(3) If $\overline{u}_\infty$ has a boundary puncture of type (P$_1$), then argument from (3B) of Lemma 3.7.10 implies an excess of n^*. If $\overline{u}_\infty$ has a boundary puncture of type (P$_2$), then the analog of Lemma 3.7.5 implies that the boundary punctures contribute $2p_{*_0}\geq 2$; there is also a large sector which contributes an additional $+1$. This gives us a total of $I>2$, a contradiction. □

Lemma 3.8.6. *If $m\gg 0$, $\overline{u}_\infty\in\partial_{\{-\infty\}}\mathcal{M}$, $\overline{v}'_1\neq\varnothing$, and $\overline{v}'_2\neq\varnothing$, then there is no level $\overline{v}_*$ such that $\overline{v}'_*\cap\overline{v}''_*\neq\varnothing$ and $\overline{v}'_*\cap\overline{v}''_*\subset int(\overline{v}'_*)$.*

Proof. Arguing by contradiction, suppose that $\overline{v}'_{*_0}\cap\overline{v}''_{*_0}\neq\varnothing$ and $\overline{v}'_{*_0}\cap\overline{v}''_{*_0}\subset int(\overline{v}'_{*_0})$ for some $\overline{v}_{*_0}$. Then $*_0=1$ and $p_1=1$ by Lemma 3.8.5(2). By Lemma 3.8.4(b), we have $p_{L,j}=1$ and $p_{R,j}=1$ for all j. Hence we may assume that $a=b=0$. Equation (3.7.2) then implies that $c=1$ and $d=0$.

By the rescaling argument with $m\to\infty$, we obtain a 2-level building $w_1\cup w_2$, where

$$w_2: cl(B_{-\infty,2})\to\mathbb{CP}^1,\quad w_1: cl(B_{-\infty,1})\to\mathbb{CP}^1,$$

and (i)–(viii), given below, hold. Here $cl(B_{-\infty,2})$ is obtained from $B_{-\infty,2}$ by adding the left and right points at infinity, denoted $t=\pm\infty$; similarly $cl(B_{-\infty,1})$ is obtained from $B_{-\infty,1}$ by adding $s=\pm\infty$ and $t=\pm\infty$. The map w_2 satisfies the following:

(i) $w_2(\overline{\mathfrak{m}}^b(-\infty))=0$;
(ii) $w_2(\partial cl(B_{-\infty,2}))\subset\{\phi=0,\rho>0\}\cup\{\infty\}$;
(iii) $w_2(t=+\infty)=\infty$ and $w_0(t=-\infty)\subset\{\phi=0,\rho>0\}$ (or vice versa);
(iv) $w_2|_{int(B_{-\infty,2})}$ is a biholomorphism onto its image.

The map w_1 satisfies the following:

(v) $w_1(t=-\infty)=0$;

(vi) $w_1(\partial cl(B_{-\infty,1})) \subset \{\phi = 0, \rho > 0\} \cup \{\infty\}$;
(vii) $w_1(\mathfrak{r}) = \infty$, where $\mathfrak{r} \in \overline{v}_1' \cap \overline{v}_1''$;
(viii) $w_1|_{int(B_{-\infty,1}}$ is a biholomorphism onto its image.

In view of the above, we consider the maps

$$\overline{\Xi}_i : \Gamma \to ([-1,1]/(-1 \sim 1)) \times [0,\infty)$$

as in Section 3.5.6. The proof strategy of Lemmas 3.5.14 and 3.5.15 carry over to give us a contradiction; this is due to the disparity in the growth rates of the left and right ends of $\overline{u}_i$ when restricted to $[-2,2] \times [1+L, r(\tau_i) - L]$. ☐

The combination of Lemmas 3.8.5 and 3.8.6 give the following, which is the analog of Lemma 3.5.4(2),(3):

Corollary 3.8.7. *If $m \gg 0$, $\overline{u}_\infty \in \partial_{\{-\infty\}}\mathcal{M}$, $\overline{v}_1' \neq \varnothing$, and $\overline{v}_2' \neq \varnothing$, then there is no fiber component, no boundary puncture of type (P$_1$) or (P$_2$), and no intersection point $\mathfrak{r} \in \overline{v}_*' \cap \overline{v}_*''$ such that $\mathfrak{r} \in \overline{v}_*' \cap \overline{v}_*'' \subset int(\overline{v}_*')$.*

Before embarking on the proof of Lemma 3.8.11, we prove Lemma 3.8.8, which is the analog of Lemma 3.5.4(4) and is a bit involved.

First we recall and modify some notation from Section I.5.7. We use the convention that $\star = (L,j)$, $j = 1,\dots,a$, or (R,j), $j = 1,\dots,b+1$. The data $\overrightarrow{\mathcal{D}}_{\star,\pm}$ at z_∞ for $\overline{v}_\star$ is given by a $p = p_{\star,\pm}$-tuple of matchings

$$\{(i_1', j_1') \to (i_1, j_1), \dots, (i_p', j_p') \to (i_p, j_p)\}, \tag{3.8.3}$$

where $i_k, i_k' \in \{1,\dots,2g\}$, $j_k, j_k' \in \{0,1\}$ for $k = 1,\dots,p$ and $i_k \neq i_l$, $i_k' \neq i_l'$ for $k \neq l$. Here the subscript $+$ (resp. $-$) in $\overrightarrow{\mathcal{D}}_{\star,\pm}$ refers to the left (resp. right) end, (i_k', j_k') corresponds to $\overline{b}_{i_k', j_k'}$, and (i_k, j_k) corresponds to $\overline{a}_{i_k, j_k}$. We write

$$\overrightarrow{\mathcal{D}}_{\star,\pm} = \overrightarrow{\mathcal{D}}'_{\star,\pm} \cup \overrightarrow{\mathcal{D}}''_{\star,\pm},$$

where $\overrightarrow{\mathcal{D}}'_{\star,\pm}$ and $\overrightarrow{\mathcal{D}}''_{\star,\pm}$ correspond to the ends of $\overline{v}_\star'$, $\overline{v}_\star''$, respectively.

Lemma 3.8.8. *If $\overline{u}_\infty \in \partial_{\{-\infty\}}\mathcal{M}$, $\overline{v}_1' \neq \varnothing$ and $\overline{v}_2' \neq \varnothing$, then every component of $\overline{v}_\star^\sharp$, $\star = (L,j)$, $j = 1,\dots,a$, or (R,j), $j = 1,\dots,b+1$, is a thin strip.*

Proof. Arguing by contradiction, suppose there exists $\overline{v}_{\star_0}^\sharp$ which is not a union of thin strips; we take $\star_0 = (R,1)$ without loss of generality. Then the following hold:

(a) $n^*(\overline{v}_{\star_0}^\sharp) = m$ and $n^{*,alt}(\overline{v}_{\star_0}^\sharp) \geq m - 2g$;
(b) each component $\widetilde{v}$ of $\overline{v}_{T,j}^\sharp$, $j = 1,\dots,d$, or $\overline{v}_{B,j}^\sharp$, $j = 1,\dots,c$, is a thin strip with z_∞ at the positive end and $I(\widetilde{v}) = 1$;
(c) each component $\widetilde{v}$ of $\overline{v}_\star^\sharp$, $\star \neq \star_0$, is a thin strip with z_∞ at the left end and $I(\widetilde{v}) = 1$;
(d) each component $\widetilde{v}$ of $\overline{v}_1^\sharp$ limits to z_∞ only at the left end and the top end, the left and top ends project to thin sectors, $\deg(\widetilde{v}) = 1$, and $I(\widetilde{v}) = 1$.

(a) follows from the argument of Lemma 3.5.4(4)(a) and (b) and (c) follow immediately from (a). (d) The assertions about the ends of $\widetilde{v}$ follow from (a). This then implies that $\widetilde{v}$ projects to the domain bounded by $\overline{b}_i$, $\overline{h}(\overline{b}_i)$, and $\overline{h}(\overline{a}_i)$ in $\overline{S}$, in view of the positions of the arcs $\overline{b}_{i,j}$, $\overline{a}_{i,j}$, $\overline{h}(\overline{b}_{i,j})$, and $\overline{h}(\overline{a}_{i,j})$ from Figure 6.

We now analyze the ends of $\overline{v}^\sharp_{\star 0}$ at z_∞ in more detail. Let l_+ and l_- be the number of left and right ends of $\overline{v}^\sharp_{\star 0}$ at z_∞ and let $\mathfrak{bp}$ be the number of boundary punctures of type (P$_3$) on $\overline{v}_{\star 0}$. In view of (c) and (d), the total number of left ends of $\overline{v}_*$ that limit to z_∞ is $l_- - l_+$. By (a slight modification of) the continuation method, we obtain a cycle $\mathcal{Z}$ consisting of $l_+ + l_- + \mathfrak{bp}$ chords, where l_+ of the chords are of type $\mathfrak{S}(\overline{a}_{k,l}, \overline{b}_{k',l'})$, l_- of the chords are of type $\mathfrak{S}(\overline{b}_{k,l}, \overline{a}_{k',l'})$, $\mathfrak{bp}$ of the chords are of type $\mathfrak{S}(\overline{a}_{k,l}, \overline{a}_{k',l'})$ or $\mathfrak{S}(\overline{b}_{k,l}, \overline{b}_{k',l'})$, and all of the chords correspond to left or right ends of $\overline{v}^\sharp_{\star 0}$ at z_∞ or boundary punctures of $\overline{v}^\sharp_{\star 0}$ of type (P$_3$).

Let us write $p_* = \deg \overline{v}'_*$, $q_* = \deg \overline{v}^\sharp_*$, and $r_* = \deg \overline{v}^\flat_*$. We have $\operatorname{ind}(\overline{v}^\sharp_{\star 0}) \geq 2$ since there is an end that limits to z_∞ or a boundary puncture of type (P$_3$) that projects to a large sector. Then $I(\overline{v}^\sharp_{\star 0}) \geq 2$ by the ECH index inequality and considerations of $\mathcal{Z}$.

(1) Suppose that $\overline{v}'_{\star 0} \neq \varnothing$. We claim the following:

Claim 3.8.9. $I(\overline{v}_{\star 0}) \geq I(\overline{v}'_{\star 0}) + I(\overline{v}''_{\star 0}) + c$, *where* $c = \mathfrak{bp}$ *if* $\mathfrak{bp} > 0$ *and* $c = 2p_{\star 0}$ *if* $\mathfrak{bp} = 0$.

Proof of Claim 3.8.9. Each boundary puncture of type (P$_3$) contributes $+1$ towards I by the argument from Lemma 3.7.5, for a total contribution of $\mathfrak{bp}$. Suppose for simplicity that $\mathfrak{bp} = 0$. Then $l_+ = l_-$. Let (s, t_2) be coordinates on $[-2,2] \times [-1,1]$. Let

$$\check{C}'_{[-1,1]}, \check{C}''_{[-1,1]} \subset [-2,2] \times [-1,1] \times \overline{S}$$

be representatives of $\overline{v}'_{\star 0}$ and $\overline{v}''_{\star 0}$ and let $\mathfrak{c}'_{\pm 1}$, $\mathfrak{c}''_{\pm 1}$ be groomings on $A_{\varepsilon/2} = \partial D_{\varepsilon/2} \times [-2,2]$ and $A_\varepsilon = \partial D_\varepsilon \times [-2,2]$ corresponding to the positive $(+1)$ and negative (-1) ends of $\check{C}'_{[-1,1]}$ and $\check{C}''_{[-1,1]}$ such that the following hold:

- $\mathfrak{c}'_{+1} = \mathfrak{c}'_{-1}$ and $w(\mathfrak{c}'_{\pm 1}) = 0$;
- $\mathfrak{c}''_{\pm 1}$ is obtained by intersecting $\pi_{[-2,2]\times\overline{S}}$-projections of the positive and negative ends of $\overline{v}''_{\star 0}$ with A_ε.

Here $\pi_{[-2,2]\times\overline{S}}$ is the projection $[-2,2] \times \mathbb{R} \times \overline{S} \to [-2,2] \times \overline{S}$. By the above description of $\mathcal{Z}$:

- $w(\mathfrak{c}''_{+1}) = 0$ or 1 and $w(\mathfrak{c}''_{-1}) = 0$ or -1;
- the endpoints of $\mathfrak{c}''_{+1}$ and $\mathfrak{c}''_{-1}$ agree and are alternating.

We now extend $\check{C}'_{[-1,1]}, \check{C}''_{[-1,1]}$ by concatenating with

$$\check{C}'_{[1,2]}, \check{C}''_{[1,2]} \subset [-2,2] \times [1,2] \times \overline{S},$$
$$\check{C}'_{[-2,-1]}, \check{C}''_{[-2,-1]} \subset [-2,2] \times [-2,-1] \times \overline{S}$$

such that the ends of $\check{C}'_{[-2,2]} \cup \check{C}''_{[-2,2]}$ are groomed and have zero winding number. Writing $\mathfrak{wr}$ for the writhe, one can verify that the only net contribution to I comes from

$$2(-\mathfrak{wr}(\mathfrak{c}'_{+1}\cup\mathfrak{c}''_{+1})+\mathfrak{wr}(\mathfrak{c}'_{-1}\cup\mathfrak{c}''_{-1})),$$

which is equal to $2p_{\star0}$. □

By Claim 3.8.9, $I(\overline{v}_{\star0})\geq 3$. Since the ECH indices of the other levels add up to at least 0 in view of (b), (c), and (d), we have a contradiction.

(2) Suppose that $\overline{v}'_{\star0}=\varnothing$. Then there are no boundary punctures of type (P_3) by definition. Since $I(\overline{v}_{\star0})=I(\overline{v}''_{\star0})\geq 2$, the only nontrivial levels are $\overline{v}_{B,j}$, $j=1,\dots,c$, which consist of thin strips and trivial strips. We claim the following:

Claim 3.8.10. *$I(\overline{v}''_{\star0})\geq 2+q_{\star0}$, where $q_{\star0}=\deg\overline{v}^\sharp_{\star0}$.*

Proof of Claim 3.8.10. Let us first consider the case where:

1. $\pi_{\overline{S}}\circ\overline{v}^\sharp_{\star0}|_{\dot{F}^\sharp_{\star0}}$ is a diffeomorphism onto its image, which contains $\overline{S}-\overline{\mathbf{a}}-\overline{\mathbf{b}}$;
2. $\pi_{\overline{S}}\circ\overline{v}^\sharp_{\star0}$ maps each component of $\partial\dot{F}^\sharp_{\star0}$ to a slit on some $\overline{a}_i$ or $\overline{b}_i$;
3. $q_{\star0}=l_+=l_-$.

If we choose a multivalued trivialization τ to be compatible with $\overrightarrow{\mathcal{D}}''_{\star0,\pm}$ (cf. Section I.5.7.3), then

$$\begin{aligned}\operatorname{ind}(\overline{v}^\sharp_{\star0})&=-\chi(F^\sharp_{\star0})+q_{\star0}+\mu_\tau(\overline{v}^\sharp_{\star0})+2c_1((\overline{v}^\sharp_{\star0})^*T\overline{S},\tau)\\&=-(1-2g)+q_{\star0}+(l_++1)+2(1-2g)\\&=2-2g+2q_{\star0}.\end{aligned}$$

Hence $I(\overline{v}^\sharp_{\star0}))\geq 2-2g+2q_{\star0}$.

Next we consider the general case. Since $I(\overline{v}''_{\star0})$ is a homological quantity, by the description of $\mathcal{Z}$ and the observation that there are no boundary punctures of type (P_3), we may replace $\overline{v}''_{\star0}$ by a homologous map $\overline{v}''_{\star0}=\overline{v}^\sharp_{\star0}\cup\overline{v}^\flat_{\star0}$ where:

1. $\overline{v}^\sharp_{\star0}$ is as in the previous paragraph;
2. $\overline{v}^\flat_{\star0}$ has $r_{\star0}$ components and each is a strip with $\operatorname{ind}=I\geq -1$;
3. $\langle\overline{v}^\sharp_{\star0},\overline{v}^\flat_{\star0}\rangle=r_{\star0}$, where $\langle,\rangle$ is the algebraic intersection number.

Summing the contributions, we obtain:

$$\begin{aligned}I(\overline{v}_{\star0})&=I(\overline{v}^\sharp_{\star0})+I(\overline{v}^\flat_{\star0})+2r_{\star0}\\&\geq 2-2g+2q_{\star0}-r_{\star0}+2r_{\star0}=2+q_{\star0},\end{aligned}$$

since $q_{\star0}+r_{\star0}=2g$. □

Since $q_{\star0}\geq 1$, $I(\overline{v}_{\star0})\geq 3$, which is a contradiction. □

Lemma 3.8.11. *If $m\gg 0$, then there is no $\overline{u}_\infty\in\partial_{\{-\infty\}}\mathcal{M}$ such that $\overline{v}'_1\neq\varnothing$ and $\overline{v}'_2\neq\varnothing$.*

Proof. Corollary 3.8.7 and Lemma 3.8.8 imply the analog of Lemma 3.5.4. The proof of Lemma 3.5.15 then carries over with no change. □

Proof of Lemma 3.3.11. Suppose $\overline{u}_\infty \in \partial_{\{-\infty\}}\mathcal{M}$. If $\overline{v}_2' = \varnothing$, then we are in the situation of Lemma 3.3.11 by Lemma 3.8.2. On the other hand, Lemmas 3.8.3 and 3.8.11 imply that $\overline{v}_2' = \varnothing$. □

3.9. **Breaking in the middle, part II.** Let $\overline{u}_\infty \in \partial_{(-\infty,+\infty)}\mathcal{M}$ be the limit of $\overline{u}_i \in \mathcal{M}_{\tau_i}$, where $\tau_i \to T'$.

Lemma 3.9.1. *If fiber components are removed from $\overline{u}_\infty$, then the ECH index of each level $\overline{v}_* \neq \overline{v}_0$ is nonnegative and the only components of $\overline{u}_\infty$ which have negative ECH index are the following:*

- *branched covers of $\sigma_\infty^{T'}$; and*
- *at most one component $\widetilde{v}$ of $\overline{v}_0''$ with $I(\widetilde{v}) = -1$.*

Proof. Similar to the proofs of Lemmas 3.7.4 and 3.7.5. □

The following lemma is analogous to Lemma 3.7.10.

Lemma 3.9.2. *If $m \gg 0$ and $\overline{u}_\infty \in \partial_{(-\infty,+\infty)}\mathcal{M}$, then the following hold:*

1. *$\overline{u}_\infty$ has no fiber components;*
2. *there is no level $\overline{v}_*$ such that $\overline{v}_*' \cap \overline{v}_*'' \neq \varnothing$ and $\overline{v}_*' \cap \overline{v}_*'' \subset int(\overline{v}_*')$; and*
3. *$\overline{u}_\infty$ has no boundary puncture which is removable at z_∞.*

Proof. (1) Arguing by contradiction, suppose that $\overline{u}_\infty$ has a fiber component $\widetilde{v}$. Then $n^*(\widetilde{v}) \geq m$ and $\widetilde{v}$ can be eliminated by arguing as in Lemma 3.7.10(1), in view of the following contributions towards I:

(γ_1) $I(\overline{v}_0') = -p_0$.

(γ_2) $\cup_{j=1}^c \overline{v}_{-1,j}^\sharp$ is a union of p_0 trivial strips and $\sum_{j=1}^c I(\overline{v}_{-1,j}^\sharp) = p_0$.

(γ_3) $\widetilde{v}$ and the intersection $\widetilde{v} \cap \overline{v}_*$ for an appropriate $\overline{v}_*$ contribute $2g+2 \geq 4$ towards I.

(γ_4) After removing the fiber component $\widetilde{v}$, all the levels $\neq \overline{v}_0$ have nonnegative ECH index and $I(\overline{v}_0'') \geq -1$ by Lemma 3.9.1.

The sum of the above ECH indices is at least 3, a contradiction.

(2), (3) Similar to (2) and (3) of Lemma 3.7.10. Observe that the sum of ECH indices of the levels was at least 4 in all the cases of Lemma 3.7.10 that were not ruled out by other means; in the present case the sum will be at least 3. □

Lemma 3.9.3. *If $\overline{u}_\infty \in \partial_{(-\infty,+\infty)}\mathcal{M}$ and $\overline{v}_0' = \varnothing$, then $\overline{u}_\infty$ is one of the following 2-level buildings:*

1. *$\overline{v}_{1,1}$ with $I=1$ from $\mathbf{z}$ to some $\mathbf{z}'$ consisting of*
    1. (i) *one thin strip and trivial strips or*
    2. (ii) *one nontrivial component of $\overline{v}_{1,1}''$ with image in W and trivial strips;*

   and $\overline{v}_0 = \overline{v}_0''$ with $I=1$ and $n^* \leq m + |\mathcal{I}|$ from $\mathbf{z}'$ to $\mathbf{y}'$.
2. *$\overline{v}_0 = \overline{v}_0''$ with $I=1$ and $n^* \leq m+|\mathcal{I}|$ from $\mathbf{z}$ to some $\mathbf{y}''$; and $\overline{v}_{-1,1}$ with $I=1$ from $\mathbf{y}''$ to $\mathbf{y}'$.*

Proof. Suppose that $\overline{v}_0' = \varnothing$. Since passing through $\overline{\mathfrak{m}}(T')$ is *effectively* a codimension two condition, we have $I(\overline{v}_0) \geq 1$. Hence there can be only one other nontrivial level — either $\overline{v}_{1,1}$ or $\overline{v}_{-1,1}$ — and $I(\overline{v}_{\pm1,1}) \geq 1$. By Lemma 3.9.1,

$I(\overline{v}_0)=1$ and $I(\overline{v}_{\pm 1,1})=1$. Moreover, $n^*(\overline{v}_{\pm 1,1})\leq |\mathcal{I}|$ since $n^*(\overline{v}_0)\geq m$. This means that there are no components $\overline{v}_*^\sharp$ that limit to z_∞ at the negative end and all the positive ends $\mathcal{E}_+$ that limit to z_∞ satisfy $n^*(\mathcal{E}_+)=1$. The lemma follows. □

Lemma 3.9.4. *For each interval $[-T,T]$, there exists $m\gg 0$ such that there is no sequence of curves $\overline{u}_i\in\mathcal{M}_{\tau_i}$, $\tau_i\to T'\in[-T,T]$, that limits to $\overline{u}_\infty$ for which $\overline{v}_0'\neq\varnothing$.*

Proof. This is similar to Lemma 3.6.5 and will be omitted. □

Proof of Lemma 3.3.12. This is argued in the same way as Lemma 3.3.8. Suppose $\overline{u}_\infty\in\partial_{(-\infty,+\infty)}\mathcal{M}$. If $\overline{v}_0'=\varnothing$, then Lemma 3.9.3 implies that $\overline{u}_\infty$ is as described in Lemma 3.3.12. On the other hand, by Lemma 3.9.4, for any $T>0$ there exists $m\gg 0$ such that $\overline{v}_0'=\varnothing$ in $\overline{u}_\infty\in\partial_{[-T,+T]}\mathcal{M}$. The possibilities where $m_i\to\infty$, $T_i\to\infty$, $\overline{u}_{ij}\to\overline{u}_{i\infty}\in\partial_{\{\pm T_i\}}\mathcal{M}$ are treated in the same way as in Lemmas 3.7.14, 3.8.3, and 3.8.11. □

4. HOMOTOPY OF COBORDISMS II

In this section we consider the homotopy of cobordisms corresponding to $\Phi\circ\Psi$. Many of the constructions for $\overline{W}_\tau^\mp$ are similar to those of $\overline{W}_\tau^\pm$ with minor modifications.

We first give a brief description of $\overline{W}_\tau=\overline{W}_\tau^\mp$. (If $\mp$ is understood, at it will be in the rest of this section, then it will be omitted.) The base B_τ of $\overline{W}_\tau$ is biholomorphic to an infinite cylinder with a disk removed. As $\tau\to+\infty$, $\overline{W}_\tau$ degenerates to the stacking of $\overline{W}_-$ "on top of" $\overline{W}_+$. On the other hand, as $\tau\to-\infty$, $\overline{W}_\tau$ degenerates to $\overline{W}_{-\infty}$, whose base $B_{-\infty}$ is given by:

$$B_{-\infty}=((\mathbb{R}\times(\mathbb{R}/2\mathbb{Z}))\sqcup E)\,/\sim, \tag{4.0.1}$$

where $E=\{|z|\leq 1\}\subset\mathbb{C}$ and $\sim$ identifies $(0,\frac{3}{2})\in\mathbb{R}\times(\mathbb{R}/2\mathbb{Z})$ with $0\in E$.

4.1. Construction of the homotopy of cobordisms for $\Phi\circ\Psi$.

4.1.1. *Definition of the family $\overline{W}_\tau$.* Let $l\in(0,\infty)$ and $r\in(0,1]$. Consider the fibration

$$\pi:\mathbb{R}\times\overline{N}\to\mathbb{R}\times(\mathbb{R}/2\mathbb{Z}),$$

where $\overline{N}$ is viewed as $(\overline{S}\times[0,2])/(x,2)\sim(\overline{h}(x),0)$ and (s,t) are coordinates on $\mathbb{R}\times(\mathbb{R}/2\mathbb{Z})$. We define $\overline{W}_{l,r}=\pi^{-1}(B_{l,r})$, where the base $B_{l,r}$ is obtained by smoothing the corners of

$$(\mathbb{R}\times(\mathbb{R}/2\mathbb{Z}))-([-l/2,l/2]\times[(3-r)/2,(3+r)/2]).$$

Next choose a function

$$\eta=(l,r):\mathbb{R}\to(0,\infty)\times(0,1], \tag{4.1.1}$$

which is obtained by smoothing

$$\eta_0(\tau)=\begin{cases}(\tau+1,1), & \text{for } \tau\geq 0;\\ (e^\tau,e^\tau), & \text{for } \tau\leq 0;\end{cases}$$

near $\tau=0$.

We then define $\overline{W}_\tau = \overline{W}_{\eta(\tau)}$ and $B_\tau = B_{\eta(\tau)}$. Let $\pi_{B_\tau} : \overline{W}_\tau \to B_\tau$ be the projection along $\{(s,t)\} \times \overline{S}$.

As $\tau \to +\infty$, the cobordism $\overline{W}_\tau$ approaches the concatenation of $\overline{W}_-$ and $\overline{W}_+$. See Figure 16. On the other hand, as $\tau \to -\infty$, the cobordism $\overline{W}_\tau$ can be viewed as degenerating to $\overline{W}_{-\infty} = (\overline{W}_{-\infty,1} \sqcup \overline{W}_{-\infty,2})/\sim$, which we describe now: Consider the base $B_{-\infty} = (B_{-\infty,1} \sqcup B_{-\infty,2})/\sim$, where $B_{-\infty,1} = \mathbb{R} \times (\mathbb{R}/2\mathbb{Z})$, $B_{-\infty,2} = E = \{|z| \leq 1\}$, and $\sim$ identifies $(0, \frac{3}{2}) \in B_{-\infty,1}$ with $0 \in B_{-\infty,2}$. We also identify the asymptotic markers ∂_s at $(0, \frac{3}{2})$ and $\{x = 0, y > 0\}$ at 0. We then set $\overline{W}_{-\infty,1} = \pi^{-1}(B_{-\infty,1})$ and $\overline{W}_{-\infty,2} = B_{-\infty,2} \times \overline{S}$, and identify $(0, \frac{3}{2}, x) \sim (0, x)$, where $(0, \frac{3}{2}, x) \in \overline{W}_{-\infty,1}$ and $(0, x) \in \overline{W}_{-\infty,2}$. We write $\pi_{B_{-\infty,i}} : \overline{W}_{-\infty,i} \to B_{-\infty,i}$ for the projection along $\overline{S}$.



FIGURE 16. The bases of the family $\overline{W}^{\mp}_\tau$. The leftmost diagram represents $B_{-\infty}$, i.e., $\tau = -\infty$, and the parameter τ increases as we go to the right. The location of $\overline{\mathfrak{m}}^b(\tau)$ is indicated by ×.

4.1.2. *Marked points.* We choose a 1-parameter family of marked points

$$\overline{\mathfrak{m}}(\tau) = (\overline{\mathfrak{m}}^b(\tau), \overline{\mathfrak{m}}^f(\tau)) = (\overline{\mathfrak{m}}^b(\tau), z_\infty) \in \overline{W}_\tau$$

for $\tau \in \mathbb{R}$, such that the following hold:

(i) $\overline{\mathfrak{m}}^b(\tau)$ is on the segment $\{s > \frac{l(\tau)}{2}, t = \frac{3}{2}\}$;

(ii) as $\tau \to +\infty$, $\overline{\mathfrak{m}}(\tau)$ limits to $\overline{\mathfrak{m}}(+\infty) = (\overline{\mathfrak{m}}^b(+\infty), \overline{\mathfrak{m}}^f(+\infty))$, where $\overline{\mathfrak{m}}^b(+\infty) = (0, \frac{3}{2}) \in B_-$ and $\overline{\mathfrak{m}}^f(+\infty) = z_\infty$;

(iii) as $\tau \to -\infty$, $\overline{\mathfrak{m}}^b(\tau)$ limits to $\overline{\mathfrak{m}}^b(-\infty) = (\overline{\mathfrak{m}}^b(-\infty), \overline{\mathfrak{m}}^f(\infty))$, where $\overline{\mathfrak{m}}^b(+\infty) = \frac{i}{2} \in B_{-\infty,2}$ and $\overline{\mathfrak{m}}^f(+\infty) = z_\infty$.

4.1.3. *Stable Hamiltonian structures and symplectic forms.* We first consider $\overline{W}_\tau$. The stable Hamiltonian structure on $\overline{N} = (\overline{S} \times [0,2])/\sim$ is obtained from $(dt, \overline{\omega})$ on $\overline{S} \times [0,2]$ by passing to the quotient. The 2-plane field is $\xi_\tau = T\overline{S}$ and the Hamiltonian vector field is $R_\tau = \partial_t$. The symplectic form $\overline{\Omega}_\tau$ is the restriction to $\overline{W}_\tau$ of $ds \wedge dt + \overline{\omega}$, defined on $\mathbb{R} \times \overline{S} \times [0,2]$ and sent to the quotient $\mathbb{R} \times \overline{N}$.

Next we consider $\overline{W}_{-\infty}$. For $\overline{W}_{-\infty,1} = \overline{N}$ the Hamiltonian structure is $(dt, \overline{\omega})$ and the symplectic form is $\overline{\Omega}_{-\infty,1} = ds \wedge dt + \overline{\omega}$. The symplectic form on $\overline{W}_{-\infty,2} = B_{-\infty,2} \times \overline{S}$ is a product symplectic form $\overline{\Omega}_{-\infty,2} = \omega_{-\infty,2} \oplus \overline{\omega}$, where $\omega_{-\infty,2}$ is the standard area form on the unit disk.

4.2. **Holomorphic curves and moduli spaces.**

4.2.1. *Lagrangian boundary conditions.* Consider the cobordism $(\overline{W}_\tau, \overline{\Omega}_\tau)$. We place a copy of $\overline{\mathbf{a}}$ on $\pi_{B_\tau}^{-1}(\frac{l(\tau)}{2}, \frac{3}{2})$ and parallel transport along the vertical boundary $\partial_v \overline{W}_\tau := \pi^{-1}(\partial B_\tau)$ using the symplectic connection $\overline{\Omega}_\tau$ to obtain $L^\tau_{\overline{\mathbf{a}}}$. Similarly, we place a copy of $\overline{\mathbf{a}}$ over $1 \in B_{-\infty,2}$ and parallel transport along $\partial B_{-\infty,2}$ to obtain $L^{-\infty,2}_{\overline{\mathbf{a}}} = \partial B_{-\infty,2} \times \overline{\mathbf{a}}$. The Lagrangian submanifolds $L^\tau_{\widehat{\mathbf{a}}}$, $L^\tau_{\widehat{a}_i}$, $L^\tau_{\overline{a}_i}$ are defined similarly.

4.2.2. *Almost complex structures.*

Definition 4.2.1.

(1) An almost complex structure $\overline{J}_\tau$ on $\overline{W}_\tau$ is *admissible* if $\overline{J}_\tau$ is the restriction of some $\overline{J'} \in \overline{\mathcal{J}'}$. In this case $\overline{J}_\tau$ is said to be *compatible* with $\overline{J'}$.
(2) An almost complex structure $\overline{J}_{-\infty,1}$ on $\mathbb{R} \times \overline{N}$ is *admissible* if $\overline{J}_{-\infty,1} \in \overline{\mathcal{J}'}$.
(3) An almost complex structure $\overline{J}_{-\infty,2}$ on $B_{-\infty,2} \times \overline{S}$ is *admissible* if it is a product complex structure.

The space of admissible $\overline{J}_\tau$, $\overline{J}_{-\infty,1}$ and $\overline{J}_{-\infty,2}$ will be denoted by $\overline{\mathcal{J}}_\tau$, $\overline{\mathcal{J}}_{-\infty,1}$ and $\overline{\mathcal{J}}_{-\infty,2}$, respectively.

Definition 4.2.2. A family $\{\overline{J}_\tau \in \overline{\mathcal{J}}_\tau\}_{\tau \in \mathbb{R}}$ of almost complex structures is *admissible* if there exist $\overline{J'} \in \overline{\mathcal{J}'}$, $\overline{J} \in \overline{\mathcal{J}}$, $\overline{J}_+ \in \overline{\mathcal{J}}_+$, $\overline{J}_- \in \overline{\mathcal{J}}_-$, $\overline{J}_{-\infty,1} \in \overline{\mathcal{J}}_{-\infty,1}$ and $\overline{J}_{-\infty,2} \in \overline{\mathcal{J}}_{-\infty,2}$ such that the following hold:

(1) $\overline{J}_\tau$ converges to $(\overline{J}_{-\infty,1}, \overline{J}_{-\infty,2})$ as $\tau \to -\infty$;
(2) $\overline{J}_\tau$ converges to $(\overline{J}_+, \overline{J}_-)$ as $\tau \to +\infty$;
(3) $\overline{J}_+$ and $\overline{J}_-$ are compatible with $\overline{J'}$ and $\overline{J}$; and
(4) $\overline{J}_\tau$ is compatible with $\overline{J'} = \overline{J}_{-\infty,1}$.

The space of all admissible $\{\overline{J}_\tau \in \overline{\mathcal{J}}_\tau\}_{\tau \in \mathbb{R}}$ will be denoted by $\overline{\mathcal{I}}$.

4.2.3. *Notation and conventions.* We will be using the notation and conventions from Section 3.2.3, with the exception of intersection numbers $n^*(\overline{u})$.

Intersection numbers. Let δ_{ρ_0,ϕ_0} be a closed orbit of the Hamiltonian vector field ∂_t which lies on the torus $\{\rho = \rho_0\} \subset \overline{S} \times [0,2]/\sim$ for $\rho_0 > 0$ sufficiently small and which passes through the point $(t, \phi) = (0, \phi_0)$. We assume additionally that δ_{ρ_0,ϕ_0} does not intersect the projections of the Lagrangians of $\overline{W}_\tau$, $\overline{W}_+$ and $\overline{W}_-$ to $\overline{N}$.

We then write $(\sigma^*_\infty)^\dagger$ for the restriction of $\mathbb{R} \times \delta_{\rho_0,\phi_0}$ to $\overline{W}_*$, where $* = \varnothing, ', \tau, +,$ $-$ or $(-\infty, 1)$. For $\overline{W}_{-\infty,2}$ we write

$$(\sigma^{-\infty,2}_\infty)^\dagger = B_{-\infty,2} \times \{\rho = \rho_0, \phi = \phi_0 + 3\pi/2m + 2\pi k/m, k \in \mathbb{Z}\}.$$

Finally we define $n^*(\overline{u}) = \langle \overline{u}, (\sigma^*_\infty)^\dagger \rangle$, where $* = \varnothing, ', \tau, +, -$ or $(-\infty, 1)$.

4.2.4. *Holomorphic maps to* $\overline{W}_\tau$. Let (F,j) be a compact Riemann surface, possibly disconnected, with $2g$ boundary components and two sets of interior punctures $\mathbf{p}^+ = \{p_1^+,\dots,p_{k_+}^+\}$ and $\mathbf{p}^- = \{p_1^-,\dots,p_{k_-}^-\}$ such that each component of F has nonempty boundary, at least one puncture from $\mathbf{p}^+$, and at least one puncture from $\mathbf{p}^-$. We write $\dot{F} = F - \mathbf{p}^+ - \mathbf{p}^-$.

Definition 4.2.3. Let $\overline{J}_\tau \in \overline{\mathcal{J}}_\tau$ and let $\gamma = \prod \gamma_i^{m_i}, \gamma' = \prod (\gamma_i')^{m_i'} \in \widehat{\mathcal{O}}_k$.

A *degree k multisection of* $(\overline{W}_\tau, \overline{J}_\tau)$ *from γ to γ'* is a holomorphic map

$$\overline{u} : (\dot{F}, j) \to (\overline{W}_\tau, \overline{J}_\tau)$$

which is a degree k multisection of $\pi_{B_\tau} : \overline{W}_\tau \to B_\tau$ and which additionally satisfies the following:

1. $\overline{u}(\partial F) \subset L^\tau_{\widehat{\mathbf{a}}}$;
2. $\overline{u}$ maps each connected component of ∂F to a different $L^\tau_{\widehat{a}_i}$;
3. $\lim_{w\to p_i^+} \pi_\mathbb{R} \circ \overline{u}(w) = +\infty$ and $\lim_{w\to p_i^-} \pi_\mathbb{R} \circ \overline{u}(w) = -\infty$;
4. $\overline{u}$ converges to a cylinder over a multiple of some γ_j near each puncture p_i^+ so that the total multiplicity of γ_j over all the p_i^+'s is m_j (and similarly for p_i^-).

Here $\pi_\mathbb{R} : \overline{W}_\tau \to \mathbb{R}$ is the projection to the s-coordinate.

A $(\overline{W}_\tau, \overline{J}_\tau)$*-curve from γ to γ'* is a degree $2g$ multisection of $(\overline{W}_\tau, \overline{J}_\tau)$ satisfying $n^*(\overline{u}) = m$.

Let $\mathcal{M}_{\overline{J}_\tau}(\gamma,\gamma')$ be the moduli space of degree k multisections of $(\overline{W}_\tau, \overline{J}_\tau)$ from γ to γ' and let $\mathcal{M}_{\overline{J}_\tau}(\gamma,\gamma';\overline{\mathfrak{m}}(\tau)) \subset \mathcal{M}_{\overline{J}_\tau}(\gamma,\gamma')$ be the subset of curves that pass through $\overline{\mathfrak{m}}(\tau)$. We write

$$\mathcal{M}_{\{\overline{J}_\tau\}}(\gamma,\gamma') = \coprod_{\tau\in\mathbb{R}} \mathcal{M}_{\overline{J}_\tau}(\gamma,\gamma'),$$

$$\mathcal{M}_{\{\overline{J}_\tau\}}(\gamma,\gamma';\overline{\mathfrak{m}}) = \coprod_{\tau\in\mathbb{R}} \mathcal{M}_{\overline{J}_\tau}(\gamma,\gamma';\overline{\mathfrak{m}}(\tau)).$$

4.2.5. *Holomorphic maps to* $\overline{W}_{-\infty}$. We now describe the curves in $\overline{W}_{-\infty,i}$.

Definition 4.2.4. Let $\overline{J}_{-\infty,1} \in \overline{\mathcal{J}}_{-\infty,1}$ and let $\gamma = \prod \gamma_i^{m_i}, \gamma' = \prod (\gamma_i')^{m_i'} \in \widehat{\mathcal{O}}_k$. A *degree k multisection of* $(\overline{W}_{-\infty,1}, \overline{J}_{-\infty,1})$ *from γ to γ'* is a holomorphic map

$$\overline{u} : (\dot{F}, j) \to (\overline{W}_{-\infty,1}, \overline{J}_{-\infty,1})$$

which is a degree k multisection of $\pi_{B_{-\infty,1}} : \overline{W}_{-\infty,1} \to B_{-\infty,1}$ and which is asymptotic to γ and γ' at the positive and negative ends. Here $(\dot{F}, j)$ is a punctured Riemann surface. A $(\overline{W}_{-\infty,1}, \overline{J}_{-\infty,1})$*-curve from γ to γ'* is a degree $2g$ multisection of $(\overline{W}_{-\infty,1}, \overline{J}_{-\infty,1})$ satisfying $n^*(\overline{u}) = 0$.

Definition 4.2.5. Let $\overline{J}_{-\infty,2} \in \overline{\mathcal{J}}_{-\infty,2}$. A *degree k multisection of* $(\overline{W}_{-\infty,2}, \overline{J}_{-\infty,2})$ is holomorphic map

$$\overline{u} : (F, j) \to (\overline{W}_{-\infty,2}, \overline{J}_{-\infty,2})$$

which is a degree k multisection of $\pi_{B_{-\infty,2}}:\overline{W}_{-\infty,2}\to B_{-\infty,2}$ and which additionally satisfies the following:

(1) (F,j) is a compact Riemann surface with k boundary components;
(2) $\overline{u}$ maps each component of ∂F to a different $L^{-\infty,2}_{\widehat{a}_i}$.

A $(\overline{W}_{-\infty,2},\overline{J}_{-\infty,2})$*-curve* is a degree $2g$ multisection of $(\overline{W}_{-\infty,2},\overline{J}_{-\infty,2})$ satisfying $n^*(\overline{u})=m$. A *degenerate* $(\overline{W}_{-\infty,2},\overline{J}_{-\infty,2})$*-curve* consists of $2g$ copies of $B_{-\infty,2}\times\{pt\}$ and a singular fiber $\{0\}\times\overline{S}$.

Let $\mathcal{M}_{\overline{J}_{-\infty,1}}(\gamma,\gamma')$ be the moduli space of degree k multisections of $(\overline{W}_{-\infty,1},\overline{J}_{-\infty,1})$ from γ to γ' and let $\mathcal{M}_{\overline{J}_{-\infty,2}}$ be the moduli space of degree k multisections of $(\overline{W}_{-\infty,2},\overline{J}_{-\infty,2})$.

Let $\mathfrak{z}$ be a formal product $\zeta_0^{r_0}\zeta_1^{r_1}\dots\zeta_l^{r_l}$, where $r_0,\dots,r_l\in\mathbb{Z}^{\geq 0}$, $r_0+\dots+r_l=k$, $\zeta_0=z_\infty$, and $\zeta_1,\dots,\zeta_l\in\overline{S}-\{z_\infty\}$. We write $\mathfrak{Z}$ for the set of all formal products $\mathfrak{z}$ on $\overline{S}$. A degree k multisection $\overline{u}\in\mathcal{M}_{\overline{J}_{-\infty,1}}(\gamma,\gamma')$ (resp. $\mathcal{M}_{\overline{J}_{-\infty,2}}$) *passes through* $\mathfrak{z}$ if it passes through $((0,\frac{3}{2}),\zeta_i)$ (resp. $(0,\zeta_i)$), $i=0,\dots,l$, with multiplicities r_i. Here the multiplicity r_i is the total degree of the projection $\pi_{B_{-\infty,i}}\circ\overline{u}$, restricted to neighborhoods of points in the domain $\dot{F}$ (resp. F) which map to $((0,\frac{3}{2}),\zeta_i)$ (resp. $(0,\zeta_i)$) under $\overline{u}$.

Let

$$\mathcal{M}_{\overline{J}_{-\infty,1}}(\gamma,\gamma',\mathfrak{z})\subset\mathcal{M}_{\overline{J}_{-\infty,1}}(\gamma,\gamma'),\quad \mathcal{M}_{\overline{J}_{-\infty,2}}(\mathfrak{z})\subset\mathcal{M}_{\overline{J}_{-\infty,2}}$$

be subsets of curves that pass through $\mathfrak{z}$. Also let

$$\mathcal{M}_{\overline{J}_{-\infty,2}}(\mathfrak{z},\overline{\mathfrak{m}}(-\infty))\subset\mathcal{M}_{\overline{J}_{-\infty,2}}(\mathfrak{z})$$

be the subset of curves that pass through $\overline{\mathfrak{m}}(-\infty)$.

4.2.6. *Indices.* We now briefly discuss the Fredholm index $\operatorname{ind}(\overline{u})$ and the ECH index $I(\overline{u})$ of a $\overline{W}_\tau$-curve $\overline{u}:\dot{F}\to\overline{W}_\tau$ from γ to γ'. This is similar to Section 3.2.6.

Let $\check{\overline{W}}_\tau=\overline{W}_\tau-\{s>\frac{l(\tau)}{2}+1\}-\{s<-\frac{l(\tau)}{2}-1\}$, where $l(\tau)$ is given in Equation (4.1.1). Let $\check{\overline{u}}:\check{F}\to\check{\overline{W}}_\tau$ be the compactification of $\overline{u}$, where $\check{F}$ is obtained by performing a real blow-up of F at its boundary punctures.

The trivialization $\tau^\star$ of $T\overline{S}$ along $L^\tau_{\widehat{\mathbf{a}}}$ is defined as in Section I.4.4.2: We pick a point $p\in\partial B_\tau$, define $\tau^\star$ of $T\overline{S}|_{\pi^{-1}_{B_\tau}(p)}$ along $\widehat{\mathbf{a}}$ by choosing a nonsingular tangent vector field along $\widehat{\mathbf{a}}$, and then parallel transport $\tau^\star$ along $\partial\overline{W}_\tau$. We also choose $\tau^\star$ along γ and γ'.

Let $Q_{\tau^\star}(\check{\overline{u}})$ be the relative intersection form given by normalizing $\check{\overline{u}}$ near $s=\pm(\frac{l(\tau)}{2}+1)$ as in [Hu1, Definition 2.4] and intersecting $\check{\overline{u}}$ and a pushoff of $\check{\overline{u}}$. Here the pushoff along $\partial\check{\overline{W}}_\tau$ is in the direction of $J\tau^\star$. Then

$$I(\overline{u})=c_1(\check{\overline{u}}^*T\check{\overline{W}}_\tau,(\tau^\star,\partial_t))+Q_{\tau^\star}(\check{\overline{u}})+\widetilde{\mu}_{\tau^\star}(\gamma)-\widetilde{\mu}_{\tau^\star}(\gamma')-2g,\tag{4.2.1}$$

$$\operatorname{ind}(\overline{u})=-\chi(\dot{F})+2c_1(\check{\overline{u}}^*T\check{\overline{W}}_\tau,(\tau^\star,\partial_t))+\mu_{\tau^\star}(\gamma)-\mu_{\tau^\star}(\gamma')-2g,\tag{4.2.2}$$

and the index inequality holds as usual:

$$\operatorname{ind}(\overline{u}) + 2\delta(\overline{u}) \leq I(\overline{u}), \tag{4.2.3}$$

where $\delta(\overline{u}) \geq 0$ and equals zero if and only if $\overline{u}$ is an embedding. We also have

$$I(\sigma_\infty^\tau) = \operatorname{ind}(\sigma_\infty^\tau) = -1. \tag{4.2.4}$$

The Fredholm and ECH indices for $\overline{W}_{-\infty,i}$-curves can be defined and computed similarly.

Remark 4.2.6. The Fredholm and ECH indices for $\overline{W}_\tau$ and $\overline{W}_{-\infty,i}$ do not take into account the point constraint $\overline{\mathfrak{m}}(\tau)$ and the condition "passing through $\overline{\mathfrak{m}}(\tau)$" is a codimension 2 condition. Moreover, the indices for $\overline{W}_{-\infty,i}$, $i = 1, 2$, do not take into account the constraints $\mathfrak{z}$.

We now have the following:

Lemma 4.2.7. *Let $\overline{u}$ be a $\overline{W}_{-\infty,2}$-curve. Then*

$$[\overline{u}] = 2g[B_{-\infty,2} \times \{pt\}] + [\{pt\} \times \overline{S}] \in H_2(\overline{W}_{-\infty,2}, \partial B_{-\infty,2} \times \overline{S}),$$

$\overline{u}$ consists of $0 \leq k < 2g$ copies of $B_{-\infty,2} \times \{pt\}$ together with an irreducible component in the class $(2g-k)[B_{-\infty,2} \times \{pt\}] + [\{pt\} \times \overline{S}]$, and $I(\overline{u}) = 4g+2$.

Proof. Let $\pi_{\overline{S}} : \overline{W}_{-\infty,2} \to \overline{S}$ be the projection along $B_{-\infty,2}$. Since $\overline{J}_{-\infty,2}$ is a split complex structure, $\pi_{\overline{S}}$ is a holomorphic map. Hence if $\overline{u}$ is a $\overline{W}_{-\infty,2}$-curve, then $\pi_{\overline{S}} \circ \overline{u}$ either maps to a point on some $\widehat{a}_i$ or to all of $\overline{S}$. The lemma follows by listing all the possibilities. □

We also have the following, which is stated without proof.

Lemma 4.2.8. *Let $\overline{u}$ be a degree $2g$ multisection of $(\overline{W}_{-\infty,2}, \overline{J}_{-\infty,2})$ satisfying $n^*(\overline{u}) = 0$. Then $\overline{u} = \overline{u}' \cup \overline{u}''$ consists of $\overline{u}'$ which is a degree k cover of $\sigma_\infty^{-\infty,2}$ and $\overline{u}''$ which is the union of $2g-k$ copies of $B_{-\infty,2} \times \{pt\}$, and $I(\overline{u}) = 2g$.*

4.2.7. *Regularity.*

Definition 4.2.9. The family $\{\overline{J}_\tau\}_{\tau\in\mathbb{R}} \in \overline{\mathcal{I}}$ is *regular* if:

1. $\mathcal{M}^\dagger_{\{\overline{J}_\tau\}}(\delta_0^p\gamma, \delta_0^q\gamma')$ is transversely cut out for all $\delta_0^p\gamma, \delta_0^q\gamma' \in \widehat{\mathcal{O}}_k$, $k \leq 2g$;
2. the restriction $\overline{J}'$ of $\overline{J}_\tau$ to the positive and negative ends (the restriction is independent of τ) is regular; and
3. $\overline{J}_-$ and $\overline{J}_+$ in the limit $\tau \to +\infty$ are regular.

Let $\overline{\mathcal{I}}^{reg}$ be the space of regular $\{\overline{J}_\tau\} \in \overline{\mathcal{I}}$. As usual we have:

Lemma 4.2.10. *The generic $\{\overline{J}_\tau\} \in \overline{\mathcal{I}}$ is regular.*

We also introduce the perturbations of $\{\overline{J}_\tau\}$ to ensure that passing through $\overline{\mathfrak{m}}(\tau)$ is a generic condition. Let $\mathfrak{p}(\tau) \in B_\tau$ be a family of points such that:

$$\lim_{\tau\to+\infty} \mathfrak{p}(\tau) = \mathfrak{p}(+\infty) \in B_-, \quad \lim_{\tau\to-\infty} \mathfrak{p}(\tau) = \mathfrak{p}(-\infty) \in B_{-\infty,1}$$

and $\mathfrak{p}(\tau) \neq \overline{\mathfrak{m}}^b(\tau)$ for all $\tau \in [-\infty, \infty]$. We then define the families $\{U_\tau = U_{\varepsilon,\delta,\mathfrak{p}(\tau)}\}$, $\{K_\tau = K_{\mathfrak{p}(\tau),\delta}\}$, and $\{\overline{J}_\tau^\Diamond = \overline{J}_\tau^\Diamond(\varepsilon, \delta, \mathfrak{p}(\tau))\}$ as in Section 3.2.7. Also, the modifier $\{K_\tau\}$ means "passing through K_τ for an appropriate τ."

Definition 4.2.11. The family $\{\overline{J}_\tau^\Diamond\}$ is $\{K_\tau\}$*-regular with respect to* $\overline{\mathfrak{m}}$ if the moduli spaces $\mathcal{M}^{\dagger,\{K_\tau\}}_{\{\overline{J}_\tau\}}(\delta_0^p\gamma, \delta_0^q\gamma'; \overline{\mathfrak{m}})$ are transversely cut out.

The following lemmas are analogous to Lemmas 3.2.16 and 3.2.17:

Lemma 4.2.12. *A generic* $\{\overline{J}_\tau^\Diamond\}_{\tau\in\mathbb{R}}$ *is* $\{K_\tau\}$*-regular with respect to* $\overline{\mathfrak{m}}$.

Lemma 4.2.13. *If* $\{\overline{J}_\tau\}$ *is a generic family, then for* $\varepsilon, \delta > 0$ *sufficiently small, there exist a generic family* $\{\overline{J}_\tau^\Diamond(\varepsilon, \delta, \mathfrak{p}(\tau))\}$ *which is* $\{K_{\mathfrak{p}(\tau),\delta}\}$*-regular with respect to* $\overline{\mathfrak{m}}$ *and disjoint finite subsets* $\mathcal{T}_1, \mathcal{T}_2 \subset \mathbb{R}$ *with the following properties:*

1. $\tau \in \mathcal{T}_1$ *if and only if there exists* $\overline{v}_\tau \in \mathcal{M}^{\dagger,s,irr,\mathrm{ind}=-1}_{\overline{J}_\tau}(\delta_0^p\gamma, \delta_0^q\gamma')$ *for some* $\delta_0^p\gamma$ *and* $\delta^q\gamma'$.
2. $\tau \in \mathcal{T}_2$ *if and only if there exists* $\overline{v}_\tau \in \mathcal{M}^{\dagger,s,irr,\{K_\tau\},\mathrm{ind}=1}_{\overline{J}_\tau^\Diamond(\varepsilon,\delta,\mathfrak{p}(\tau))}(\delta_0^p\gamma, \delta_0^q\gamma')$ *for some* $\delta_0^p\gamma$ *and* $\delta^q\gamma'$.

Moreover, for each $\tau \in \mathcal{T}_i$ *there is a unique such irreducible curve* $\overline{v}_\tau$.

4.3. **Proof of the other half of Theorem 1.0.1.** In this subsection we prove the "other half" of Theorem 1.0.1, i.e., $\Phi \circ \Psi = id$ on the level of homology, assuming the results of Sections 4.4–4.5.

We make the following simplifying assumption, which is possible by Theorem I.2.5.2.

($\dagger\dagger$) All the elliptic orbits of the stable Hamiltonian vector field corresponding to $h : S \xrightarrow{\sim} S$ that intersect $S \times \{0\}$ at most $2g$ times have linearized first return maps which are ε-rotations with $\varepsilon > 0$ small.

Theorem 4.3.1. *Suppose (*$\dagger\dagger$*) holds. If* $m \gg 0$*, then there exists a chain homotopy*

$$K : PFC_{2g}(N) \to PFC_{2g}(N),$$

such that the following holds:

$$\partial_{ECH} \circ K + K \circ \partial_{ECH} = \Phi \circ \Psi + id. \tag{4.3.1}$$

Proof. Suppose $m \gg 0$. Fix $\mathfrak{p}(\tau)$ and choose $\{\overline{J}_\tau\} \in \overline{\mathcal{I}}^{reg}$ satisfying Lemma 4.2.13. For sufficiently small $\varepsilon, \delta > 0$ (which depend on the choices of m and $\{\overline{J}_\tau\}$), there exists $\{\overline{J}_\tau^\Diamond(\varepsilon, \delta, \mathfrak{p}(\tau))\}$ so that Lemma 4.2.13 holds.

Fix $\gamma, \gamma' \in \widehat{\mathcal{O}}_{2g}$ and abbreviate

$$\mathcal{M} = \mathcal{M}^{I=2,n^*=m}_{\{\overline{J}_\tau^\Diamond(\varepsilon,\delta,\mathfrak{p}(\tau))\}}(\gamma, \gamma'; \overline{\mathfrak{m}}),\ \mathcal{M}^{\{K_{\mathfrak{p}(\tau),\delta}\}} = \mathcal{M}^{I=2,n^*=m,\{K_{\mathfrak{p}(\tau),\delta}\}}_{\{\overline{J}_\tau^\Diamond(\varepsilon,\delta,\mathfrak{p}(\tau))\}}(\gamma, \gamma'; \overline{\mathfrak{m}}).$$

Let $\overline{\mathcal{M}}$ be the SFT compactification of $\mathcal{M}$ and let $\partial\mathcal{M} = \overline{\mathcal{M}} - \mathcal{M}$ be the boundary of $\mathcal{M}$. If $U \subset [-\infty, +\infty]$, then we write $\partial_U\mathcal{M}$ for the set of $\overline{u}_\infty \in \partial M$ where $\overline{u}_\infty$ is a building which corresponds to some $\tau \in U$. By Lemma 4.2.12, we may take $\mathcal{M}^{\{K_{\mathfrak{p}(\tau),\delta}\}}$ to be regular.

In view of the considerations from Claim 3.3.4, we assume that all of $\mathcal{M}$ is regular.

Step 1 (Breaking at $+\infty$). The following is proved in Section 4.4.

Lemma 4.3.2. $\partial_{\{+\infty\}}\mathcal{M} \subset A_1$, *where*

$$A_1 = \coprod_{\mathbf{y}\in\mathcal{S}_{\mathbf{a},h(\mathbf{a})}} \left(\mathcal{M}^{I=2,n^*=m}_{\overline{J}^{\diamondsuit}_-(\varepsilon,\delta,\mathfrak{p}(+\infty))}(\gamma,\mathbf{y};\overline{\mathfrak{m}}(+\infty)) \times \mathcal{M}^{I=0}_{J_+}(\mathbf{y},\gamma') \right).$$

Gluing the pairs in A_1 accounts for the term $\Phi\circ\Psi$ in Equation (4.3.1).

Step 2 (Breaking at $-\infty$). The following lemmas are proved in Section 4.5.

Lemma 4.3.3. $\partial_{\{-\infty\}}\mathcal{M} \subset A_2 \sqcup A_3$, *where*

$$A_2 = \coprod_{\gamma\in\widehat{\mathcal{O}}_{2g},\mathfrak{z}\in\mathfrak{Z},r_0=0} \left(\mathcal{M}^{I=0,n^*=0}_{\overline{J}^{\diamondsuit}_{-\infty,1}(\varepsilon,\delta,\mathfrak{p}(-\infty))}(\gamma,\gamma,\mathfrak{z}) \times \mathcal{M}^{I=4g+2,n^*=m}_{\overline{J}_{-\infty,2}}(\mathfrak{z};\overline{\mathfrak{m}}(-\infty)) \right).$$

$$A_3 = \coprod_{\gamma,\gamma'\in\widehat{\mathcal{O}}_{2g},\mathfrak{z}\in\mathfrak{Z},r_0=1} \left(\mathcal{M}^{I=2g+2,n^*=m}_{\overline{J}^{\diamondsuit}_{-\infty,1}(\varepsilon,\delta,\mathfrak{p}(-\infty))}(\gamma,\gamma',\mathfrak{z}) \times \mathcal{M}^{I=2g,n^*=0}_{\overline{J}_{-\infty,2}}(\mathfrak{z}) \right).$$

Note that if $\overline{v}_2 \in \mathcal{M}^{I=2g,n^*=0}_{\overline{J}_{-\infty,2}}(\mathfrak{z})$, then $\overline{v}_2 = B_{-\infty,2}\times\mathfrak{z}$.

Lemma 4.3.4. *Gluing the pairs in A_2 accounts for the term id in Equation* (4.3.1).

Lemma 4.3.5. *Gluing the pairs in A_3 gives a total of* 0 *mod* 2.

Step 3 (Breaking in the middle). The following is proved in Section 4.6.

Lemma 4.3.6. $\partial_{(-\infty,+\infty)}\mathcal{M} \subset A_4 \sqcup A_5$, *where*

$$A_4 = \coprod_{\gamma''\in\widehat{\mathcal{O}}_{2g}} \left(\mathcal{M}^{I=1,n^*=m}_{\{\overline{J}^{\diamondsuit}_\tau(\varepsilon,\delta,\mathfrak{p}(\tau))\}}(\gamma,\gamma'';\overline{\mathfrak{m}}(+\infty)) \times \mathcal{M}^{I=1}_{J'}(\gamma'',\gamma') \right);$$

$$A_5 = \coprod_{\gamma''\in\widehat{\mathcal{O}}_{2g}} \left(\mathcal{M}^{I=1}_{J'}(\gamma,\gamma'') \times \mathcal{M}^{I=1,n^*=m}_{\{\overline{J}^{\diamondsuit}_\tau(\varepsilon,\delta,\mathfrak{p}(\tau))\}}(\gamma'',\gamma';\overline{\mathfrak{m}}(+\infty)) \right).$$

Gluing the pairs in A_4 and A_5 accounts for the terms $\partial_{ECH}\circ K$ and $K\circ\partial_{ECH}$ in Equation (4.3.1).

This completes the proof of Theorem 4.3.1, modulo the results from Sections 4.4–4.6. $\square$

4.4. **Degeneration at $+\infty$.** In this subsection we study the limit of holomorphic maps to $\overline{W}_\tau$ as $\tau\to+\infty$, i.e., when $\overline{W}_\tau$ degenerates into a concatenation of $\overline{W}_-$ with $\overline{W}_+$ along the HF-type end. This will prove Lemma 4.3.2.

We assume that $m \gg 0$; $\varepsilon, \delta > 0$ are sufficiently small; and $\{\overline{J}_\tau\} \in \overline{\mathcal{I}}^{reg}$ and $\{\overline{J}^\diamondsuit_\tau(\varepsilon, \delta, \mathfrak{p}(\tau))\}$ satisfy Lemma 4.2.13. Fix $\gamma, \gamma' \in \widehat{\mathcal{O}}_{2g}$ and let

$$\mathcal{M} = \mathcal{M}^{I=2, n^*=m}_{\{\overline{J}^\diamondsuit_\tau(\varepsilon,\delta,\mathfrak{p}(\tau))\}}(\gamma, \gamma'; \overline{\mathfrak{m}}), \quad \mathcal{M}_\tau = \mathcal{M}^{I=2, n^*=m}_{\overline{J}^\diamondsuit_\tau(\varepsilon,\delta,\mathfrak{p}(\tau))}(\gamma, \gamma'; \overline{\mathfrak{m}}).$$

We will analyze $\partial_{\{+\infty\}}\mathcal{M}$.

Let $\overline{u}_i$, $i \in \mathbb{N}$, be a sequence of curves in $\mathcal{M}$ such that $\overline{u}_i \in \mathcal{M}_{\tau_i}$ and $\lim\limits_{i\to\infty} \tau_i = +\infty$, and let

$$\overline{u}_\infty = (\overline{v}_{-1,1} \cup \dots \cup \overline{v}_{-1,c}) \cup \overline{v}_+ \cup (\overline{v}_{0,1} \cup \dots \cup \overline{v}_{0,b}) \cup \overline{v}_- \cup (\overline{v}_{1,1} \cup \dots \cup \overline{v}_{1,a})$$

be the limit holomorphic building in order from bottom to top, where $\overline{v}_{-1,j}$, $j = 1, \dots, c$, maps to $\overline{W'}$; $\overline{v}_+$ maps to $\overline{W}_+$; $\overline{v}_{0,j}$, $j = 1, \dots, b$, maps to $\overline{W}$; $\overline{v}_-$ maps to $\overline{W}_-$; and $\overline{v}_{1,j}$, $j = 1, \dots, a$ maps to $\overline{W'}$. Here we are allowing the possibility that a, b, or $c = 0$. For notational convenience, sometimes we refer to $\overline{v}_+$ as $\overline{v}_{-1,c+1}$ or $\overline{v}_{0,0}$ and $\overline{v}_-$ as $\overline{v}_{0,b+1}$ or $\overline{v}_{1,0}$.

As before, we have the following constraints:

$$n^*(\overline{u}_i) = \sum_{\overline{v}_*} n^*(\overline{v}_*) = m; \tag{4.4.1}$$

$$I(\overline{u}_i) = \sum_{\overline{v}_*} I(\overline{v}_*) = 2, \tag{4.4.2}$$

where the summations are over all the levels $\overline{v}_*$ of $\overline{u}_\infty$.

The following is the analog of Lemma 3.4.17, with a similar proof (omitted):

Lemma 4.4.1. *If $\overline{v}'_* \cup \overline{v}^\sharp_* = \varnothing$ for all levels $\overline{v}_*$ of $\overline{u}_\infty$, then $a = b = c = 0$; $I(\overline{v}_+) = 0$; $I(\overline{v}_-) = 2$; $\overline{v}_+$ is a W_+-curve; and $\overline{v}_-$ is a $\overline{W}_-$-curve.*

Lemma 4.4.2. *If $\overline{v}'_* \cup \overline{v}^\sharp_* \neq \varnothing$ for some level $\overline{v}_*$ of $\overline{u}_\infty$, then:*

(1) $p_- = \deg(\overline{v}'_-) > 0$;
(2) *some $\overline{v}_{1,j_0}$, $j_0 > 0$, has a negative end $\mathcal{E}_-$ that limits to δ_0^p for some $p > 0$ and satisfies $n^*(\mathcal{E}_-) \geq m - p$;*
(3) *$\overline{u}_\infty$ has no boundary puncture that is removable at z_∞;*
(4) *$\overline{u}_\infty$ has no fiber components and no components of $\overline{v}''_*$ that intersect the interior of a section at infinity;*
(5) *each component of $\overline{v}^\sharp_{0,j}$, $1 \leq j \leq b$, is a thin strip;*
(6) *each component of $\overline{v}^\sharp_+$ is an $n^* = 1$, $I = 0$ or 1 section from z_∞ to h or e which is contained in $\overline{W}_+ - W_+$;*
(7) *each component of $\overline{v}^\sharp_{-1,j}$, $0 \leq j \leq c$, is an $n^* = 1$, $I = 1$ or 2 cylinder from δ_0 to h or e which is contained in $\mathbb{R} \times (\overline{N} - N)$;*
(8) *h appears at most once at the negative end of $\overline{v}^\sharp_{-1,1}$.*

Proof. The proof is based on Equation (4.4.1). First observe that:

(*) either there is a negative end $\mathcal{E}_-$ that limits to δ_0^p for some $p > 0$ and satisfies $n^*(\mathcal{E}_-) \geq m - p$ by Lemma 3.4.1; or

(**) there is a negative end $\mathcal{E}_-$ that limits to z_∞ and the sum of $n^*(\mathcal{E}_i)$ over all the ends $\mathcal{E}_i$ that limit to z_∞ and $n^*(\mathcal{E}'_i)$ over all the neighborhoods of the boundary punctures is $\geq m-2g$.

(1) If (*) or (**) holds, then $\overline{v}'_- \neq \varnothing$, since otherwise the neighborhood of $\overline{\mathfrak{m}}(+\infty)$ contributes m towards $n^*(\overline{v}_-)$.

(2) is a consequence of (1) and is a subcase of (*). In particular (**) does not hold. (3)–(7) follow from (2). (8) follows from the definition of the ECH differential. ☐

Lemma 4.4.3. *If $\overline{v}'_*\cup\overline{v}^\sharp_* \neq \varnothing$ for some level $\overline{v}_*$, then $\overline{u}_\infty$ cannot have the following subbuildings:*

1. *a degree one component of $\overline{v}^\sharp_+$ from z_∞ to h; and*
2. *$\overline{v}'_+$ which has degree p_+ and $\overline{v}^\sharp_{-1,1}$ which is a union of p_+ cylinders from δ_0 to h or e.*

Proof. This is due to the positioning of h, given in Section 3.1.4. (1) was proved in Lemma I.6.6.5. (2) is due to Lemma 3.7.8: the usual rescaling procedure with fixed $m \gg 0$, together with Lemma 4.4.2, gives rise to an SFT limit $w_+ : \Sigma_+ \to \mathbb{CP}^1$, $\pi_+ : \Sigma_+ \to cl(B_+)$, where Σ_+ consists of p_+ copies of $cl(B_+)$ (and hence π_+ is a trivial branched cover) and the restriction of w_+ to each component of Σ_+ satisfies the conditions of Lemma 3.7.9. ☐

Lemma 4.4.4. *If $\overline{v}'_* \cup \overline{v}^\sharp_* \neq \varnothing$ for some level $\overline{v}_*$, then $\overline{v}'_- \neq \varnothing$ and $\overline{u}_\infty$ contains one of the following subbuildings:*

1. *A 3-level building consisting of $\overline{v}^\sharp_{1,1}$ with $I=1$ and a negative end $\delta_0\gamma'$; $\overline{v}'_- = \sigma^-_\infty$; and a thin strip.*
2. *A 3-level building consisting of $\overline{v}^\sharp_{1,1}$ with $I=1$ and a negative end $\delta_0\gamma'$; $\overline{v}'_- = \sigma^-_\infty$; and a component of $\overline{v}^\sharp_+$ with $I=1$ from z_∞ to e.*

Here we are omitting levels which are connectors.

Proof. The lemma is a consequence of Lemmas 4.4.2 and 4.4.3, subject to the conditions given by Equations (4.4.1) and (4.4.2). ☐

Lemma 4.4.5. *If $m \gg 0$, then there is no $\overline{u}_\infty \in \partial_{\{+\infty\}}\mathcal{M}$ such that $\overline{v}'_* \cup \overline{v}^\sharp_* \neq \varnothing$ for some level $\overline{v}_*$.*

Proof. The proofs for Cases (1) and (2) are similar to those of Theorem I.7.11.1 and will be omitted. ☐

Proof of Lemma 4.3.2. This is a combination of Lemmas 4.4.1 and 4.4.5. ☐

4.5. **Degeneration at $-\infty$.** In this subsection we study the limit of holomorphic maps to $\overline{W}_\tau$ as $\tau \to -\infty$. This will prove Lemma 4.3.3.

We assume that $m \gg 0$; $\varepsilon, \delta > 0$ are sufficiently small; and $\{\overline{J}_\tau\} \in \overline{\mathcal{I}}^{reg}$ and $\{\overline{J}^\Diamond_\tau(\varepsilon,\delta,\mathfrak{p}(\tau))\}$ satisfy Lemma 4.2.13. Fix $\gamma,\gamma' \in \widehat{\mathcal{O}}_{2g}$ and let

$$\mathcal{M} = \mathcal{M}^{I=2,n^*=m}_{\{\overline{J}^\Diamond_\tau(\varepsilon,\delta,\mathfrak{p}(\tau))\}}(\gamma,\gamma';\overline{\mathfrak{m}}), \quad \mathcal{M}_\tau = \mathcal{M}^{I=2,n^*=m}_{\overline{J}^\Diamond_\tau(\varepsilon,\delta,\mathfrak{p}(\tau))}(\gamma,\gamma';\overline{\mathfrak{m}}).$$

We will analyze $\partial_{\{-\infty\}}\mathcal{M}$.

Let $\overline{u}_i$, $i \in \mathbb{N}$, be a sequence of curves in $\mathcal{M}$ such that $\overline{u}_i \in \mathcal{M}_{\tau_i}$ and $\lim_{i\to\infty} \tau_i = -\infty$, and let

$$\overline{u}_\infty = (\overline{v}_{-1,1} \cup \cdots \cup \overline{v}_{-1,c}) \cup \overline{v}_1 \cup \overline{v}_2 \cup (\overline{v}_{1,1} \cup \cdots \cup \overline{v}_{1,a})$$

be the limit holomorphic building in order from bottom to top, where $\overline{v}_{-1,j}$, $j = 1, \dots, c$, maps to $\overline{W'}$; $\overline{v}_j$ maps to $\overline{W}_{-\infty,j}$; and $\overline{v}_{1,j}$, $j = 1, \dots, a$ maps to $\overline{W'}$. Sometimes we refer to $\overline{v}_1$ as $\overline{v}_{-1,c+1}$ or $\overline{v}_{1,0}$.

We write $\overline{v}_* = \overline{v}'_* \cup \overline{v}''_*$, $\overline{v}''_* = \overline{v}^\sharp_* \cup \overline{v}^\flat_*$, where:

- $\overline{v}'_*$ is the union of branched covers of a section at infinity;
- $\overline{v}^\sharp_*$ is the union of components that are not in $\overline{v}'_*$ and are asymptotic to some multiple of δ_0 or pass through $\mathfrak{z} = z_\infty^{r_0}\zeta_1^{r_1}\cdots\zeta_l^{r_l}$ with $r_0 > 0$; and
- $\overline{v}^\flat_*$ is the union of the remaining components of $\overline{v}_*$.

Remark 4.5.1. Strictly speaking, in the limit $\tau \to -\infty$, there exist levels between $\overline{v}_1$ and $\overline{v}_2$, i.e., levels that map to $\mathbb{R} \times S^1 \times \overline{S}$. By considerations of n^*, these levels are connectors (i.e., map to $\mathbb{R} \times S^1 \times \{pt\}$) and will be ignored until we consider gluing.

The following are the analogs of Lemmas 4.4.1 and 4.4.2:

Lemma 4.5.2. *If* $\overline{v}'_* \cup \overline{v}^\sharp_* = \varnothing$ *for all levels* $\overline{v}_*$ *of* $\overline{u}_\infty$*, then* $a = c = 0$*;* $I(\overline{v}_1) = 0$*;* $I(\overline{v}_2) = 4g + 2$*; and* $\overline{v}_2$ *is a* $\overline{W}_2$*-curve.*

Proof. Suppose $\overline{v}'_* \cup \overline{v}^\sharp_* = \varnothing$ for all levels $\overline{v}_*$ of $\overline{u}_\infty$. The following are immediate from considerations of n^*:

(1) $\overline{v}_2$ is a $\overline{W}_2$-curve or a degenerate $\overline{W}_2$-curve; and
(2) $\overline{v}_* = \overline{v}''_*$ and $n^*(\overline{v}_*) = 0$ for $* = (-1, j)$, 1, and $(1, j)$.

(1) implies that $I(\overline{v}_2) = 4g + 2$ by Lemma 4.2.7. Since there are $2g$ codimension two gluing conditions between $\overline{v}_1$ and $\overline{v}_2$, it follows that

$$\sum_* I(\overline{v}_*) = 4g + 2, \tag{4.5.1}$$

where the summation is over all the levels $*$. Hence $I(\overline{v}_1) = 0$ and each component $\widetilde{v}$ of $\overline{v}_1$ is a branched cover of a trivial cylinder with possibly empty branch locus. (Here we are assuming without loss of generality that the almost complex structure on $\overline{W}_1$ is $\overline{J}_1$.) On the other hand, each non-fiber component of a degenerate $\overline{W}_2$-curve is of the type $B_{-\infty,2} \times \{q_l\}$ for some $q_l \in \widehat{a}_l$ and has Fredholm index one. Since we may assume that $\widehat{\mathcal{O}}_{2g}$ is disjoint from $\mathbf{a} \times \{1\} \subset \overline{S} \times \{1\}$ by genericity, the gluing condition for $\overline{v}_1$ and $\overline{v}_2$ is not satisfied. □

Lemma 4.5.3. *If* $\overline{v}'_* \cup \overline{v}^\sharp_* \neq \varnothing$ *for some level* $\overline{v}_*$ *of* $\overline{u}_\infty$*, then:*

(1) $p_2 = \deg(\overline{v}'_2) > 0$*;*
(2) $\overline{u}_\infty$ *has no boundary puncture that is removable at* z_∞*;*
(3) $\overline{u}_\infty$ *has no fiber components and no components of* $\overline{v}''_*$ *that intersect the interior of a section at infinity; and*

(4) *$\overline{v}''_2$ is the union of $2g-p_2$ copies of $B_{-\infty,2}\times\{pt\}$, and $I(\overline{v}_2)=2g$.*

Proof. (1)–(3) are analogs of Lemma 4.4.2 and (4) is a consequence of Lemma 4.2.8. □

The following is the analog of Lemma 4.4.4.

Lemma 4.5.4. *If $\overline{v}'_*\cup\overline{v}^\sharp_*\neq\varnothing$ for some level $\overline{v}_*$, then $\overline{v}'_2\neq\varnothing$ and $\overline{v}''_2$ is a union of components $B_{-\infty,2}\times\{pt\}$, and $\overline{u}_\infty$ contains one of the following subbuildings:*

(1_i) *A 2-level building consisting of $\overline{v}^\sharp_1$ with $I=i+(\deg(\overline{v}^\sharp_1)-1)$, $i=2,3$, which passes through $\mathfrak{z}$ with $r_0=1$; and $\overline{v}'_2=\sigma_\infty^{-\infty,2}$ with $I=1$.*

(2) *A 2-level building consisting of $\overline{v}^\sharp_1$ with $I=4+(\deg(\overline{v}^\sharp_1)-2)$ which passes through $\mathfrak{z}$ with $r_0=2$; and $\overline{v}'_2$ with $I=2$ which is a branched double cover of $\sigma_\infty^{-\infty,2}$.*

(3_i) *A 4-level building consisting of $\overline{v}^\sharp_{1,1}$ with $I=1$ or 2 and a negative end at δ_0; $\overline{v}'_1=\sigma_\infty^{-\infty,1}$; $\overline{v}'_2=\sigma_\infty^{-\infty,2}$; and a cylinder of $\overline{v}^\sharp_{-1,1}$ from δ_0 to h or e with $I=1$ or 2.*

(4) *A 4-level building consisting of $\overline{v}^\sharp_{1,1}$ with $I=1$ and a negative end at δ_0^2; $\overline{v}'_1=\sigma_\infty^{-\infty,1}$; a cylinder of $\overline{v}^\sharp_1$ from δ_0 to h or e with $I=1,2$; $\overline{v}'_2=\sigma_\infty^{-\infty,2}$; a cylinder of $\overline{v}^\sharp_{-1,1}$ from δ_0 to h with $I=1$; and a cylinder of $\overline{v}''_{-1,1}$ from h or e to e with $I=1,0$.*

(5) *A 4-level building consisting of $\overline{v}^\sharp_{1,1}$ with $I=1$ and a negative end at δ_0^2; $\overline{v}'_1$ with $I=0$ which is a branched double cover of $\sigma_\infty^{-\infty,1}$; $\overline{v}'_2$ with $I=2$ which is a branched double cover of $\sigma_\infty^{-\infty,2}$; and two cylinder components of $\cup_{j=1}^c\overline{v}^\sharp_{-1,j}$ from δ_0 to h or e, each with $I=1$ or 2.*

(6) *A 5-level building consisting of $\overline{v}^\sharp_{1,2}$ with $I=1$ and a negative end at δ_0^2; $\overline{v}'_{1,1}=\mathbb{R}\times\delta_0$; a component of $\overline{v}^\sharp_{1,1}$ with $I=1$ which is a cylinder from δ_0 to h; $\overline{v}'_1=\sigma_\infty^{-\infty,1}$; $\overline{v}'_2=\sigma_\infty^{-\infty,2}$; and a cylinder of $\overline{v}^\sharp_{-1,1}$ from δ_0 to h with $I=1$.*

We are omitting levels which are branched covers of trivial cylinders. Moreover, each gluing condition reduces the sum of ECH indices by 2.

See Figure 17.

Proof. The proof is similar to that of Lemma 4.4.4 and we only give a sketch.

We have $\overline{v}'_2\neq\varnothing$ by Lemma 4.5.3(1). By Lemma 4.2.8, $\overline{v}''_2$ is a union of components $B_{-\infty,2}\times\{q_l\}$, $q_l\in\widehat{a}_l$, and $I(\overline{v}_2)=2g$.

First suppose that $\overline{v}'_1=\varnothing$. Then $\overline{v}^\sharp_1$ passes through $((0,\frac{3}{2}),z_\infty)$ and $\pi_{B_{-\infty,1}}\circ\overline{v}^\sharp_1$ has a k-fold branch point at $(0,\frac{3}{2})$ for some $k>0$, where $\pi_{B_{-\infty,1}}:\overline{W}_{-\infty,1}\to B_{-\infty,1}$ is the projection along $\overline{S}$. The k-fold branch point at $((0,\frac{3}{2}),z_\infty)$ contributes $2k$ towards the Fredholm index and each of the remaining $2g-k$ intersection points with $\{(0,\frac{3}{2})\}\times\overline{S}$ passes through some $\widehat{a}_l$ and contributes $+1$ towards the Fredholm index. Hence $I(\overline{v}_1)\geq\operatorname{ind}(\overline{v}_1)\geq 2k+(2g-k)$. Since the gluing



FIGURE 17. Schematic diagrams for the possible types of degenerations. The dotted lines indicate the section at infinity. The quantities -2 and -4 along the intersections of $\overline{v}_1$ and $\overline{v}_2$ indicate the reduction of I due the gluing conditions. For simplicity we are assuming that $\deg \overline{v}_1^\sharp = 1$ in (1_i) and $\deg \overline{v}_1^\sharp = 2$ in (2).

conditions contribute $-4g$, the total ECH contribution of the three terms is k. This implies that $k \leq 2$, giving us (1_i), $i = 2, 3$, or (2).

Next suppose that $\overline{v}_1' \neq \varnothing$. Then $I(\overline{v}_1') = 0$, $I(\overline{v}_2') = k$, and the gluing conditions give $-2k$ for some $k > 0$. Hence (3_i), $i = 1, 2$, and (4)–(6) follow from enumerating all the possibilities, subject to the condition that h appears only once at the negative end of $\overline{v}_{-1,1}$. $\square$

Lemma 4.5.5. *If $m \gg 0$, $\overline{u}_\infty \in \partial_{\{-\infty\}}\mathcal{M}$, $\overline{v}_*' \cup \overline{v}_*^\sharp \neq \varnothing$ for some level $\overline{v}_*$, then the only possibility is Case (1_3).*

Proof. Cases (1_2) and (2). We will treat Case (1_2); Case (2) is similar. We apply the usual rescaling argument with $m \gg 0$ fixed to obtain a function $w_2 : B_{-\infty,2} \to \mathbb{CP}^1$ satisfying the following:

(i_2) $w_2(0) = \infty$;
(ii_2) $w_2(\frac{i}{2}) = 0$;
(iii_2) $w_2(\partial B_{-\infty,2}) \subset \mathbb{R}^+ \cdot e^{i\phi(\overline{a}_{k,l})}$ for some (k, l);
(iv_2) w_2 is a biholomorphism away from $\partial B_{-\infty,2}$.

We now observe that w_2 is uniquely determined by (i_2)–(iv_2), up to multiplication by a positive real constant. This implies that:

(v_2) w_2 maps the asymptotic marker ∂_y at $0 \in B_{-\infty,2}$ to the asymptotic marker $\dot{\mathcal{R}}_{\pi+\phi(\overline{a}_{k,l})}(\infty)$.

Here $z = x + iy$ is the complex coordinate on $B_{-\infty,2}$. (v_2) translates into an asymptotic condition for $\overline{v}_1^\sharp$ at $((0, \frac{3}{2}), z_\infty)$. Hence $I(\overline{v}_1^\sharp)$ is at least 3.

Cases (3$_i$), (4)–(6). We will treat Case (3$_i$). The rescaling argument gives $w_2 \cup w_1$, where w_2 is as before and $w_1 : cl(B_{-\infty,1}) \to \mathbb{CP}^1$ satisfies the following:

- (i$_1$) $w_1(0, \frac{3}{2}) = 0$ and $w_1(+\infty) = +\infty$;
- (ii$_1$) w_1 is a biholomorphism.

(iv$_2$) implies the following asymptotic condition for w_1:

- (iii$_1$) w_1 maps the marker ∂_s at $(0, \frac{3}{2})$ to the marker $\dot{\mathcal{R}}_{\pi+\phi(\overline{a}_{k,l})}(0)$.

Hence w_1 is uniquely determined by (i$_1$)–(iii$_1$) up to multiplication by a positive real constant.

As a consequence of the uniqueness of w_1 up to multiplication by a positive real constant, the following are uniquely determined:

- (a) the asymptotic eigenfunction of $\overline{v}^\sharp_{1,1}$ at the negative end δ_0;
- (b) the asymptotic eigenfunction of $\overline{v}^\sharp_{-1,1}$ at the positive end δ_0;
- (c) the next higher order term in the asymptotic expansion of $\overline{v}^\sharp_{-1,1}$ at the positive end δ_0.

(a) is determined by the image of the asymptotic marker $\dot{\mathcal{L}}_{3/2}(+\infty)$ at $+\infty \in cl(B_{-\infty,1})$, (b) by the radial ray that contains $w_1(-\infty)$, and (c) by the image of the asymptotic marker $\dot{\mathcal{L}}_{3/2}(-\infty)$ at $-\infty \in cl(B_{-\infty,1})$. (a)–(c) give rise to one constraint on $\overline{v}^\sharp_{1,1}$ and two constraints on $\overline{v}^\sharp_{-1,1}$. Hence $I(\overline{v}^\sharp_{1,1}) \geq 2$ and $I(\overline{v}^\sharp_{-1,1}) \geq 3$, which is a contradiction. □

Proof of Lemma 4.3.3. This is a combination of Lemmas 4.5.2 and 4.5.5. □

Proof of Lemma 4.3.4. We claim that the mod 2 count of $\mathcal{M}^{I=4g+2,n^*=m}_{\overline{J}_{-\infty,2}}(\mathfrak{z};\overline{\mathfrak{m}}(-\infty))$ is 1 when $\mathfrak{z}$ is generic and $r_0 = 0$. This is proved by reducing to the calculation of Theorem 2.3.3 as follows: Degenerate $B_{-\infty,2}$ into a sphere $B_{-\infty,21}$ and a disk $B_{-\infty,22}$ which are identified at one point and degenerate $\overline{W}_{-\infty,2} = B_{-\infty,2} \times \overline{S}$ into $(B_{-\infty,21} \times \overline{S}) \cup (B_{-\infty,22} \times \overline{S})$. We assume that the marked point is in $B_{-\infty,21} \times \overline{S}$. Then a curve $\overline{v}_2 \in \mathcal{M}^{I=4g+2,n^*=m}_{\overline{J}_{-\infty,2}}(\mathfrak{z};\overline{\mathfrak{m}}(-\infty))$, degenerates into a pair $(\overline{v}_{21}, \overline{v}_{22})$, where $\overline{v}_{2i}$, $i = 1, 2$, maps to $B_{-\infty,2i} \times \overline{S}$ and $\overline{v}_{22}$ is a union of constant sections $B_{-\infty,22} \times \{q_l\}$, where $q_l \in \widehat{a}_l$. Hence $\overline{v}_{21}$ is a curve in $B_{-\infty,21} \times \overline{S}$ with exactly the same type of constraints as in Theorem 2.3.3. This implies the claim.

In the rest of the proof we discuss how to glue pairs $(\overline{v}_1, \overline{v}_2) \in A_2$. For simplicity we work with $\overline{J}_{-\infty,1}$ instead of $\overline{J}^\Diamond_{-\infty,1}(\varepsilon, \delta, \mathfrak{p}(-\infty)))$. Since $I(\overline{v}_1) = 0$, each component $\widetilde{v}$ of $\overline{v}_1$ is a branched cover of a trivial cylinder with possibly empty branch locus. If $\widetilde{v}$ is simple, then it glues to $\overline{v}_2$ in the usual manner. On the other hand, if $\widetilde{v}$ is multiply covered, then it is the cylinder over an elliptic orbit which is the same type as e by (††). Suppose $\widetilde{v}$ covers $\mathbb{R} \times e$ with multiplicity k. By the partition conditions, the incoming partition is (k) and the outgoing partition is $(1, \dots, 1)$. The outgoing partition condition applies to the positive end of $\widetilde{v}$ and the incoming partition condition applies to the negative end of $\widetilde{v}$; this is the same partition condition as that of $\overline{u}_i$ for $i \gg 0$ where $\overline{u}_i \to \overline{u}_\infty$. Hence $\widetilde{v}$ has k positive ends, 1 negative end, and $\mathfrak{b} \geq k - 1$ branch points (in the sense of Definition 3.4.22).

For consistency with $\overline{v}_2$, we must have a branch point at $(0,\frac{3}{2})$ which contributes $k-1$ towards $\mathfrak{b}$.

Next we define $\operatorname{ind}'(\widetilde{v})$ as the Fredholm index when:

- $\widetilde{v}$ is viewed as a curve in the fibration
$$\overline{W}_{-\infty,1}-(\{(0,3/2)\}\times\overline{S})\to B_{-\infty,1}-\{(0,3/2)\};$$
- $\{(0,\frac{3}{2})\}\times\overline{S}$ is in one-to-one correspondence with a Morse-Bott family of orbits and the orbits of $\{(0,\frac{3}{2})\}\times\overline{S}$ are viewed as orbits of the positive end;
- the end $\zeta\in\{(0,\frac{3}{2})\}\times\overline{S}$ of $\widetilde{v}$ is viewed as an elliptic orbit with Conley-Zehnder index 1.

By the usual Fredholm index calculation, we obtain $\operatorname{ind}'(\widetilde{v})\geq 0$, with equality if and only if $\mathfrak{b}=k-1$. We can similarly define ind' for $\overline{v}_2$. Then $\operatorname{ind}'(\overline{v}_2)=2$. When we use ind' instead of ind, the gluing conditions are 0-dimensional instead of $4g$-dimensional. This implies that $\operatorname{ind}'(\widetilde{v})=0$, $\mathfrak{b}=k-1$, and $\widetilde{v}$ has no branch points besides $(0,\frac{3}{2})$.

Finally we discuss the automatic transversality of $\widetilde{v}$. Recall from [We3, Theorem 1] that automatic transversality holds if

$$\operatorname{ind}'(\widetilde{v})\geq 2g+\#\Gamma_0-1, \tag{4.5.2}$$

where g is the genus of $\widetilde{v}$ and $\#\Gamma_0$ is the number of punctures (of $\widetilde{v}$ with $(0,\frac{3}{2})$ removed) with even Conley-Zehnder index. Since the right-hand side of Equation (4.5.2) is equal to -1, automatic transversality holds and $\widetilde{v}$ glues in the usual manner (without any concerns of inserting branch-covered cylinders) to $\overline{v}_2$. This implies the lemma. □

Sketch of proof of Lemma 4.3.5. We use some considerations of Section I.7.13. For simplicity we assume we are gluing degree one curves. Fix $\gamma,\gamma'\in\widehat{\mathcal{O}}_1$, $k\in\{1,\dots,2g\}$, and $l\in\{0,1\}$. We consider the gluing parameter space

$$\mathfrak{P}_{k,l}:=\coprod_{\mathfrak{z}_0\in\overline{\mathfrak{u}}_{k,l}}\left((-\infty,-r]\times\mathcal{M}_1(\mathfrak{z}_0)\times\mathcal{M}_2(\mathfrak{z}_0)\right),$$

where $\mathfrak{u}_{k,l}\subset\vec{a}_{k,l}$ is a small open interval containing z_∞,

$$\mathcal{M}_1(\mathfrak{z}_0):=\mathcal{M}^{I=3,n^*=m}_{\overline{J}^{\diamond}_{-\infty,1}(\varepsilon,\delta,\mathfrak{p}(-\infty))}(\gamma,\gamma',\mathfrak{z}_0),$$
$$\mathcal{M}_2(\mathfrak{z}_0):=\mathcal{M}^{I=1,n^*=0,ext}_{\overline{J}_{-\infty,2}}(\mathfrak{z}_0),$$

ext means the boundary of the holomorphic curve is mapped to $L^{-\infty,2}_{\overline{a}_k\cup\vec{a}_{k,l}}$, and $(-\infty,-r]$ is viewed as a subset of $(-\infty,\infty)$ with parameter τ. For each $\mathfrak{z}_0\in\mathfrak{u}_{k,l}$ there is a covering map

$$\pi_{\mathfrak{z}_0}:\mathcal{M}_1(\mathfrak{z}_0)\to S^1=\mathbb{R}/2\pi\mathbb{Z}$$

which maps $\overline{v}_1$ to its asymptotic marker at $\mathfrak{z}_0$ and is continuous in $\mathfrak{z}_0$. Note that $\mathcal{M}_2(\mathfrak{z}_0)$ consists of a single point $B_{-\infty,2}\times\{\mathfrak{z}_0\}$.

Let

$$G_{k,l}:\mathfrak{P}_{k,l}\to\coprod_{\tau\in(-\infty,r]}\mathcal{M}^{I=2,n^*=m,ext}_{\{\overline{J}^{\diamondsuit}_\tau(\varepsilon,\delta,\mathfrak{p}(\tau))\}}(\gamma,\gamma'),\quad \mathfrak{d}=(\tau,\overline{v}_1,\overline{v}_2)\mapsto\overline{u}(\mathfrak{d})$$

be the gluing map, defined as usual. Also let $\mathfrak{P}'_{k,l}\subset\mathfrak{P}_{k,l}$ be the subset of gluing parameters $\mathfrak{d}$ such that $G_{k,l}(\mathfrak{d})\in\displaystyle\coprod_{\tau\in(-\infty,r]}\mathcal{M}^{I=2,n^*=m}_{\{\overline{J}^{\diamondsuit}_\tau(\varepsilon,\delta,\mathfrak{p}(\tau))\}}(\gamma,\gamma')$.

Let $\widetilde{\mathcal{N}}$ be the space of holomorphic maps $w_2:B_{-\infty,2}=\overline{\mathbb{D}}\to\mathbb{CP}^1$ satisfying (i_2) and (iv_2) of Lemma 4.5.5 and

(iii'_2) $w_2(\partial B_{-\infty,2})\subset\mathbb{R}\cdot e^{i\phi(\overline{a}_{k,l})}$ for some (k,l).

For simplicity we assume that $\phi(\overline{a}_{k,l})=0$. We leave it to the reader to verify the following:

Claim 4.5.6. $\widetilde{\mathcal{N}}=\{w_2(z)=az+b+\frac{\overline{a}}{z}\mid a\in\mathbb{C}^\times,b\in\mathbb{R}\}$. *Hence* $\dim\widetilde{\mathcal{N}}=3$ *and* $\widetilde{\mathcal{N}}$ *admits an* $\mathbb{R}^\times$*-action which is multiplication by* $c\in\mathbb{R}^\times$.

For $\tau_0\ll 0$, the restrictions of $\overline{u}(\mathfrak{d})$ to a neighborhood of the section at infinity are approximated by elements of $\widetilde{\mathcal{N}}$. More precisely, for $\tau_0\ll 0$ we define:

$$g_{\tau_0,k,l}:\mathfrak{P}_{k,l}\cap\{\tau=\tau_0\}\to\widetilde{\mathcal{N}},$$

as follows: Given $\mathfrak{d}\in\mathfrak{P}_{k,l}\cap\{\tau=\tau_0\}$, restrict $G_{k,l}(\mathfrak{d})$ to a neighborhood of the section at infinity so that the domain of $G_{k,l}(\mathfrak{d})$ is $\overline{\mathbb{D}}^\varepsilon:=\{\varepsilon\leq|z|\leq 1\}\subset\overline{\mathbb{D}}$ for $\varepsilon>0$ small. Let us write $\pi_{\overline{S}}\circ G_{k,l}(\mathfrak{d})|_{\overline{\mathbb{D}}^\varepsilon}$ as a Laurent series $\sum_{i=-\infty}^{\infty}c_i(\mathfrak{d})z^i$. Then we set

$$g_{\tau_0,k,l}(\mathfrak{d})=\overline{c_{-1}(\mathfrak{d})}\cdot z+\mathrm{Re}(c_0(\mathfrak{d}))+\frac{c_{-1}(\mathfrak{d})}{z}.$$

The definition makes sense in view of the following (proof omitted):

Claim 4.5.7.

(1) $\displaystyle\lim_{\tau\to-\infty}\frac{c_i(\mathfrak{d})}{c_{-1}(\mathfrak{d})}=0$ *if* $i>1$ *or* $i<-1$ *and* $\displaystyle\lim_{\tau\to-\infty}\frac{\overline{c_1(\mathfrak{d})}}{c_{-1}(\mathfrak{d})}=1$, *where the convergence is uniform in* $\displaystyle\coprod_{\mathfrak{z}_0\in\overline{\mathfrak{u}}_{k,l}}(\mathcal{M}_1(\mathfrak{z}_0)\times\mathcal{M}_2(\mathfrak{z}_0))$.

(2) *If* $\mathrm{Im}(\overline{v}_2)=\{\mathfrak{z}_0\}\neq\{z_\infty\}$, *then* $\displaystyle\lim_{\tau_0\to-\infty}\frac{c_{-1}(\mathfrak{d})}{c_0(\mathfrak{d})}=0$ *and* $\displaystyle\lim_{\tau_0\to-\infty}\frac{\mathrm{Im}(c_0(\mathfrak{d}))}{\mathrm{Re}(c_0(\mathfrak{d}))}=0$, *where the convergence is uniform in* $\mathcal{M}_1(\mathfrak{z}_0)$.

Now define the evaluation map:

$$ev_{\tau_0,k,l}:\mathfrak{P}_{k,l}\cap\{\tau=\tau_0\}\to\mathbb{C},$$

which sends $\mathfrak{d}$ to $g_{\tau_0,k,l}(\mathfrak{d})(\frac{i}{2})$. One can verify that the local degree of $ev_{\tau_0,k,l}$ near $0\in\mathbb{C}$ is $2\cdot\deg\pi_{\mathfrak{z}_0}$ and the local degree of $(ev_{\tau_0,k,l})|_{\mathfrak{P}'_{k,l}\cap\{\tau=\tau_0\}}$ is $\deg\pi_{\mathfrak{z}_0}$. We will informally say that "$\mathcal{M}$ has $\deg\pi_{\mathfrak{z}_0}$ ends near $\mathfrak{P}'_{k,l}$".

Finally observe that an element of $\mathcal{M}_2(z_\infty)$ can be viewed as a map to $\overline{a}_{i,0}$ or to $\overline{a}_{i,1}$. Hence $\mathcal{M}$ has $\deg\pi_{\mathfrak{z}_0}$ ends near $\mathfrak{P}'_{k,l}$ and $\deg\pi_{\mathfrak{z}_0}$ ends near $\mathfrak{P}'_{k,1-l}$ for a total of $2\cdot\deg\pi_{\mathfrak{z}_0}$ ends. This proves Lemma 4.3.5. $\square$

4.6. **Breaking in the middle.** In this subsection we study the limit of holomorphic maps to $\overline{W}_\tau$ as $\tau \to T'$ for some $T' \in (-\infty, \infty)$. This will prove Lemma 4.3.6.

We assume that $m \gg 0$; $\varepsilon, \delta > 0$ are sufficiently small; and $\{\overline{J}_\tau\} \in \overline{\mathcal{I}}^{reg}$ and $\{\overline{J}_\tau^\Diamond(\varepsilon, \delta, \mathfrak{p}(\tau))\}$ satisfy Lemma 4.2.13. Fix $\gamma, \gamma' \in \widehat{\mathcal{O}}_{2g}$ and let

$$\mathcal{M} = \mathcal{M}^{I=2, n^*=m}_{\{\overline{J}_\tau^\Diamond(\varepsilon,\delta,\mathfrak{p}(\tau))\}}(\gamma, \gamma'; \overline{\mathfrak{m}}), \quad \mathcal{M}_\tau = \mathcal{M}^{I=2, n^*=m}_{\overline{J}_\tau^\Diamond(\varepsilon,\delta,\mathfrak{p}(\tau))}(\gamma, \gamma'; \overline{\mathfrak{m}}).$$

We will analyze $\partial_{(-\infty,\infty)}\mathcal{M}$.

Let $\overline{u}_i$, $i \in \mathbb{N}$, be a sequence of curves in $\mathcal{M}$ such that $\overline{u}_i \in \mathcal{M}_{\tau_i}$ and $\lim_{i\to\infty} \tau_i = T'$, and let

$$\overline{u}_\infty = (\overline{v}_{-1,1} \cup \dots \cup \overline{v}_{-1,c}) \cup \overline{v}_0 \cup (\overline{v}_{1,1} \cup \dots \cup \overline{v}_{1,a})$$

be the limit holomorphic building in order from bottom to top, where $\overline{v}_{-1,j}$ and $\overline{v}_{1,j}$ map to $\overline{W'}$ and $\overline{v}_0$ maps to $\overline{W}_{T'}$. Sometimes we refer to $\overline{v}_0$ as $\overline{v}_{-1,c+1}$ or $\overline{v}_{1,0}$.

The following is the analog of Lemma 4.4.1, and is stated without proof.

Lemma 4.6.1. *If $\overline{v}'_* \cup \overline{v}^\sharp_* = \varnothing$ for all levels $\overline{v}_*$ of $\overline{u}_\infty$, then $\overline{u}_\infty$ is one of the following:*

1. *$a = 0$, $c = 1$; $\overline{v}_0$ is a $\overline{W}_{T'}$-curve with $I = 1$ which passes through $\overline{\mathfrak{m}}(T')$; and $\overline{v}_{-1,1}$ is a W'-curve with $I = 1$; or*
2. *$a = 1$, $c = 0$; $\overline{v}_{1,1}$ is a W'-curve with $I = 1$; and $\overline{v}_0$ is a $\overline{W}_{T'}$-curve with $I = 1$ which passes through $\overline{\mathfrak{m}}(T')$.*

Here either $T' \in \mathcal{T}_2$ and a component of $\overline{v}_0$ is in

$$\mathcal{M}^{\dagger, s, irr, \mathrm{ind}=1, n^*=m}_{\overline{J}_{T'}^\Diamond(\varepsilon,\delta,\mathfrak{p}(T'))}(\delta_0^p\gamma_1, \delta_0^q\gamma_2; \overline{\mathfrak{m}}(T'))$$

from Lemma 4.2.13(2), for some $\delta_0^p\gamma_1$, $\delta_0^q\gamma_2$; or $T' \in \mathcal{T}_1$ and a component of $\overline{v}_0$ does not pass through $\overline{\mathfrak{m}}(T')$ but is in

$$\mathcal{M}^{\dagger, s, irr, \mathrm{ind}=-1, n^*=0}_{\overline{J}_{T'}^\Diamond(\varepsilon,\delta,\mathfrak{p}(T'))}(\delta_0^p\gamma_1, \delta_0^q\gamma_2)$$

from Lemma 4.2.13(1).

The following is the analog of Lemma 4.4.2 and is stated without proof.

Lemma 4.6.2. *If $\overline{v}'_* \cup \overline{v}^\sharp_* \neq \varnothing$ for some level $\overline{v}_*$ of $\overline{u}_\infty$, then:*

1. *$p_0 = \deg(\overline{v}'_0) > 0$;*
2. *some $\overline{v}_{1,j_0}$, $j_0 > 0$, has a negative end $\mathcal{E}_-$ that limits to δ_0^p for some $p > 0$ and satisfies $n^*(\mathcal{E}_-) \geq m - p$;*
3. *$\overline{u}_\infty$ has no boundary puncture that is removable at z_∞;*
4. *$\overline{u}_\infty$ has no fiber components and no components of $\overline{v}''_*$ that intersect the interior of a section at infinity;*
5. *each component of $\overline{v}^\sharp_{-1,j}$, $0 \leq j \leq c$, is an $n^* = 1$, $I = 1$ or 2 cylinder from δ_0 to h or e which is contained in $\mathbb{R} \times (\overline{N} - N)$;*
6. *h appears at most once at the negative end of $\overline{v}^\sharp_{-1,1}$.*

The following is the analog of Lemma 4.4.4 and is stated without proof.

Lemma 4.6.3. *If $\overline{v}'_* \cup \overline{v}^\sharp_* \neq \varnothing$ for some level $\overline{v}_*$, then $\overline{v}'_0 \neq \varnothing$ and $\overline{u}_\infty$ contains one of the following subbuildings, subject to two conditions:*

- *the sum of the ECH indices of the components of the subbuildings is at most 3;*
- *the total multiplicity of h at the negative end of $\overline{v}^\sharp_{-1,1}$ is at most 1.*

(1_i) *A 3-level building consisting of $\overline{v}^\sharp_{1,1}$ with $I = i$, $i = 1, 2$, and a negative end $\delta_0\gamma'$; $\overline{v}'_0 = \sigma^\tau_\infty$; and a cylinder component of $\overline{v}^\sharp_{-1,1}$ with $I = 1$ or 2 from δ_0 to h or e.*

(2_i) *A 4-level building consisting of $\overline{v}^\sharp_{1,2}$ with $I = i$, $i = 1, 2$, and a negative end $\delta_0^2\gamma'$; $\overline{v}'_{1,1} = \mathbb{R} \times \delta_0$; a cylinder component of $\overline{v}^\sharp_{1,1}$ with $I = 1$ or 2 from δ_0 to h or e; $\overline{v}'_0 = \sigma^\tau_\infty$; and a cylinder component of $\overline{v}^\sharp_{-1,1}$ with $I = 1$ or 2 from δ_0 to h or e.*

(3_i) *A 3-level building consisting of $\overline{v}^\sharp_{1,1}$ with $I = i$, $i = 1, 2$, and a negative end $\delta_0^2\gamma'$; $\overline{v}'_0 = \sigma^\tau_\infty$; a component of $\overline{v}^\sharp_0$ with $I = 0$ or 1 from δ_0 to h or e; and a cylinder component of $\overline{v}^\sharp_{-1,1}$ with $I = 1$ or 2 from δ_0 to h or e.*

(4_i) *A 3-level building consisting of $\overline{v}^\sharp_{1,1}$ with $I = i$, $i = 1, 2$, and a negative end $\delta_0^2\gamma'$; $\overline{v}'_0$ with $I = -2$ which is a degree 2 branched cover of σ^τ_∞; and two cylinder components of $\cup_{j=1}^c \overline{v}^\sharp_{-1,j}$ from δ_0 to h or e, each with $I = 1$ or 2.*

(5) *A 5-level building consisting of $\overline{v}^\sharp_{1,3}$ with $I = 1$ and a negative end $\delta_0^3\gamma'$; $\overline{v}'_{1,2}$ which is a degree 2 branched cover of $\mathbb{R} \times \delta_0$; a cylinder component of $\overline{v}^\sharp_{1,2}$ with $I = 1$ from δ_0 to h; $\overline{v}'_{1,1} = \mathbb{R} \times \delta_0$; a cylinder component of $\overline{v}^\sharp_{1,1}$ with $I = 1$ from δ_0 to h; $\overline{v}'_0 = \sigma^\tau_\infty$; and a cylinder component of $\overline{v}^\sharp_{-1,1}$ with $I = 1$ from δ_0 to h.*

(6) *A 4-level building consisting of $\overline{v}^\sharp_{1,2}$ with $I = 1$ and a negative end $\delta_0^3\gamma'$; $\overline{v}'_{1,1} = \mathbb{R} \times \delta_0$; two cylinder components of $\overline{v}^\sharp_{1,1}$ from δ_0 to h, each with $I = 1$; $\overline{v}'_0 = \sigma^\tau_\infty$; and a cylinder component of $\overline{v}^\sharp_{-1,1}$ with $I = 1$ from δ_0 to h.*

(7) *A 4-level building consisting of $\overline{v}^\sharp_{1,2}$ with $I = 1$ and a negative end $\delta_0^3\gamma'$; $\overline{v}'_{1,1}$ which is a degree 2 branched cover of $\mathbb{R} \times \delta_0$; a cylinder component of $\overline{v}^\sharp_{1,1}$ with $I = 1$ or 2 from δ_0 to h or e; $\overline{v}'_0 = \sigma^\tau_\infty$; a component of $\overline{v}^\sharp_0$ with $I = 0$ or 1 from δ_0 to h or e; and a cylinder component of $\overline{v}^\sharp_{-1,1}$ with $I = 1$ or 2 from δ_0 to h or e.*

(8) *A 4-level building consisting of $\overline{v}^\sharp_{1,2}$ with $I = 1$ and a negative end $\delta_0^3\gamma'$; $\overline{v}'_{1,1}$ which is a degree 2 branched cover of $\mathbb{R} \times \delta_0$; a cylinder component of $\overline{v}^\sharp_{1,1}$ with $I = 1$ from δ_0 to h; $\overline{v}'_0$ with $I = -2$ which is a degree 2 branched cover of σ^τ_∞; and two cylinder components of $\cup_{j=1}^c \overline{v}^\sharp_{-1,j}$ from δ_0 to h or e, each with $I = 1$ or 2.*

(9) *A* 3*-level building consisting of* $\overline{v}^\sharp_{1,1}$ *with* $I=1$ *and a negative end* $\delta_0^3\gamma'$*;* $\overline{v}'_0=\sigma^\tau_\infty$*; two components of* $\overline{v}^\sharp_0$ *from* δ_0 *to* h *or* e*, each with* $I=0$ *or* 1*; and a cylinder component of* $\overline{v}^\sharp_{-1,1}$ *with* $I=1$ *or* 2 *from* δ_0 *to* h *or* e*.*

(10) *A* 3*-level building consisting of* $\overline{v}^\sharp_{1,1}$ *with* $I=1$ *and a negative end* $\delta_0^3\gamma'$*;* $\overline{v}'_0$ *with* $I=-2$ *which is a degree* 2 *branched cover of* σ^τ_∞*; a component of* $\overline{v}^\sharp_0$ *with* $I=0$ *or* 1 *from* δ_0 *to* h *or* e*; and two cylinder components of* $\cup_{j=1}^c\overline{v}^\sharp_{-1,j}$ *from* δ_0 *to* h *or* e*, each with* $I=1$ *or* 2*.*

(11) *A* 3*-level building consisting of* $\overline{v}^\sharp_{1,1}$ *with* $I=1$ *and a negative end* $\delta_0^3\gamma'$*;* $\overline{v}'_0$ *with* $I=-3$ *which is a degree* 3 *branched cover of* σ^τ_∞*; and three cylinder components of* $\cup_{j=1}^c\overline{v}^\sharp_{-1,j}$ *from* δ_0 *to* h *or* e*, each with* $I=1$ *or* 2*.*

We are omitting levels which are connectors.

See Figure 18. We will write $(1_{i,e})$, $(1_{i,h})$, etc. to indicate that we are in Case (1_i) and the negative ends of the lowest level are e, h, etc.



(1_i) (2_i) (3_i) (4_i) (5) (6)

(7) (8) (9) (10) (11)

FIGURE 18. Schematic diagrams for the possible types of degenerations. For simplicity we have drawn only one level $\overline{v}_{-1,1}$ to indicate the cylindrical components of $\cup_{j=1}^c\overline{v}^\sharp_{-1,j}$ from δ_0 to h or e.

Lemma 4.6.4. *If $m \gg 0$, then there is no $\overline{u}_\infty \in \partial_{(-\infty,+\infty)}\mathcal{M}$ such that $\overline{v}'_* \cup \overline{v}^\sharp_* \neq \varnothing$ for some level $\overline{v}_*$.*

Proof. Cases (1_i), (2_i), (3_i), (5), (6), (7) and (9). We will treat Case (1_i); the rest of the cases are similar. The key observation here is that $\deg(\overline{v}'_0) = 1$ and $\overline{v}^\sharp_{-1,1}$ is a cylinder from δ_0 to h or e. Applying the usual rescaling argument with $m \gg 0$ fixed, we obtain a holomorphic map $w_0 : cl(B_{T'}) \to \mathbb{CP}^1$ which satisfies the following:

(i) $w_0(+\infty) = \infty$ and $w_0(\overline{\mathfrak{m}}^b(T')) = 0$;
(ii) $w_0(s,t) \in int(\mathcal{R}_{\eta(t)})$ for all $(s,t) \in \partial B_{T'}$;
(iii) $\deg(w_0) = 1$ away from $\mathfrak{S}(\overline{a}_{k,l}, \overline{h}(\overline{a}_{k,l}))$ for some (k,l).

Here $\eta(t) = \phi_0 + \frac{\pi}{m}(t-1)$, where ϕ_0 is the ϕ-coordinate of $\overline{a}_{k,l}$. (Recall that we are projecting to $\pi_{D_{\rho_0}}$ using balanced coordinates.)

We now observe that $w_0 : cl(B_{T'}) \to \mathbb{CP}^1$ is uniquely determined by (i)–(iii), up to multiplication by a positive real constant; this is argued in the same way as in Lemma 3.7.9. Using the same method as in Case (4_i) of Lemma 4.5.5, we then obtain that $I(\overline{v}^\sharp_{1,1}) \geq 2$ and $I(\overline{v}^\sharp_{-1,1}) \geq 3$. This is a contradiction.

Cases (4_i), (8), (10), and (11). We will treat Case (4_i). In this case we obtain a holomorphic map $w_0 : \Sigma_0 \to \mathbb{CP}^1$ and a branched double cover $\pi_0 : \Sigma_0 \to cl(B_{T'})$ such that:

(i) $w_0(z_0) = \infty$ for some $z_0 \in \pi_0^{-1}(+\infty)$;
(ii) $w_0(z_1) = 0$ for some $z_1 \in \pi_0^{-1}(\overline{\mathfrak{m}}^b(T'))$;
(iii) $w_0(\pi_0^{-1}(s,t)) \in int(\mathcal{R}_{\eta_1(t)} \cup \mathcal{R}_{\eta_2(t)})$ for all $(s,t) \in \partial B_{T'}$;
(iv) $\deg(w_0) = 1$ away from $\mathfrak{S}(\overline{a}_{i_k,j_k}, \overline{h}(\overline{a}_{i_k,j_k}))$, $k = 1,2$, for some (i_k, j_k).

Here $\eta_k(t) = \phi_k + \frac{\pi}{m}(t-1)$, $k = 1,2$, where ϕ_k is the ϕ-coordinate of $\overline{a}_{i_k,j_k}$. By a Fredholm index calculation, one verifies that $\pi_0^{-1}(+\infty)$ consists of one point and that π_0 has one simple branch point $p \in B_{T'}$. For each choice of branch point p, there is a unique complex structure on Σ_0 and a unique holomorphic map w_0, up to multiplication by a positive real constant. (The argument is the same as before.) We now have one constraint for the end of $\overline{v}^\sharp_{1,1}$ at δ_0, two constraints for each end of $\overline{v}^\sharp_{-1,1}$ at δ_0, and two degrees of freedom from the position of the branch point p. In total we have:

$$I(\overline{v}^\sharp_{1,1}) + I(\overline{v}'_0) + I(\overline{v}^\sharp_{-1,1}) = 2 + (-2) + 2(3) - 2 = 4,$$

which is a contradiction. ☐

Proof of Lemma 4.3.6. This is a combination of Lemmas 4.6.1 and 4.6.4. ☐

5. STABILIZATION

The goal of this section is to prove Theorem 1.0.2. Let $N = N_{(S,h)}$ be the suspension of (S,h), where S is a bordered surface of genus g with connected boundary and $h : (S,\omega) \xrightarrow{\sim} (S,\omega)$ is a symplectomorphism which has zero flux and restricts to the identity on ∂S.

The strategy of the proof is to apply two positive stabilizations to (S,h) — corresponding to the connected sum with a trefoil knot — to obtain (S',h'), where S' has connected boundary and genus $g+1$. We then compare $\Phi_{(S,h)}$ and $\Phi_{(S',h')}$, which both induce isomorphisms on the level of homology. Here $\Phi_{(S,h)}$ is the Φ map for (S,h).

5.1. **The setup.** Let T be a genus one surface with connected boundary and let η_0, η_1 be two essential simple closed curves on T which intersect transversely in one point. A positive Dehn twist along a closed curve η will be denoted by τ_η. Let $h_T : T \overset{\sim}{\to} T$ be the first return map of a Reeb vector field on the suspension

$$N_T = N_{(T,h_T)} = (T\times[0,1])/((x,1)\sim(h_T(x),0))$$

so that the following hold:

1. h_T is isotopic to $\tau_{\eta_0}\circ\tau_{\eta_1}$ relative to the boundary;
2. all the Reeb orbits in the interior of N_T which intersect $T\times\{0\}$ at most $2g+2$ times are nondegenerate;
3. $h_T|_{\partial T} = id$ and ∂N_T is foliated by a negative Morse-Bott family of slope ∞.

In view of the discussion in Section I.3, h_T will be viewed interchangeably as (i) the first return map of a Reeb vector field or (ii) the time-1 map of a stable Hamiltonian vector field with zero flux.

Now let S' be the boundary connected sum of S and T. More precisely, if P is a pair-of-pants with boundary $\partial P = \partial_1 P \sqcup \partial_2 P \sqcup \partial_3 P$, then S' is obtained from $S\sqcup T\sqcup P$ by identifying $\partial S \simeq -\partial_1 P$ and $\partial T \simeq -\partial_2 P$. Observe that $\partial S' = \partial_3 P$ is connected and $g(S') = g+1$. See Figure 19.

We now define a symplectomorphism $h' : (S',\omega') \overset{\sim}{\to} (S',\omega')$ with zero flux (with respect to some ω') as follows: First set $h'|_S = h$ and $h'|_T = h_T$. Then define $h'|_P$ as the first return map of a Reeb flow R_α on

$$N_P = N_{(P,id)} = (P\times[0,1])/((x,1)\sim(x,0)).$$

The contact form α is given by $f_0 dt + \beta$, where f_0 is a function and β is a 1-form on P, and both f_0 and β do not depend on t. We choose a Morse-Bott function $f_0 : P \to \mathbb{R}$ which satisfies the following:

1. f_0 is C^k-close to 1 for $k \gg 0$;
2. f_0 attains its minimum along the Morse-Bott family ∂P;
3. the critical points of f_0 in $int(P)$ are isolated and consist of the maximum e_P and two saddles h_{1P}, h_{2P}.

The Reeb orbits corresponding to e_P, h_{1P}, h_{2P} will also be denoted by e_P, h_{1P}, h_{2P}.

Let $\mathcal{N}_i$, $i = 1,2,3$, be the negative Morse-Bott family of Reeb orbits corresponding to $\partial_i P$. If f_0 is C^k-close to 1 for $k \gg 0$, then the only orbits that intersect $P\times\{0\}$ at most $2g+2$ times are: e_P, h_{iP} and the orbits of $\mathcal{N}_i$. We pick two orbits in each $\mathcal{N}_i$ and label them e_i, h_i; they will become elliptic and hyperbolic when the Morse-Bott function f_0 is perturbed into a Morse function, which we call f. For convenience we write N_S for the suspension of (S,h), $N_{S'}$ for the suspension of (S',h'), etc.



FIGURE 19. The page S'. The gradient trajectories of the Morse function f are given.

Since (S',h') is obtained from (S,h) by applying two positive stabilizations, the corresponding contact structures $\xi_{(S,h)}$ and $\xi_{(S',h')}$ are isomorphic. As a special case, observe that if $(S,h)=(D^2,id)$, then $(S',h')=(T,h_T)$ and (T,h_T) is an open book decomposition for the standard tight contact structure on S^3.

5.2. **Morse-Bott theory.** Let J be an adapted almost complex structure on $\mathbb{R}\times N_{S'}$ and let $\pi:\mathbb{R}\times N_{S'}\to N_{S'}$ be the projection onto the second factor. Also let u be a finite energy Morse-Bott building in $\mathbb{R}\times N_{S'}$ from γ to γ', where both orbit sets intersect $S'\times\{0\}$ in $2g+2$ points.

The following lemma partitions the irreducible components of u into three regions. When u is a Morse-Bott building, then we say that u is "irreducible" if the holomorphic curve, obtained by perturbing the Morse-Bott contact form to a nondegenerate one and correspondingly gluing up the levels of the Morse-Bott building, is irreducible.

Lemma 5.2.1. *Every irreducible component of the Morse-Bott building u in $\mathbb{R}\times N_{S'}$ has image in one of $\mathbb{R}\times N_S$, $\mathbb{R}\times N_T$, or $\mathbb{R}\times N_P$.*

Proof. The lemma is an application of the winding number wind_π of [HWZ1] and the positivity of intersections. (For example, see [CGH1, Lemma 5.5.1].)

Suppose without loss of generality that $u:\dot{F}\to\mathbb{R}\times N_{S'}$ is a single-level Morse-Bott building and that $u(\dot{F})$ nontrivially intersects $\mathbb{R}\times N_P$; the case of a multiple-level Morse-Bott building only differs in notation. If u corresponds to a gradient trajectory from h_i to e_i, then the lemma holds. Otherwise let $P_\varepsilon\subset P$ be a slight retraction of P so that $\partial P_\varepsilon=\partial_1^\varepsilon P\sqcup\partial_2^\varepsilon P\sqcup\partial_3 P$. Let $\Pi: N_P=P\times S^1\to P$ be the projection onto the first factor, let $T_i^\varepsilon=\Pi^{-1}(\partial_i^\varepsilon P)$, $i=1,2$, and let $N_{P_\varepsilon}=\Pi^{-1}(P_\varepsilon)$. We may assume that T_i^ε is foliated by (not necessarily closed) Reeb orbits. We consider the intersection $\delta_i = \pi(u(\dot{F})) \cap T_i^\varepsilon$, where δ_i is given the boundary orientation of $\pi(u(\dot{F})) \cap N_{P_\varepsilon}$. The curve δ_i is transverse to the Reeb vector field away from a finite number of points by [CGH1, Lemma 5.5.1]. If δ_i is not homologous to a multiple of $\{pt\} \times S^1$ in $H_1(T_i^\varepsilon)$, then $\Pi(\delta_i)$ is in the homology class $k[\partial_i P_\varepsilon] \in H_1(P_\varepsilon)$, $k > 0$, by the positivity of intersections in dimension four. This implies that $\Pi(\pi(u(\dot{F})) \cap \partial N_{S'})$ is in the class $k[\partial_3 P] \in H_1(P)$ for $k > 0$. Since this is a contradiction, we must have $[\delta_i] = l[\{pt\} \times S^1] \in H_1(T_i^\varepsilon)$, for some $l \geq 0$. This implies that u has negative ends along $\mathcal{N}_i$, $i = 1, 2$. Hence u has image in $\mathbb{R} \times N_P$. □

We now use Morse-Bott theory (cf. Bourgeois [Bo1, Bo2]) to analyze holomorphic curves on $\mathbb{R} \times N_P$. In particular, we consider a perturbation of the Morse-Bott family of orbits on N_P, perturbed by the Morse function $f : P \to \mathbb{R}$ as described above.

Lemma 5.2.2. *Let $f : P \to \mathbb{R}$ be C^k-close to 1 for $k \gg 0$. Then there is an adapted almost complex structure J on $\mathbb{R} \times N_P$ such that every $I_{ECH} = 1$ finite energy J-holomorphic curve u whose image is in $\mathbb{R} \times N_P$, is a simply-covered cylinder which corresponds to a gradient flow line between critical points of f of adjacent index. The complete list is as follows:*

1. *one cylinder from e_P to h_i, $i = 1, 2, 3$, and two cylinders from e_P to h_{iP}, $i = 1, 2$;*
2. *two cylinders from h_i to e_i, $i = 1, 2, 3$;*
3. *one cylinder from h_{1P} to e_1 and one cylinder from h_{1P} to e_3;*
4. *one cylinder from h_{2P} to e_2 and one cylinder from h_{2P} to e_3.*

Moreover, all the $I_{ECH} = 1$ curves above are regular.

See Figure 19 for the gradient trajectories of f which correspond to the above holomorphic cylinders.

Proof. The proof is very similar to that of [CGH1, Lemma 9.2.2]. We use the following fact, which can be proved using Morse-Bott theory or by a direct computation:

Fact. There is an adapted almost complex structure J on $\mathbb{R} \times N_P$ such that there is a one-to-one correspondence between (parametrized) gradient trajectories $\delta : \mathbb{R} \to P$ of f and finite energy J-holomorphic cylinders Z_δ in $\mathbb{R} \times N_P$ which intersect each $\{(s,t)\} \times P$ exactly once and project to $\mathrm{Im}(\delta)$ under the projection $\Pi \circ \pi : \mathbb{R} \times N_P \to P$, $(s, x, t) \mapsto x$. Moreover, the cylinders Z_δ, together with the trivial cylinders over the orbits corresponding to the Morse critical points, give a finite energy foliation of $\mathbb{R} \times N_P$.

Fix f and J as above. We will use the notation $s\delta : \mathbb{R} \to P$ for the translation $(s\delta)(\tau) = \delta(\tau + s)$.

Let $u : \dot{F} \to \mathbb{R} \times N_P$ be a finite energy J-holomorphic curve. Let $D_\varepsilon \subset P$ be an arbitrarily small disk centered at the point e_P and let $N(e_P) = \Pi^{-1}(D_\varepsilon)$ be a solid torus neighborhood of the orbit e_P. We assume that $\partial N(e_P)$ is foliated by (not necessarily closed) orbits of the Reeb vector field. We identify $\partial N(e_P) \simeq \mathbb{R}^2/\mathbb{Z}^2$ so that the meridian has slope zero and a fiber $\{pt\} \times S^1$ has slope ∞. Consider $\eta = \pi(u(\dot{F})) \cap \partial N(e_P)$, where η is given the boundary orientation of $\pi(u(\dot{F})) \cap \Pi^{-1}(P - int(D_\varepsilon))$. If the projection $[\Pi(\eta)] \in H_1(P - int(D_\varepsilon))$ is nonzero, then $[\Pi(\eta)] = -k[\partial D_\varepsilon]$, $k > 0$, by the positivity of intersections. This is a contradiction as in the proof of Lemma 5.2.1. Hence $[\eta] = l[\{pt\} \times S^1] \in H_1(\partial N(e_P))$ for some $l \geq 0$, i.e., has slope ∞. In other words, u cannot intersect $\mathbb{R} \times e_P$ and can only have e_P at the positive end. Similar considerations hold for $N(e_i)$, $i = 1, 2, 3$, where $D_{i,\varepsilon} \subset P$ is a half-disk centered at e_i and $N(e_i) = \Pi^{-1}(D_{i,\varepsilon})$.

We now claim that u is some multiple cover of some Z_δ with multiplicity ≥ 1. Arguing by contradiction, suppose u does not multiply cover any Z_δ. Let us first consider the case where $\Pi \circ \pi(u(\dot{F}))$ does not equal $\mathrm{Im}(\delta)$ for any δ. Then there is some Z_δ from e_P to some e_i such that the intersection $u(\dot{F}) \cap Z_\delta$ is nonempty; moreover, in view of the asymptotics on $N(e_P)$ and $N(e_i)$, we may assume that $K = \pi(u(\dot{F}) \cap Z_\delta)$ is compact. This implies that $u(\dot{F})$ and $Z_{s\delta}$ do not intersect for sufficiently large s. On the other hand, since the intersection pairing $\langle u(\dot{F}), Z_{s\delta}\rangle$ is a homological quantity and does not depend on s due to the asymptotics, it follows that $K = \varnothing$, which is a contradiction. This implies that $\Pi \circ \pi(u(\dot{F})) = \mathrm{Im}(\delta)$ for some δ. Now $\mathbb{R} \times \Pi^{-1}(\delta)$ is a 3-manifold which is foliated by $Z_{s\delta}$, $s \in \mathbb{R}$, and if u does not multiply cover any $Z_{s\delta}$, then u intersects some $Z_{s\delta}$ along a 1-manifold, a contradiction. We conclude that u is a multiple cover of some Z_δ.

Now Fredholm index one cylinders Z_δ — i.e., those that correspond to δ connecting two Morse critical points of adjacent index — are regular by the automatic transversality results of Wendl [We2, We3]. Moreover, u cannot multiply cover Z_δ with multiplicity > 1 by [HT1, Proposition 7.15], since otherwise $I(u) > 1$. This implies that u is equal to some Z_δ, thereby completing the proof of the lemma. □

5.3. **A commutative diagram.** We assume that h has no elliptic periodic points of period $\leq 2g + 2$ in $int(S)$.

Consider the following (not commutative) diagram of chain complexes:

(5.3.1)
$$\begin{tikzcd}
\widehat{CF}(S, \mathbf{a}, h(\mathbf{a})) \arrow[r, "\Theta"] \arrow[d, "\Phi_{(S,h)}"'] & \widehat{CF}(S', \mathbf{a}', h'(\mathbf{a}')) \arrow[d, "\Phi_{(S',h')}"] \\
PFC_{2g}(S) \arrow[r, "i \circ j"] & PFC_{2g+2}(S')
\end{tikzcd}$$

Here we are writing $PFC_j(S)$ for $PFC_j(N_S)$ and $PFC_j(S')$ for $PFC_j(N_{S'})$. The chain maps Θ, j, and i which appear Diagram (5.3.1) are defined as follows:

The map Θ. The chain map

$$\Theta : \widehat{CF}(S, \mathbf{a}, h(\mathbf{a})) \to \widehat{CF}(S', \mathbf{a}', h'(\mathbf{a}'))$$

is defined as follows: Let $\mathbf{a} = \{a_1, \dots, a_{2g}\}$ be a basis of S. We extend $\mathbf{a}$ to $\overline{\mathbf{a}} = \{\overline{a}_1, \dots, \overline{a}_{2g}\}$ so that $\overline{a}_i$ is a properly embedded arc with boundary on $\partial S'$ and then complete $\overline{\mathbf{a}}$ to a basis $\mathbf{a}'$ of S' by adding the extensions $\overline{a}_{2g+1}$ and $\overline{a}_{2g+2}$ of the basis arcs a_{2g+1} and a_{2g+2} of T, subject to the following conditions:

- $\overline{a}_i$ and $h'(\overline{a}_i)$ have two extra pairs of canceling intersections in $S'-(S\cup T)$ for $i=1,\dots,2g+2$;
- $\overline{a}_i - a_i$ and $h'(\overline{a}_j - a_j)$ are disjoint for $1\leq i\neq j\leq 2g+2$.

Let $x_i\sim x_i'$, $i=2g+1,2g+2$, be the intersections of $h'(\overline{a}_i)$ and $\overline{a}_i$ on $\partial S'=\partial_3 P$, which comprise the contact class. We then define

$$\Theta(\mathbf{y})=\mathbf{y}\cup\{x_{2g+1},x_{2g+2}\}.$$

It is easy to see that Θ is a chain map. (Compare with the gluing map from [HKM2].)

The map j. Next define the map

$$j: PFC_{2g}(S)\to PFC_{2g+2}(S),$$

$$j(\gamma)=e_1^2\gamma,$$

where we are using multiplicative notation for orbit sets. The map j is the composition $\mathfrak{I}_{2g+1}\circ\mathfrak{I}_{2g}$, where $\mathfrak{I}_i$ is the map

$$\mathfrak{I}_i: PFC_i(S)\to PFC_{i+1}(S),$$

$$\gamma\mapsto e_1\gamma.$$

The elliptic orbit e_1 can only appear at the negative asymptotic end of a holomorphic curve, provided the curve is not a trivial cylinder. Hence j is a chain map in view of Lemma 5.2.1.

The map i. The map i is the inclusion map on the chain level:

$$i: PFC_{2g+2}(S)\to PFC_{2g+2}(S'),$$

$$\gamma\mapsto\gamma.$$

The map i is chain map by Lemma 5.2.1. The analogous map for the inclusion $T\subset S'$ is also a chain map.

Although Diagram (5.3.1) is not quite commutative on the chain level, we have the following:

Lemma 5.3.1. *Diagram (5.3.1) induces the following commutative diagram on the level of homology:*

(5.3.2)
$$\begin{tikzcd}
\widehat{HF}(S,\mathbf{a},h(\mathbf{a})) \arrow[r, "\Theta_*"] \arrow[d, "(\Phi_{(S,h)})_*"'] & \widehat{HF}(S',\mathbf{a}',h'(\mathbf{a}')) \arrow[d, "(\Phi_{(S',h')})_*"] \\
PFH_{2g}(S) \arrow[r, "(i\circ j)_*"] & PFH_{2g+2}(S')
\end{tikzcd}$$

Proof. We consider the W_+-curves for (S', h') that limit to the tuple $\Theta(\mathbf{y}) = \mathbf{y} \cup \{x_{2g+1}, x_{2g+2}\}$ at the positive end. Since x_{2g+1} and x_{2g+2} are components of the contact class, there are no holomorphic curves, besides restrictions of trivial cylinders, that limit to x_{2g+1} or x_{2g+2} at the positive end. The argument is the same as that of Lemma I.6.2.3. The trivial cylinder over x_i, $i = 2g+1, 2g+2$, "maps" the component x_i to a generic point p_i of the Morse-Bott family $\mathcal{N}_3$ and is concatenated with a cylinder corresponding to a gradient trajectory from p_i to e_3 to give a Morse-Bott building. Moreover, by automatic transversality [We2, Theorem 4.5.36] and the discussion from Lemma I.5.8.9, the above Morse-Bott building is Morse-Bott regular. Hence the pair $\{x_{2g+1}, x_{2g+2}\}$ is "mapped" to e_3^2. By Lemma 5.2.1 we have:

$$\Phi_{(S',h')}(\mathbf{y} \cup \{x_{2g+1}, x_{2g+2}\}) = e_3^2 \cdot \Phi_{(S,h)}(\mathbf{y}).$$

On the other hand, $e_3^2\gamma$ is homologous to $e_1^2\gamma$ in $PFH_{2g+2}(S')$, since there is one cylinder each from h_{1P} to e_1 and e_3. This proves the commutativity of the diagram. □

Corollary 5.3.2. *The composition $i_* \circ j_*$ is an isomorphism.*

Proof. This follows from the commutativity of Diagram (5.3.2). Since (S, h) and (S', h') are both open book decompositions for M and we may assume that h and h' both have no periodic points of period $\leq 2g+2$ in $int(S)$ or $int(S')$, the cobordism maps $(\Phi_{(S,h)})_*$ and $(\Phi_{(S',h')})_*$ are both isomorphisms. The map Θ_* is also an isomorphism since $\widehat{HF}(S', \mathbf{a}', h'(\mathbf{a}'))$ can be computed as the tensor product of the S and T sides and the T side gives $\widehat{HF}(S^3)$, which is generated by $\{x_{2g+1}, x_{2g+2}\}$. Since three sides of the diagram are isomorphisms, it follows that $i_* \circ j_*$ is an isomorphism. □

5.4. **An isomorphism.**

To simplify notation let us write $V_i = PFH_i(S)$ and $W_j = PFH_j(T)$. The goal of this subsection is to prove the following key proposition.

Proposition 5.4.1. *The induced map*

$$j_* = (\mathfrak{I}_{2g+1})_* \circ (\mathfrak{I}_{2g})_* : V_{2g} \to V_{2g+2}$$

is an isomorphism.

5.4.1. *Description of the differential.* Given two orbit sets $\gamma' = \prod \gamma_i^{m_i'}$ and $\gamma = \prod \gamma_i^{m_i}$, we set $\gamma/\gamma' = \prod \gamma_i^{m_i - m_i'}$ if $m_i' \leq m_i$ for all i; otherwise we set $\gamma/\gamma' = 0$.

The chain group $PFC_{2g+2}(S')$ can be written as:

$$PFC_{2g+2}(S') = \bigoplus_{m+i+j=2g+2} \left(\mathbb{F}[h_{1P}, h_{2P}, h_3, e_P, e_3]_m \otimes PFC_i(S) \otimes PFC_j(T)\right).$$

$\mathbb{F}[h_{1P}, h_{2P}, h_3, e_P, e_3]$ is a polynomial ring where h_{1P}, h_{2P}, h_3 (resp. e_P, e_3) are considered as Grassmann variables of odd degree (resp. even degree) and the subscript m indicates the subspace spanned by monomials with total exponent m.

Let us write a generator of $PFC_{2g+2}(S')$ as $\gamma\otimes\Gamma_1\otimes\Gamma_2$, where $\Gamma_1\in PFC_i(S)$, $\Gamma_2\in PFC_j(T)$, and γ is constructed from orbits passing through $S'-S-T$. Using Lemma 5.2.1 and the description of ECH index one curves in $\mathbb{R}\times N_P$ from Lemma 5.2.2, we write the differential ∂ of $PFC_{2g+2}(S')$ as follows:

$$\begin{aligned}\partial(\gamma\otimes\Gamma_1\otimes\Gamma_2) =&\gamma\otimes(\partial_S\Gamma_1)\otimes\Gamma_2+\gamma\otimes\Gamma_1\otimes(\partial_T\Gamma_2)\\ &+(\gamma/e_P)(h_3\otimes\Gamma_1\otimes\Gamma_2+1\otimes h_1\Gamma_1\otimes\Gamma_2+1\otimes\Gamma_1\otimes h_2\Gamma_2)\\ &+(\gamma/h_{1P})(e_3\otimes\Gamma_1\otimes\Gamma_2+1\otimes e_1\Gamma_1\otimes\Gamma_2)\\ &+(\gamma/h_{2P})(e_3\otimes\Gamma_1\otimes\Gamma_2+1\otimes\Gamma_1\otimes e_2\Gamma_2).\end{aligned}$$

Here ∂_S and ∂_T are the differentials on $PFC(S)$ and $PFC(T)$.

5.4.2. *Spectral sequence calculation.* In this subsection we use spectral sequences to interpret $PFH_{2g+2}(S')$ in terms of V_i and W_j. More precisely, we prove the following:

Lemma 5.4.2. *There is a natural inclusion of* $\left(\bigoplus_{i+j=2g+2}V_i\otimes W_j\right)/\sim$ *into* $PFH_{2g+2}(S')$*, where the equivalence relation* $\sim$ *is given by* $e_1\Gamma_1\otimes\Gamma_2\sim\Gamma_1\otimes e_2\Gamma_2$*. Here* $\Gamma_1\in V_i$ *and* $\Gamma_2\in W_j$ *and the equivalence* $\sim$ *is generated by terms in adjacent* $V_{i+1}\otimes W_j$ *and* $V_i\otimes W_{j+1}$*.*

Proof. Let $\mathcal{F}$ be a filtration on $(PFC_{2g+2}(S'),\partial)$ which, on the generators, counts the multiplicity of h_{1P}. This means that $\mathcal{F}$ takes values in $\{0,1\}$. We write $(E^r(\mathcal{F}),\partial_r)$ for the E^r-term and the E^r-differential of the spectral sequence for $\mathcal{F}$. Each page $E^r(\mathcal{F})$ has a grading coming from $\mathcal{F}$ and $E^r_k(\mathcal{F})$ is the degree k component of $E^r(\mathcal{F})$ with respect to this grading. We remark that the spectral sequence associated to $\mathcal{F}$ is nothing but a long exact sequence in homology, and its use is motivated by our wish to give a parallel treatment of the cases where we filter by the multiplicity of a hyperbolic orbit or by the multiplicity of an elliptic orbit.

Next let $\mathcal{G}$ be a filtration on $(E^0(\mathcal{F}),\partial_0)$ which counts the multiplicity of h_{2P}. Again, $\mathcal{G}$ takes values in $\{0,1\}$. We write $(E^r(\mathcal{G}),\partial_{0r})$ for the E^r-term and the E^r-differential of the spectral sequence for $\mathcal{G}$. Finally, let $\mathcal{H}$ be the filtration on $(E^0(\mathcal{G}),\partial_{00})$ which counts the multiplicity of e_P, and let $(E^r(\mathcal{H}),\partial_{00r})$ be the E^r-term and the E^r-differential of the spectral sequence for $\mathcal{H}$.

We first consider $(E^0(\mathcal{H}),\partial_{000})$, where:

$$\partial_{000}(\gamma\otimes\Gamma_1\otimes\Gamma_2)=\gamma\otimes(\partial_S\Gamma_1)\otimes\Gamma_2+\gamma\otimes\Gamma_1\otimes(\partial_T\Gamma_2).$$

By the Künneth formula, we have:

$$E^1(\mathcal{H})=\bigoplus_{m+i+j=2g+2}\left(\mathbb{F}[h_{1P},h_{2P},h_3,e_P,e_3]_m\otimes V_i\otimes W_j\right).$$

Next consider $(E^1(\mathcal{H}),\partial_{001})$, where:

$$\begin{aligned}\partial_{001}(e_P^n\gamma\otimes\Gamma_1\otimes\Gamma_2)=&e_P^{n-1}h_3\gamma\otimes\Gamma_1\otimes\Gamma_2+e_P^{n-1}\gamma\otimes(h_1\Gamma_1)\otimes\Gamma_2\\ &+e_P^{n-1}\gamma\otimes\Gamma_1\otimes(h_2\Gamma_2),\end{aligned}$$

Here $\Gamma_1 \in V_i$, $\Gamma_2 \in W_j$, and γ has no e_P term. Note that any $e_P^{n-1}h_3\gamma \otimes \Gamma_1 \otimes \Gamma_2$ is homologous (with respect to the differential ∂_{001}) to a linear combination of $e_P^{n-1}\gamma' \otimes \Gamma_1' \otimes \Gamma_2'$, where γ' does not have any e_P and h_3 terms, Γ_1' is in some V_i, and Γ_2' is in some W_j. Hence every element of $E^1(\mathcal{H})/\operatorname{Im}(\partial_{001})$ can be represented by $w = \sum_{i\geq 0} e_P^i w_i$, where w_i has no e_P and h_3 terms, and we can write

$$\partial_{001}(w) = \sum_{i>0} e_P^{i-1} h_3 w_i + w',$$

where w' has no terms which contain h_3. If $\partial_{001}(w) = 0$, then all the w_i, $i > 0$, must be zero. We therefore obtain:

$$E^2(\mathcal{H}) = \bigoplus_{m+i+j=2g+2} \left(\mathbb{F}[h_{1P}, h_{2P}, e_3]_m \otimes V_i \otimes W_j\right).$$

Since $E^2(\mathcal{H})$ is supported in degree 0, the spectral sequence for $\mathcal{H}$ converges at the E^2-term and we have $E^1(\mathcal{G}) \simeq E^2(\mathcal{H})$. Moreover, $E^2(\mathcal{H}) = E_0^2(\mathcal{H})$ is naturally isomorphic to $E^1(\mathcal{G})$ since 0 is the lowest filtration level.

Now consider $(E^1(\mathcal{G}), \partial_{01})$, where

$$\partial_{01}(\gamma \otimes \Gamma_1 \otimes \Gamma_2) = e_3(\gamma/h_{2P}) \otimes \Gamma_1 \otimes \Gamma_2 + (\gamma/h_{2P}) \otimes \Gamma_1 \otimes e_2\Gamma_2.$$

The calculation of $E^2(\mathcal{G})$ is similar to the calculation of $E^2(\mathcal{H})$ in the previous paragraph. Any $e_3^{n+1}\gamma \otimes \Gamma_1 \otimes \Gamma_2$ is homologous to a linear combination of $e_3^n\gamma' \otimes \Gamma_1' \otimes \Gamma_2'$, where γ, γ' do not have any h_{2P} and e_3 terms. Hence every element of $E^1(\mathcal{G})/\operatorname{Im}(\partial_{01})$ can be represented by $w = w' + \sum_{i=0}^k h_{2P} e_3^i w_i$, where w' and w_i have no h_{2P} or e_3 terms, and we can write

$$\partial_{01} w = \sum_{i=0}^k e_3^{i+1} w_i + \sum_{i=1}^k e_3^i e_2 w_i.$$

If $\partial_{01} w = 0$, then all the w_i must be zero. Only the w' term remains, and we have:

$$E^2(\mathcal{G}) = \bigoplus_{m+i+j=2g+2} \left(\mathbb{F}[h_{1P}]_m \otimes V_i \otimes W_j\right),$$

which we can write as a direct sum $\mathfrak{L}_0 \oplus \mathfrak{L}_1$, where:

$$\mathfrak{L}_0 = \bigoplus_{i+j=2g+2} V_i \otimes W_j,$$

$$\mathfrak{L}_1 = \bigoplus_{i+j=2g+1} \mathbb{F}\{h_{1P}\} \otimes V_i \otimes W_j.$$

Since $E^2(\mathcal{G})$ is supported in the lowest degree 0, the spectral sequence for $\mathcal{G}$ converges at the E^2-term and $E^2(\mathcal{G}) = E_0^2(\mathcal{G})$ is naturally isomorphic to $E^1(\mathcal{F})$.

Finally, $E^1(\mathcal{F}) \simeq \mathfrak{L}_0 \oplus \mathfrak{L}_1$ has differential ∂_1 given by:

$$\partial_1(\Gamma_1 \otimes \Gamma_2) = 0, \tag{5.4.1}$$

$$\partial_1(h_{1P} \otimes \Gamma_1 \otimes \Gamma_2) = e_1\Gamma_1 \otimes \Gamma_2 + \Gamma_1 \otimes e_2\Gamma_2, \tag{5.4.2}$$

since $\partial(h_{1P} \otimes \Gamma_1 \otimes \Gamma_2) = e_1\Gamma_1 \otimes \Gamma_2 + e_3 \otimes \Gamma_1 \otimes \Gamma_2$ and $\partial(h_{2P} \otimes \Gamma_1 \otimes \Gamma_2) = \partial_0(h_{2P} \otimes \Gamma_1 \otimes \Gamma_2) = e_3 \otimes \Gamma_1 \otimes \Gamma_2 + \Gamma_1 \otimes e_2\Gamma_2$. Viewing the spectral sequence as a long exact sequence with connecting homomorphism $\mathfrak{L}_1 \xrightarrow{\delta} \mathfrak{L}_0$, where δ is given by Equation (5.4.2), we see that $E^2(\mathcal{F})$ is isomorphic to $\mathfrak{L}_0/\operatorname{Im}(\delta) = \mathfrak{L}_0/\sim$, where the equivalence relation is given by $e_1\Gamma_1 \otimes \Gamma_2 \sim \Gamma_1 \otimes e_2\Gamma_2$. As argued previously, the spectral sequence converges at the E^2-term and $E^2(\mathcal{F})$ is naturally isomorphic to $PFH_{2g+2}(S')$. □

5.4.3. *Isomorphism of* j_*. We now complete the proof of Proposition 5.4.1.

Proof of Proposition 5.4.1. Since $i_* \circ j_* : V_{2g} \to PFH_{2g+2}(S')$ is an isomorphism by Corollary 5.3.2, $j_* : V_{2g} \to V_{2g+2}$ is injective. It remains to show that j_* is surjective.

First note that $W_0 = \mathbb{F}\{1\}$. We also have $W_2 = \mathbb{F}\{e_2^2\}$, since $\widehat{HF}(S^3) \simeq \mathbb{F}$, generated by the contact class $c(\xi_{std})$ of the standard tight contact structure ξ_{std} on S^3, and e_2^2 is the image of $c(\xi_{std})$ under the isomorphism

$$(\Phi_{(T,h_T)})_* : \widehat{HF}(S^3) = \widehat{HF}(T, h_T) \xrightarrow{\sim} PFH_2(T).$$

Now, since the isomorphism $W_0 \xrightarrow{\sim} W_2$, $1 \mapsto e_2^2$, factors as $W_0 \to W_1 \to W_2$, $1 \mapsto e_2 \mapsto e_2^2$, it follows that $W_1 = \mathbb{F}\{e_2\} \oplus W_1'$ for some $\mathbb{F}$-vector space W_1'.

By Lemma 5.4.2,

$$PFH_{2g+2}(S') \simeq \left(\bigoplus_{i+j=2g+2} V_i \otimes W_j \right) / \sim .$$

Moreover, $V_{2g} \otimes W_2 = V_{2g} \otimes \mathbb{F}\{e_2^2\}$ is the image of V_{2g} under the map $i_* \circ j_*$. Since $i_* \circ j_*$ is an isomorphism, every element of $V_{2g+2} \otimes W_0 = V_{2g+2} \otimes \mathbb{F}\{1\}$ is equivalent to some element of $V_{2g} \otimes W_2$, i.e., if $v_{2g+2} \in V_{2g+2}$, then $v_{2g+2} \otimes 1 \sim v_{2g} \otimes e_2^2$ for some $v_{2g} \in V_{2g}$. More explicitly,

$$\begin{aligned} v_{2g+2} \otimes 1 + v_{2g} \otimes e_2^2 = &(e_1 v''_{2g+1} \otimes 1 + v''_{2g+1} \otimes e_2) + (e_1 v''_{2g} \otimes e_2 + v''_{2g} \otimes e_2^2) \\ &+ \sum_i (e_1 v''_{2g,i} \otimes w'_{1,i} + v''_{2g,i} \otimes e_2 w'_{1,i}) + \dots , \end{aligned}$$

where $v''_{2g+1} \in V_{2g+1}$; $v''_{2g}, v''_{2g,i} \in V_{2g}$; $\{w'_{1,i}\}$ is a basis for W_1'; and $\dots$ is a linear combination of terms which are not in $V_{2g+2} \otimes W_0$ and $V_{2g+1} \otimes W_1$. A term-by-term comparison gives $v_{2g+2} = e_1 v''_{2g+1}$ and $v''_{2g+1} = e_1 v''_{2g}$. Hence $v_{2g+2} = e_1^2 v''_{2g}$ and $v_{2g+2} \in \operatorname{Im}(j_*)$. This completes the proof of Proposition 5.4.1. □

5.5. **Proof of Theorem 1.0.2.** We use the notation from I.2.5.3.

We first consider the ECH cobordism map

$$\mathfrak{I}''_j : ECC_j(N, f_j\alpha) \to ECC_j(N, f_{j+2}\alpha),$$

defined via Seiberg-Witten Floer homology as in [HT3, Theorem 2.4].

More precisely, let $\delta_k > 0$ be constants satisfying $\lim_{k\to\infty} \delta_k = 0$. Let $V_{j+2}^{\delta_k}$ be a δ_k-tubular neighborhood of the elliptic orbits of $R_{f_j\alpha}$ with $\mathcal{F} = j+1$ or $j+2$, where δ_k is measured with respect to a fixed Riemannian metric on N, and let $f_{j+2,k} : N \to (0, +\infty)$ be a function such that the following hold:

(1) $f_{j+2,k} = f_j$ on $N - V^{\delta_k}_{j+2}$;
(2) $R_{f_{j+2,k}\alpha}$ has no elliptic orbits with $\mathcal{F} \leq j+2$; and
(3) $\lim\limits_{k\to\infty} f_{j+2,k} = f_j$ in C^1.

Let J_j be an $f_j\alpha$-adapted almost complex structure on $\mathbb{R}\times N$ and let $J_{j+2,k}$ be $f_{j+2,k}\alpha$-adapted almost complex structures on $\mathbb{R}\times N$ which agree with J_j on $\mathbb{R}\times(N - V^{\delta_k}_{j+2})$ and satisfy $J_{j+2,k}\to J_j$ in C^0 as $k\to\infty$. Also let $\widetilde{J}_{j,k}$ be an almost complex structure on $\mathbb{R}\times N$ which adapted to an exact symplectic cobordism $\omega_{j,k}$ from $f_j\alpha$ to $f_{j+2,k}\alpha$, agrees with J_j on $\mathbb{R}\times(N - V^{\delta_k}_{j+2})$, and satisfies $\widetilde{J}_{j,k}\to J_j$ as $k\to\infty$.

Observe that the generators of $ECC_j(N, f_j\alpha)$ and $ECC_j(N, f_{j+2,k}\alpha)$ agree for $k \gg 0$. Let γ and γ' denote orbit sets for either $f_j\alpha$ or $f_{j+2,k}\alpha$.

Lemma 5.5.1. *Let u_k either be a sequence of $J_{j+2,k}$-holomorphic curves from γ to γ' or a sequence of $\widetilde{J}_{j,k}$-holomorphic curves from γ to γ'. Then, after passing to a subsequence, u_k converges to a J_j-holomorphic building u from γ to γ'.*

Sketch of proof. This follows from Gromov compactness for C^0-convergence of almost complex structures, due to Ivashkovich-Shevchishin [IS].

Alternatively, we can argue as follows, using the usual Gromov compactness for C^∞-convergence of almost complex structures: We treat the case where u_k is a $J_{j+2,k}$-holomorphic curve. By the compactness argument from I.3.4, we may assume that $[u_k]\in H_2(N,\gamma,\gamma')$ and the topological type of the domain F_k of u_k are fixed. We then restrict u_k to the preimage G_k of $\mathbb{R}\times(N - V^{\delta_k}_{j+2})$. We may assume that G_k is obtained from a compact Riemann surface with boundary by removing interior punctures. (This is possible by taking δ_k to be generic.)

We claim that $|\chi(G_k)|$ is bounded above. This is equivalent to an upper bound on the number of disk components of $F_k - G_k$. Let γ'' be the union of core orbits of $V^{\delta_k}_{j+2}$, which is independent of k. The number of disk components is bounded above by the intersection number with $\mathbb{R}\times\gamma''$, which in turn is controlled by the homology class $[u_k]\in H_2(N,\gamma,\gamma')$.

Since $J_{j+2,k} = J_j$ on $\mathbb{R}\times(N - V^{\delta_k}_{j+2})$, we can take the limit of $u_k|_{G_k}$ to obtain $u|_G$. For simplicity assume that u has only one level. Then ∂G is mapped to $\mathbb{R}\times\gamma''$, which contradicts the positivity of intersections unless $\partial G = \varnothing$. Hence $\partial G = \varnothing$ and all the interior punctures of G which do not to limit to γ or γ' are removable. $\square$

Lemma 5.5.2. *For $k\gg 0$, the ECH cobordism map*

$$\mathfrak{I}''_j : ECC_j(N, f_j\alpha) \to ECC_j(N, f_{j+2,k}\alpha)$$

satisfies $\mathfrak{I}''_j(\gamma) = \gamma$ and induces an isomorphism on the level of homology.

Proof. We use the Holomorphic Curves Property in [HT3, Theorem 2.4], which states that $\langle \mathfrak{I}''_j(\gamma), \gamma'\rangle$ is a sum of weights $w(v_i)$ over all $I = 0$ $\widetilde{J}_{j,k}$-holomorphic buildings $v_1,\dots,v_l$ from γ to γ'. By Lemma 5.5.1, for $k\gg 0$, the only $I=0$ $\widetilde{J}_{j,k}$-buildings are curves supported on trivial cylinders. Moreover, the Holomorphic

Curves Property also guarantees that $w(v)=1$ if v is supported on trivial cylinders. This implies that $\mathfrak{I}''_j(\gamma)=\gamma$.

Finally, since the chain map $\mathfrak{I}''_j$ is the identity, the differentials on $ECC_j(N, f_j\alpha)$ and $ECC_j(N, f_{j+2,k}\alpha)$ must agree. This implies the second assertion of the lemma. $\square$

We now let $f_{j+2}=f_{j+2,k}$ for $k\gg 0$. In view of Theorem I.2.5.6, we can write

$$\widehat{ECH}(M)\simeq \lim_{j\to\infty} ECH_{2j}(N, f_{2j}\alpha),$$

where the map $ECH_{2j}(N,f_{2j}\alpha)\to ECH_{2j+2}(N,f_{2j+2}\alpha)$ is given by the top row of:
(5.5.1)

$$\begin{tikzcd}
ECH_{2j}(N,f_{2j}\alpha) \arrow[r, "(\mathfrak{I}''_{2j})_*"] & ECH_{2j}(N,f_{2j+2}\alpha) \arrow[r, "j'_*"] \arrow[d, "\simeq"'] & ECH_{2j+2}(N,f_{2j+2}\alpha) \\
 & PFH_{2j}(N,f_{2j+2}\alpha) \arrow[r, "j_*"] & PFH_{2j+2}(N,f_{2j+2}\alpha) \arrow[u, "\simeq"']
\end{tikzcd}$$

Here j'_* and j_* are maps on homology induced by the inclusions $\gamma\mapsto e^2\gamma$, and the square commutes. Finally, Proposition 5.4.1 (and a similar argument) implies that j'_* is an isomorphism. This proves Theorem 1.0.2.

Acknowledgements. We are indebted to Michael Hutchings for many helpful conversations and for our previous collaboration which was a catalyst for the present work. We also thank Denis Auroux, Tobias Ekholm, Dusa McDuff, Ivan Smith and Jean-Yves Welschinger for illuminating exchanges. Part of this work was done while KH and PG visited MSRI during the academic year 2009–2010. We are extremely grateful to MSRI and the organizers of the "Symplectic and Contact Geometry and Topology" and the "Homology Theories of Knots and Links" programs for their hospitality; this work probably would never have seen the light of day without the large amount of free time which was made possible by the visit to MSRI. KH also thanks the Simons Center for Geometry and Physics for their hospitality during his visit in May 2011.

REFERENCES

[Ab] C. Abbas, *Finite energy surfaces and the chord problem*, Duke Math. J. **96** (1999), 241–316.

[Bo1] F. Bourgeois, *A Morse Bott approach to contact homology*, from: "Symplectic and contact topology: interactions and perspectives (Toronto, ON/Montréal, QC, 2001)", Fields Inst. Comm. **35**, Amer. Math. Soc., Providence, RI (2003) 55–77.

[Bo2] F. Bourgeois, *A Morse-Bott approach to contact homology*, Ph.D. thesis, 2002.

[BEE] F. Bourgeois, T. Ekholm and Y. Eliashberg, *Effect of Legendrian surgery*. With an appendix by Sheel Ganatra and Maksim Maydanskiy. Geom. Topol. **16** (2012) 301-389.

[BEHWZ] F. Bourgeois, Y. Eliashberg, H. Hofer, K. Wysocki and E. Zehnder, *Compactness results in symplectic field theory*, Geom. Topol. **7** (2003), 799–888 (electronic).

[CGH1] V. Colin, P. Ghiggini and K. Honda, *Embedded contact homology and open book decompositions*, preprint 2010. `ArXiv:1008.2734.`

[CGH2] V. Colin, P. Ghiggini and K. Honda, *The equivalence of Heegaard Floer homology and embedded contact homology I*, preprint 2011.

[CGH3] V. Colin, P. Ghiggini and K. Honda, *The equivalence of Heegaard Floer homology and embedded contact homology III: from hat to plus*, in preparation.

[CGHH] V. Colin, P. Ghiggini, K. Honda and M. Hutchings, *Sutures and contact homology I*, Geom. Topol. **15** (2011), 1749-1842.

[CHL] V. Colin, K. Honda and F. Laudenbach, *On the flux of pseudo-Anosov homeomorphisms*, Algebr. Geom. Topol. **8** (2008), 2147–2160.

[DS] S. Donaldson and I. Smith, *Lefschetz pencils and the canonical class for symplectic four-manifolds*, Topology **42** (2003), 743–785.

[Dr] D. Dragnev, *Fredholm theory and transversality for noncompact pseudoholomorphic maps in symplectizations*, Comm. Pure Appl. Math. **57** (2004), 726–763.

[EGH] Y. Eliashberg, A. Givental and H. Hofer, *Introduction to symplectic field theory*, GAFA 2000 (Tel Aviv, 1999), Geom. Funct. Anal. 2000, Special Volume, Part II, 560–673.

[Gi1] E. Giroux, *Convexité en topologie de contact*, Comment. Math. Helv. **66** (1991), 637–677.

[Gi2] E. Giroux, *Géométrie de contact: de la dimension trois vers les dimensions supérieures*, Proceedings of the International Congress of Mathematicians, Vol. II (Beijing, 2002), 405–414, Higher Ed. Press, Beijing, 2002.

[Gr] M. Gromov, *Pseudo holomorphic curves in symplectic manifolds*, Invent. Math. **82** (1985), 307–347.

[Ho1] H. Hofer, *Pseudoholomorphic curves in symplectizations with applications to the Weinstein conjecture in dimension three*, Invent. Math. **114** (1993), 515–563.

[Ho2] H. Hofer, *Holomorphic curves and dynamics in dimension three*, Symplectic geometry and topology (Park City, UT, 1997), 35–101, IAS/Park City Math. Ser., 7, Amer. Math. Soc., Providence, RI, 1999.

[HLS] H. Hofer, V. Lizan and J.-C. Sikorav, *On genericity for holomorphic curves in four-dimensional almost complex manifolds*, J. Geom. Anal. **7** (1997), 149–159.

[HWZ1] H. Hofer, K. Wysocki and E. Zehnder, *Properties of pseudoholomorphic curves in symplectisations. I. Asymptotics*, Ann. Inst. H. Poincaré Anal. Non Linéaire **13** (1996), 337–379.

[HWZ2] H. Hofer, K. Wysocki and E. Zehnder, *Properties of pseudo-holomorphic curves in symplectizations. II. Embedding controls and algebraic invariants*, Geom. Funct. Anal. **5** (1995), 270–328.

[HKM1] K. Honda, W. Kazez and G. Matić, *On the contact class in Heegaard Floer homology*, J. Differential Geom. **83** (2009), 289–311.

[HKM2] K. Honda, W. Kazez and G. Matić, *Contact structures, sutured Floer homology, and TQFT*, preprint 2008. `ArXiv:0807.2431`.

[Hu1] M. Hutchings, *An index inequality for embedded pseudoholomorphic curves in symplectizations*, J. Eur. Math. Soc. (JEMS) **4** (2002), 313–361.

[Hu2] M. Hutchings, *The embedded contact homology index revisited*, New perspectives and challenges in symplectic field theory, 263–297, CRM Proc. Lecture Notes, 49, AMS, 2009.

[HS1] M. Hutchings and M. Sullivan, *The periodic Floer homology of a Dehn twist*, Algebr. Geom. Topol. **5** (2005), 301–354.

[HS2] M. Hutchings and M. Sullivan, *Rounding corners of polygons and the embedded contact homology of* T^3, Geom. Topol. **10** (2006), 169–266 (electronic).

[HT1] M. Hutchings and C. Taubes, *Gluing pseudoholomorphic curves along branched covered cylinders I*, J. Symplectic Geom. **5** (2007), 43–137.

[HT2] M. Hutchings and C. Taubes, *Gluing pseudoholomorphic curves along branched covered cylinders II*, J. Symplectic Geom. **7** (2009), 29–133.

[HT3] M. Hutchings and C. Taubes, *Proof of the Arnold chord conjecture in three dimensions 1*, Math. Res. Lett. **18** (2011), 295-313.

[IP1] E. Ionel and T. Parker, *Relative Gromov-Witten invariants*, Ann. of Math. (2) **157** (2003), 45–96.

[IP2] E. Ionel and T. Parker, *The symplectic sum formula for Gromov-Witten invariants*, Ann. of Math. (2) **159** (2004), 935–1025.

[IS] S. Ivashkovich and V. Shevchishin, *Gromov compactness theorem for J-complex curves with boundary*, Internat. Math. Res. Notices **2000**, 1167-1206.

[KM] M. Kontsevich and Y. Manin, *Gromov-Witten classes, quantum cohomology, and enumerative geometry*, Commun. Math. Phys. **164** (1994), 525–562.

[KrM] P. Kronheimer and T. Mrowka, *Monopoles and three-manifolds*, New Mathematical Monographs, 10. Cambridge University Press, Cambridge, 2007.

[KLT1] C. Kutluhan, Y. Lee and C. Taubes, *HF=HM I: Heegaard Floer homology and Seiberg-Witten Floer homology*, preprint 2010. `ArXiv:1007.1979.`

[KLT2] C. Kutluhan, Y. Lee and C. Taubes, *HF=HM II: Reeb orbits and holomorphic curves for the ech/Heegaard-Floer correspondence*, preprint 2010. `ArXiv:1008.1595.`

[KLT3] C. Kutluhan, Y. Lee and C. Taubes, *HF=HM III: Holomorphic curves and the differential for the ech/Heegaard Floer correspondence*, preprint 2010. `ArXiv:1010.3456.`

[KLT4] C. Kutluhan, Y. Lee and C. Taubes, *HF=HM IV: The Seiberg-Witten Floer homology and ech correspondence*, preprint 2011. `ArXiv:1107.2297`.

[KLT5] C. Kutluhan, Y. Lee and C. Taubes, *HF=HM V: Seiberg-Witten-Floer homology and handle addition*, preprint 2012. `ArXiv:1204.0115`.

[LT] Y. Lee and C. Taubes, *Periodic Floer homology and Seiberg-Witten Floer cohomology*, preprint 2009. `ArXiv:0906.0383.`

[LR] A. Li and Y. Ruan, *Symplectic surgery and Gromov-Witten invariants of Calabi-Yau 3-folds*, Invent. Math. **145** (2001), 151–218.

[Li] R. Lipshitz, *A cylindrical reformulation of Heegaard Floer homology*, Geom. Topol. **10** (2006), 955–1097.

[M1] D. McDuff, *Singularities and positivity of intersections of J-holomorphic curves*. With an appendix by Gang Liu. Progr. Math., 117, Holomorphic curves in symplectic geometry, 191–215, Birkhäuser, Basel, 1994.

[M2] D. McDuff, *Comparing absolute and relative Gromov-Witten invariants*, preprint 2008.

[MS] D. McDuff and D. Salamon, *J-holomorphic curves and symplectic topology*, American Mathematical Society Colloquium Publications, 52. American Mathematical Society, Providence, RI, 2004.

[MW] M. Micallef and B. White, *The structure of branch points in minimal surfaces and in pseudo-holomorphic curves*, Ann. of Math. (2) **141** (1995), 35–85.

[OSz1] P. Ozsváth and Z. Szabó, *Holomorphic disks and topological invariants for closed three-manifolds*, Ann. of Math. (2) **159** (2004), 1027–1158.

[OSz2] P. Ozsváth and Z. Szabó, *Holomorphic disks and three-manifold invariants: properties and applications*, Ann. of Math. (2) **159** (2004), 1159–1245.

[Pa] B. Parker, *Exploded fibrations*, Proceedings of Gökova Geometry-Topology Conference 2006, 5290, Gökova Geometry/Topology Conference (GGT), Gökova, 2007.

[Ra] J. Rasmussen, *Floer homology and knot complements*, Ph.D. thesis, 2003. `ArXiv:math.GT/0306378.`

[Se1] P. Seidel, *A long exact sequence for symplectic Floer cohomology*, Topology **42** (2003), 1003–1063.

[Se2] P. Seidel, *Fukaya categories and Picard-Lefschetz theory.* Zürich Lectures in Advanced Mathematics. European MathematicalSociety (EMS), Zürich, 2008.

[Si] R. Siefring, *Relative asymptotic behavior of pseudoholomorphic half-cylinders*, Comm. Pure Appl. Math. **61** (2008), 1631-1684.

[T1] C. Taubes, *The Seiberg-Witten equations and the Weinstein conjecture*, Geom. Topol. **11** (2007), 2117–2202.

[T2] C. Taubes, *Embedded contact homology and Seiberg-Witten Floer cohomology I–V*, Geom. Topol. **14** (2010) 2497–3000.

[T3] C. Taubes, *The structure of pseudoholomorphic subvarieties for a degenerate almost complex structure and symplectic form on* $S^1 \times B^3$, Geom. Topol. **2** (1998), 221–332.

[U] I. Ustilovsky, *Infinitely many contact structures on* S^{4m+1}, Internat. Math. Res. Notices **14** (1999), 781–791.

[We1] C. Wendl, *Finite energy foliations on overtwisted contact manifolds*, Geom. Topol. **12** (2008) 531–616.

[We2] C. Wendl, *Finite energy foliations and surgery on transverse links*, Ph.D. thesis, 2005.

[We3] C. Wendl, *Automatic transversality and orbifolds of punctured holomorphic curves in dimension four*, preprint 2008. `ArXiv:0802.3842`.

[We4] C. Wendl, *Punctured holomorphic curves with boundary on* 3*-manifolds: Fredholm theory and embeddedness*, in preparation.

UNIVERSITÉ DE NANTES, 44322 NANTES, FRANCE
E-mail address: `Vincent.Colin@univ-nantes.fr`

UNIVERSITÉ DE NANTES, 44322 NANTES, FRANCE
E-mail address: `paolo.ghiggini@univ-nantes.fr`
URL: `http://www.math.sciences.univ-nantes.fr/˜Ghiggini`

UNIVERSITY OF SOUTHERN CALIFORNIA, LOS ANGELES, CA 90089
E-mail address: `khonda@usc.edu`
URL: `http://www-bcf.usc.edu/˜khonda`